

PAPER • OPEN ACCESS

Preface International Conference on Environmental, Energy and Earth Science (ICEEES)

To cite this article: 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1041** 011001

View the [article online](#) for updates and enhancements.

You may also like

- [Small and Medium Enterprises in Pekanbaru: Implementing an Integrated Work Program](#)
Hardi
- [Analyzing the critical locations in response of constructed and planned dams on the Mekong River Basin for environmental integrity](#)
Yuan Gao, Shiblu Sarker, Tanni Sarker et al.
- [Nutrient testing of organic liquid fertilizer in tempeh factory in Pekanbaru City](#)
Neng Susi, Vonny Indah Sari, Indra Purnama et al.

ECS Toyota Young Investigator Fellowship



For young professionals and scholars pursuing research in batteries, fuel cells and hydrogen, and future sustainable technologies.

At least one \$50,000 fellowship is available annually.
More than \$1.4 million awarded since 2015!



Application deadline: January 31, 2023

Learn more. Apply today!

Preface International Conference on Environmental, Energy and Earth Science (ICEEES)

Universitas Lancang Kuning Pekanbaru organized with **the International Conference on Environmental, Energy and Earth Science (ICEEES)** on September 22, 2021 in Pekanbaru, Indonesia. The conference is aims to exchange knowledge and research finding among academicians, researchers, professionals, policy makers, and postgraduate students.

The Awareness to increase the number of publications on the results of research that has been done, deserves appreciation by all parties. Because of that, Universitas Lancang Kuning Pekanbaru provide motivation and space for researcher to disseminate their research and accommodate the result of research that has been done. **The International Conference on Environmental, Energy and Earth Science (ICEEES)**, is another International Conference held by Universitas Lancang Kuning Pekanbaru.

The International Conference on Environmental, Energy and Earth Science (ICEEES) was chosen to be implemented virtually, this is because the cov-19 pandemic is still spreading. The conference was perform using zoom. **The International Conference on Environmental, Energy and Earth Science (ICEEES)** event is virtually implemented with a model that all invited speakers are given time to present their material for 30 minutes every invited speaker and after that a question and answer is carried out with the participants with a direct questioning system, through chat forums and Q&A forums provided by the zoom application. Overall, the conference took 5 hours the number of participants who joined the zoom room was recorded at participants. Participants came from few countries, namely Indonesia, Malaysia, India, Egypt and Australia.

The International Conference on Environmental, Energy and Earth Science (ICEEES) is implemented with the support of a stable internet network system and a zoom application. In the implementation there were several technical obstacles encountered by the participants, namely the difficulty of joining the zoom application due to the unstable internet signal. The holding of a virtual conference felt less meaningful, due to the lack of interaction between speakers and participants

The International Conference on Environmental, Energy and Earth Science (ICEEES) committee received 124 manuscripts and a total of 117 papers were presented and discussed. The papers were authored by researchers from Indonesia, Malaysia, India, Egypt, japan and Australia.

All papers have been scrutinized by a panel of reviewers who provide critical comments and corrections, and thereafter contributed to the improvement of the quality of the papers. Based on the reviewer's reports, 92 papers were selected and eligible to be published in the proceeding.

We sincerely express our gratitude to the international/national advisory committee, presenters, organizing committee members, session chairs, all members of organization, participants, contributors and all the members of **The International Conference on Environmental, Energy and Earth Science (ICEEES)**. Last but not the least, we are thankful to IOP EES Conference Series for producing the proceeding.



Organizing Committee

General Chair :

Dr. Anto Ariyanto, M.Si (Universitas Lancang Kuning, Indonesia)

General Co-Chair:

Dr. H. Jeni Wardi, S.E., M.Ak., Ak., CA (Universitas Lancang Kuning, Indonesia)

Secretary :

Dr. David Setiawan, M.T (Universitas Lancang Kuning, Indonesia)

Treasurer and Finance Chair :

Ambar Tri Ratnaningsih, S.Hut., M.Si (Universitas Lancang Kuning, Indonesia)

Publication Chair :

Robbi Rahim, S.Kom., M.Kom (STIM Sukma Medan, Indonesia)

Website and Social Media Chair :

Guntoro, S.T., M.Kom (Universitas Lancang Kuning, Indonesia)

Febrizal Alfarasy Syam, S.Kom., M.Kom (Universitas Lancang Kuning, Indonesia)

Ismar Husnudin, S.T (Universitas Lancang Kuning, Indonesia)

Editors

1. Prof. Dr. Joran (Linkoping University, Swedia)
2. Prof. Dr. Amran bin Hamzah (Universitas Teknologi Malaysia, Malaysia)
3. Prof. Dr. Khaerurijal (Institut Teknologi Bandung, Indonesia)
4. Prof. Dr. Asep Saefuddin (Institut Pertanian Bogor, Indonesia)
5. Prof. Dr. Mauridi (Institut Teknologi Sepuluh November, Indonesia)
6. Prof. Dr. Adi Soeprijanto (Institut Teknologi Sepuluh November, Indonesia)
7. Prof. Dr. Syafrani (Universitas Lancang Kuning, Indonesia)
8. Prof. Dr. Sudi Fahmi (Universitas Lancang, Kuning, Indonesia)
9. Prof. Dr. Hariadi Kartodiharjo (Institut Pertanian Bogor, Indonesia)
10. Prof. Dr. Sumbangan Baja (Universitas Hasanuddin, Indonesia)
11. Prof. Wiendu Nuryanti Ph.D (Universitas Gajah Mada, Indonesia)
12. Prof. Dr. Chairul Saleh (Universitas Islam Indonesia, Indonesia)
13. Prof. Dr. Richardus Eko Indrajit (STMIK Perbanas, Indonesia)

Organizing Institutions

- Universitas Lancang Kuning

PAPER • OPEN ACCESS

Peer Review Statement

To cite this article: 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1041** 011002

View the [article online](#) for updates and enhancements.

You may also like

- [Peer review declaration](#)
- [Peer review declaration](#)
- [Peer Review Declaration Form Guidance](#)

ECS Toyota Young Investigator Fellowship



For young professionals and scholars pursuing research in batteries, fuel cells and hydrogen, and future sustainable technologies.

At least one \$50,000 fellowship is available annually.
More than \$1.4 million awarded since 2015!



Application deadline: January 31, 2023

Learn more. Apply today!

Peer Review Statement

All papers published in this volume have been reviewed through processes administered by the Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

- **Type of peer review:** Single Anonymous
- **Conference submission management system:** <https://iceees.unilak.ac.id/paper-submission/>
- **Number of submissions received:** 129
- **Number of submissions sent for review:** 129
- **Number of submissions accepted:** 87
- **Acceptance Rate (Submissions Accepted / Submissions Received × 100):** 67.4
- **Average number of reviews per paper:** 2
- **Total number of reviewers involved:** 15
- **Contact person for queries:**
Name: Robbi Rahim
Email: robbirahim@ieee.org
Affiliation: Sekolah Tinggi Ilmu Manajemen Sukma



Table of contents

Volume 1041

2022

◀ Previous issue Next issue ▶

International Conference on Environmental, Energy and Earth Science 22/09/2021 - 23/09/2021

Online

Accepted papers received: 27 May 2022

Published online: 23 June 2022

Open all abstracts

Preface

OPEN ACCESS 011001

Preface International Conference on Environmental, Energy and Earth Science (ICEEES)

+ Open abstract  View article  PDF

OPEN ACCESS 011002

Peer Review Statement

+ Open abstract  View article  PDF

Agriculture

OPEN ACCESS 012001

The effect of chitosan addition on physicochemical properties, proximate and antioxidant of functional drinks of red and purple roselle (*Hibiscus sabdariffa* L.) extracts

Kusnadi, Aldi Budi Riyanta and Rizki Febriyanti

+ Open abstract  View article  PDF

OPEN ACCESS 012002

The characteristics of bio-oil hydrothermal pyrolysis empty fruit bunches based on powder size and hydrogen gas pressure

Rina Novia Yanti, Erliza Hambali, Gustan Pari and Ani Suryani

+ Open abstract  View article  PDF

OPEN ACCESS 012003

The effect of proportion of ganyong starch and waste of straw rice on biodegradable foam production as sustainable packaging

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Gresita Fitaloka Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012004

Physicochemical and antioxidant characteristics of black garlic from indigenous Indonesian garlic

K S Sasmitaloka, SM Widayanti, I Mulyawanti and ES Iriani

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012005

Efficiency analysis of vannamei shrimp supply chain performance in PT Sari Indonesia Group, Suak Pandan Village, West Aceh Regency, Aceh Province

Yoga Nugroho, Teuku Athaillah and Muhibbul Qadri

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012006

Nutrient testing of organic liquid fertilizer in tempeh factory in Pekanbaru City

Neng Susi, Vonny Indah Sari, Indra Purnama and Hazra Yuvendus

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012007

Extraction of catfish bone waste (*pangasius hypophtalmus*) by utilizing an organic liquid of pineapple peel waste (*ananas comosus*) into gelatin

Ira Oktaviani Rz, Rahimatul Uthia, Fathul Jannah and Alexsander Yandra

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012008

The effect of orange peel waste addition on ZnO characteristics and its ability to degrade methyl orange

Aulia Dewi Rosanti, Fahmi Hidayat, Yuly Kusumawati, Arif Fadlan, Anggita R.K Wardani and Eva Umi Latifah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012009

Carbon sequestration of city agriculture: between farming and non-farming land

D.R.D. Hastuti, R. Darma, D. Salman, S. Santoso and A. Rahim

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012010

Comparison predicted using nonlinear model and in vivo studies growth production of broiler after fed different levels of fermented tamarind flour as replacing soya bean meal

Onsijofan, D.N. Adlis, MBH Natsim, Y.F. Nuringtiasari and A.D. Radianto

Use of cookies. To find out more, see our Privacy and Cookies policy.



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012011

Dry matter (DM), crude protein (CP) and water consumption in sheep which supplemented from *Moringa* leaves meal and palm oil in elephant silage-based feed

S Rizqiana, W Suryapratama and F M Suhartati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012012

Effect of various packaging materials for storing ground yellow corn of hybrid C-1 variety on water and amyllum content

Munasik, N Nurbaety, N Hidayat and E Susanti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012013

Exploring the herbaceous plant height and dry matter relationship on a sub-humid rangeland in Zimbabwe

T A Zezai, P H Mugabe and V E Imbayarwo-Chikosi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012014

Potential development based on the agricultural waste in the eastern region of West Java Province

F. D. Perwitasari, A. R. S. Putra, B. Suwignyo and R. Widiati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012015

Farming in Kediri Indonesia: analysis of cluster k-means

N Solikin, B Hartono, Sugiono and Linawati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012016

The Identification and Role pf Mold in Lowering the HCN Levels in Black Cassava Based on the Immersion Duration

Mery Pattipeilohy and Alamanda Pelamonia

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012017

The Effects of Some Additives On Characteristics RVA Profile of Cassava Flour

H Herawati and E. Kamsiati

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Agriculture Engineering and Technology

OPEN ACCESS

012018

A comprehensive review on power tillers usage – bottlenecks and prospects in India

Krishnadas R and Renganathan R

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012019

Android-based feasibility study transformation analysis case study of the RASI development in Cirendeu-Kampung Cimahi

H Herawati, A S Soemantri, E Kamsiati and M. Bachtiar

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012020

Water pump control system in hydroponic plants using the ebb and flow method

David Setiawan, Hamzah, Latifa Siswati, Anto Ariyanto and Guntoro

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012021

Resetting the ground fault relay on a 28 MVA power transformer due to grounding method retrofit

Hamzah Eteruddin, Arlenny, Hadi Siswanto, Yanuar Z. Arief and Fathimah Hasanti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012022

Intelligent ubiquitous technology as a precision agri-food framework: a proposed framework

T A Cahyanto, S Wahjuni, H Sukoco, H Rahmawan and S N Neyman

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012023

Development of internet of things-based monitoring system for application solar energy technology in Bunaken island

M Rumbayan, I Pundoko, D Ruindungan and N V Panjaitan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012024

Technical analysis on household energy consumption: LMDI decomposition index and innovative

Yetty Murni, Eka sudarmaji, Sri Ambarwati and Ismiriati Nasip

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012025

Optimization of electrical energy in the fan using fuzzy logic controllers

Arlenny and David Setiawan

[+ Open abstract](#) [View article](#) [PDF](#)

Animal Biodiversity

OPEN ACCESS

012026

The effect of giving chia seeds on the bodyweight of pregnant mice (*Mus Musculus L.*)

R Uthia and I O Rz

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012027

Supplementing mineral selenium and vitamin e in diets on *in vivo digestibility*, blood glucose, and urea levels of cows

D. Pamungkas, A.S. Putri, F. Firdaus, R. Widiyawati and D.M. Dikman

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012028

Effect of *Saccharomyces cerevisiae* on Probiotic Properties of Goat Milk Kefir

L E Radiati, D L Hati, D Fardiaz and L R H Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012029

Cattle production system in semi-arid area of Timor Island

Y Ngonjo, S Ratnawaty and PR Matitaputty

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012030

Sustainable indigenous cattle production in Brebes Regency, Indonesia: opportunities and threats during pandemic

M Sugiarto, Y. N Wakhidati, O. E. Djatmiko, S. Nur and A Einstein

[+ Open abstract](#) [View article](#) [PDF](#)

Disaster Management

OPEN ACCESS

012031

The role of philanthropy in reducing impact of disaster on economies of ASEAN-9

M S A Majid, R Nurdin, T Azhar and S Sartiyah

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS 012032
Community participation in disaster management in Gondoriyo village administration, Ngaliyan sub-district, Semarang city
Supratiwi, Yuwanto and Kushandajani
[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012033
Assessing and preparedness for earthquake disaster in Salaf-Khalaf Islamic Boarding School
Fatiya Rosyida, Neni Wahyuningtyas, Ardyanto Tanjung and Abdul Kodir
[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012034
Disaster education in disaster-prone schools: a systematic review
R Yusuf, Razali, Sanusi, Maimun, I Fajri and S A Gani
[+ Open abstract](#) [View article](#) [PDF](#)

Environmental Issue and challenge

OPEN ACCESS 012035
LCA methodology for detecting environmental impacts on natural gas drilling process
Masayu Rosyidah, Andiyan Andiyan, Haniek Listyorini, Pranoto Hadi Prayitno, Yuswardi Yuswardi and Yuhanah Yuhanah
[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012036
Community based ecotourism management to strengthen environmental ethics and supports sustainable development in Pelalawan district, Indonesia
Nawari, Thamrin, Nofrizal, Almasdi Syahza, Juandi Muhammad and Nur Islami
[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012037
Community forestry for environmental sustainability and ecotourism: the context and problems in Indonesia
R Sulistiowati, S Wahyuni, M K Yunanto, M Elsera, W E Yudiatmaja and T Samnuzulsari
[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012038
Study on the commitment of oil palm companies to achieve sustainable agriculture in Riau Province from the perspective of pesticide use
S Syafrani, I Purnama, A Mutamima and W N Dewi
[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



OPEN ACCESS

012039

Assessing the Feasibility of Migration Policy from LPG Stoves to Induction Stoves in Indonesia

Tri Anggono, Iyung Ruslan, Chrisnawan Anditya, Dian Galuh Cendrawati and Muhammad Indra al Irsyad

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012040

The innovation of ecological fiscal transfers policy at Siak Regency

Muhammad Ichsan Kabullah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012041

Farmers' perceptions of the performance of agricultural instructors in mangosteen farmers' groups in Segati Village, Langgam District, Pelalawan Regency

Asgami Putri, Latifa Siswati and Hamdan Yasid

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012042

Environment Conservation Responsibility in Community Development

N Sutisna, H Herdiansyah and P Sanjatmiko

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012043

Environmental Conservation in Ciletuh Geopark, West Java

R A Zuvara, H Herdiansyah and D Asteria

[+ Open abstract](#) [View article](#) [PDF](#)

Food Science and Technology

OPEN ACCESS

012044

Effectiveness of silica gel from palm kernel shell ash as a moisture absorber of bottle packaging medicine

Lidya Novita and Iswadi Idris

[+ Open abstract](#) [View article](#) [PDF](#)

Land Use Management

OPEN ACCESS

012045

Comparative study of village community participation in peat restoration in the Peat Swamp Forest Management Unit (PSFMU) Tebing Tinggi Island, Indonesia

Elhi Malik, Agus Iko Skripriyana, Sindangdji, Jc. Ws Satriyasa, Igr Witayonarko and M. Nasirul find out more, see our Privacy and Cookies policy.



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012046

Swot analysis of utilization of palm oil mill effluent to improve soil quality and crop productivity (case study at PT X, Lampung, Indonesia)

Dela Febriana, Sylvia Madusari and Vira Irma Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012047

Groundwater level fluctuations in peatlands as responses to canal block construction in Sungai Tohor, Tebing Tinggi Island, Indonesia

H. Malik, Ig. L. S. Purnama, Sudarmadji, L. W. Santosa and M. Naufal

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012048

Identification of land use and land cover using the image Landsat 8 in upstream Lematang sub-watershed by support vector machine and random trees methods

A Dinata, F Dhiniati and L E Diansari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012049

Conservation and utilization on Sawahlunto ex mining area for cultural landscape

Rika Cheris and Amanda Rosetia

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012050

Measuring cross-sector collaboration an effective solution for solving unlicensed mining problems in Kuantan Singingi, Indonesia

MY Tiyas Tinov, Aras Mulyadi, Sujianto, Zulkarnain, Ardi Putra, Tika Mutia and Tito Handoko

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012051

Helminthiasis of rabbits on the upland and lowland areas and the risk factors

D Indrasanti, M Indradji, Sufiriyanto, M Samsi and E Yuwono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012052

Water Quality Analysis Based on Land Use in Sekampung River, Lampung, Indonesia

K Fitriani, T K Nufutomo and R Putra

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



Oil Plantation and Management

OPEN ACCESS

012053

Application of oil palm empty bunches as organic mulch in oil palm plantation (*Elaeis guineensis* Jacq.): an evaluation and SWOT analysis

Lala Novita Sari, Sylvia Madusari and Vira Irma Sari

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012054

Oil palm trunk replanting as brown sugar raw materials

Latifa Siswati, Enny Insusanty, Neng Susi and Nopryanti

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012055

Prediction of palm oil production in Riau Province using the single exponential smoothing method

Guntoro, Lisnawita, Zamzami and David Setiawan

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012056

Study of biogas utilization as waste-to-energy plant and transport modelling of iron (Fe), lead (Pb) and copper (CU) in leachate at Muara Fajar landfill Pekanbaru

H Yuvendius, E Zondra, Zainuri and V I Sari

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012057

Utilization of oil palm waste as animal feed for Bali cattle in Margomulyo Village, Central Bengkulu

E Ramon, Z Efendi, H Kusnadi, S Yuliasari, A Ishak and WA Wulandari

[+ Open abstract](#)[View article](#)[PDF](#)

Plant Biodiversity

OPEN ACCESS

012058

Etnobotany of medicine plants in the community of Rawang Kao Village, Siak, Indonesia

Sri Wahyuni and Mar'atul Afidah

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012059

Can mangrove intensive policies in Indonesia benefit more coastal communities? environmental policy analysis

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012060

Validation of UV-VIS spectrophotometric method for the determination of Calcium in Maman (cleome gynandra l)

Fathul Jannah, S Meri Agritubella, Ira Oktaviana and Lily Restusari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012061

Utilization of bamboo charcoal to decrease tannin content of lindur fruit flour (Bruguiera gymnorrhiza)

Niken Pujirahayu, Zakiah Uslinawaty, Nuhayati Hadjar and Muhtadir

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012062

Supplementing safflower oil (*Carthamus Tinctori L*) and inocytol in feed and its effect

Ning Iriyanti, Ismoyowati, Efka Aris Rimbawanto and Modawy Abdelgader Albasheer Altayb

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012063

The effects of nano encapsulated *Syzygium Polyanthum* leaves extract inclusion in drinking water

A N 'Afifah, Zuprizal and N D Dono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012064

Effects of plant extracts against gastrointestinal nematodes of livestock: a meta-analysis

A Jayanegara, D M Sinaga and E B Laconi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012065

Coconut oil effects on in vitro ruminal fermentation and methanogenesis

A Cusiayuni, RK Nurfatahillah, D Evvyernie, A Jayanegara and KG Wiryawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012066

Effect of palm oil supplementation level on in vitro ruminal fermentability

RK Nurfatahillah, A Cusiayuni, A Jayanegara, KG Wiryawan and D Evvyernie

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

012066

The Composition of Goat Milk Kefir Added with *Moringa oleifera* Leaf Powder

S R N Endah, P D Wulasari, A Nofriyadi, Nurliyani and E Harmayani

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012068

The effect of safflower oil (*Carthamus Tinctorius* L.) and inositol supplementation on egg production

Ismoyowati, N Iriyanti and M Abdelgader

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012069

The Anti-Inflammatory Activity of Cherry Leaf Extract (*Muntingia Calabura* L.) Balm Stick

Fitria Nugrahaeni, Kriana Efendi and Abdul Kholik Aziz

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012070

The Effect of HPMC Concentration as a Gelling Agent on Color Stability of Copigmented Blush Gel Extract of Purple Sweet (*Ipomoea Batatas* (L.) Lam.)

Fitria Nugrahaeni, Nining and Redina Okvianida

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012071

Sunscreen Factor Formulation and Test of Gel Preparations of 70% Ethanol Extract on Arabica Coffee Leaf (*Coffea arabica* L.)

Sofia Fatmawati, Fitria Nugrahaeni, Fith Khaira Nursal and Ana Fitriana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012072

Antioxidant Activity and Sun Protection Factor (SPF) Graded Extract of Katuk Leaves (*Sauropus androgynus* (L.) Merr.)

Sofia Fatmawati, Ni Putu Ermi Hikmawanti, Akbar Fadillah and Awlia Mustia Putri

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012073

Anti-Inflammatory Activity of Water Extract of *Talinum paniculatum*. (Jact). Gaertn Leaves on Wistar Rat

A Emelda, Sukmawati, A L Dongke and I Marzuki

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Sustainable Food Systems

OPEN ACCESS

012074

Assessing food safety compliance for food SMEs in Indonesia

Tri Fajarwaty and David Jukes

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012075

In-silico screening of inhibitor on protein epidermal growth factor receptor (EGFR)

Jeremi Ongko, Jesica Viona Setiawan, Alfina Gracia Feronytha, Agnes Juliana, Andy Effraim, Mariana Wahjudi and Yulanda Antonius

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012076

Effect of Various Feed Additives on Carcass and Meat Quality of Two Different Strains of Chickens

E Tugiyanti and E Susanti

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012077

Microbiological profile of concentrated yoghurt manufactured from low and full fat milk during storage

T Setyawardani, J Sumarmono and D R Kusuma

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012078

Effect of dietary supplementation with flying fish (exocoetidae) silage on the egg quality and duck performance

L Ambarwati and S P Syah

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012079

Production profile and microbiological properties of locally produced pork sausage in the Philippines

MM Dealino and CM Bueno

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012080

Farming Sustainability: A Sensitivity Analysis

Y N Wakhidati, M Sugiarto, H Aunurrohman, S Mastuti and J A Suryani

[+ Open abstract](#)[View article](#)[PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



-
- OPEN ACCESS** 012081
Improving genetic quality of local livestock for achieving food security
E Kurnianto
[+ Open abstract](#) [View article](#) [PDF](#)

-
- OPEN ACCESS** 012082
Current goat milk production, characteristics, and utilization in Indonesia
J Sumarmono
[+ Open abstract](#) [View article](#) [PDF](#)

Waste Management

-
- OPEN ACCESS** 012083
The impact of agricultural waste on river water quality of kreo watershed in Semarang city
W Setyaningsih and R S Sanjaya
[+ Open abstract](#) [View article](#) [PDF](#)

-
- OPEN ACCESS** 012084
Utilization of plastic waste as an eco-friendly construction material
Zainuri, G Yanti and S W Megasari
[+ Open abstract](#) [View article](#) [PDF](#)

-
- OPEN ACCESS** 012085
Biogas as methane mitigation path on dairy cattle waste management
A Herliatika and Y Widiawati
[+ Open abstract](#) [View article](#) [PDF](#)

Water, Climate and Emission Science

-
- OPEN ACCESS** 012086
Does rebound effect influence the factors of carbon emission in Indonesia? Kaya index and LMDI decomposition
Eka Sudarmaji, Noer Azam Achسانی, Yandra Arkeman and Idqan Fahmi
[+ Open abstract](#) [View article](#) [PDF](#)

-
- OPEN ACCESS** 012087
Study of phytoplankton biology index and water quality parameters of kali Surabaya River
O H Cahyonugroho, S Hariyanto and G Supriyanto
[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012088

Factor and perception analysis of the impact of boat parking on river pollution

B Rahman and M S Aryanto

[+ Open abstract](#)[View article](#)[PDF](#)

JOURNAL LINKS

[Journal home](#)[Journal scope](#)[Information for organizers](#)[Information for authors](#)[Contact us](#)[Reprint services from Curran Associates](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



PAPER • OPEN ACCESS

Antioxidant Activity and Sun Protection Factor (SPF) Graded Extract of Katuk Leaves (*Sauropus androgynus* (L.) Merr.)

To cite this article: Sofia Fatmawati *et al* 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1041** 012072

View the [article online](#) for updates and enhancements.

You may also like

- [Phytochemical Screening and Antioxidant Activity of Ethanol Extract of Leilem \(*Clerodendrum minahassae* Teijsm. & Binn\) as an Antihyperlipidemic and Antiatherosclerotic Agent](#)
C F Kairupan, F R Mantiri and R R H Rumende
- [Comparison of Antibacterial Activity in Ethanol Extract and Essential Oil of *Citrus sinensis* \(L.\) Peels Obtained by Soxhlet and Distillation Methods](#)
E Kusri, DP Mawarni, M Mamat et al.
- [Evaluation and Determination of Total Antioxidant in Anting-Anting \(*Acalypha indica* L.\) Leaf Extract](#)
Refilda, F Ilahi, D Hanifa et al.



The Electrochemical Society
Advancing solid state & electrochemical science & technology

242nd ECS Meeting

Oct 9 – 13, 2022 • Atlanta, GA, US

Early hotel & registration pricing
ends September 12

Presenting more than 2,400
technical abstracts in 50 symposia

The meeting for industry & researchers in

BATTERIES
ENERGY TECHNOLOGY
SENSORS AND MORE!



Register now!



ECS Plenary Lecture featuring
M. Stanley Whittingham,
Binghamton University
Nobel Laureate –
2019 Nobel Prize in Chemistry



Antioxidant Activity and Sun Protection Factor (SPF) Graded Extract of Katuk Leaves (*Sauropus androgynus* (L.) Merr.)

Sofia Fatmawati*, Ni Putu Ermi Hikmawanti, Akbar Fadillah, Awlia Mustia Putri

Universitas Muhammadiyah Prof. DR. HAMKA, Jakarta, Indonesia

*sofia.fatmawati@uhamka.ac.id

Abstract. Katuk leaf (*Sauropus androgynus* (L.) Merr.) is believed to have medicinal properties, one of which is as an antioxidant. Its efficacy as an antioxidant cannot be separated from the phenolic and flavonoid compounds contained in katuk leaves. This study aims to determine the total phenolic and flavonoid levels as well as the antioxidant activity of graded extracts, namely *n*-hexane, ethyl acetate, and 70% ethanol extract of katuk leaves. The results showed that the total phenolic content and total flavonoid content of each extract were significantly different. Antioxidant activity by the DPPH method was calculated at IC₅₀ using quercetin as a comparison. The highest antioxidant potential was shown in the ethanol extract. Each extract has the potential as a sunscreen and ethanol extract provides the highest SPF value.

1. Introduction

Phenol compounds are characterized by the presence of an aromatic ring and one or two hydroxyl groups. Phenol compounds that have more than two hydroxyl groups are called polyphenols, for example the tannin, flavonoid, melanin, and lignin groups [1]. Flavonoid compounds have a characteristic structure of two aromatic rings connected by three C atoms, usually with O atomic bonds in the form of heterocyclic oxygen bonds [1]. Flavonoids are a group of secondary metabolites produced by plants which are included in the large group of polyphenols. Flavonoids have the ability to scavenge free radicals and inhibit lipid oxidation [2].

Sunscreen contains compounds that can protect the skin by absorbing ultraviolet (UV) rays emitted by the sun [3]. Compounds contained in sunscreen can be used to prevent various skin diseases and to protect human skin health from the negative effects of UV rays. Sunscreens are divided into 2 groups, namely physical sunscreens and chemical sunscreens. Chemical sunscreens are sunscreens that absorb ultraviolet light, such as PABA, PABA esters, benzophenone, avobenzone, salicylate, cinnamate and camphor derivatives [4].

Katuk leaves (*Sauropus androgynus* (L.) Merr.) contain secondary metabolites such as tannins, saponins, alkaloids, flavonoids, glycosides, and phenols [5]. The 95% ethanol extract of katuk leaves has a high content of phenolic compounds with 1.49 mgGAE/g fresh weight and 8.71 mgGAE/g dry weight [6]. Antioxidant activity in previous studies stated that the IC₅₀ value of katuk leaf methanol extract had a value of 80.81 ppm [7]. Variations in ethanol concentration and extraction methods resulted in different phenolic and flavonoid content in katuk leaves [8], [9]. Multilevel extraction gives better results than individual extraction with each solvent [10].



The aim of the study was to determine the levels of phenolic, flavonoid, as well as antioxidant activity and SPF testing of the *n*-hexane, ethyl acetate, and 70% ethanol extract of katuk leaves from the extraction process with graded maceration, using UV-Vis spectrophotometry method.

2. Materials and Methods

2.1 Preparation Extract

Katuk (*Sauropus androgynus* (L) Merr) leaves were collected and determined at the Institut Pertanian, Bogor, West Java. Simplicia processing begins with fresh katuk leaves taken and cleaned of impurities, then washed with water until clean, drained and finely chopped, and then dried again by aerating. After dried then powdered.

The graded extract was obtained by a stepwise method using solvents with different polarity levels, namely *n*-hexane, ethyl acetate, and 70% ethanol (1:10). The material is soaked for 24 hours, stirring occasionally for the first 6 hours. The residue is then separated from the filtrate. The process is repeated until the solvent is clear (3 repetitions). The residue was then extracted with ethyl acetate and 70% ethanol using the same procedure. Each maserate obtained was then concentrated with a vacuum rotary evaporator at a temperature of 40°C - 50°C to obtain a thick extract.

2.2 Characterization of Extract

The organoleptic test was carried out by observing the shape, color, smell, and taste using the five senses. Each extract was weighed carefully as much as 2 grams, then put into a silicate crucible that had been sized and compared, incandescent in a kiln and the temperature was gradually increased to 600°C (temperature difference of approximately 25°C) until carbon-free. Next, it was cooled in a desiccator and then weighed. The total ash content is calculated against the weight of the test material and is expressed in % w/w. Each 1.0 gram extract was placed in a moisture content balance container which had been previously sized and the initial weight of the extract was seen. Then the extract was dried at a temperature of 105°C at the moisture content balance until a constant extract weight value was obtained, afterward the final weight of the extract was seen. Results are viewed as % MC values [11].

2.3 Phytochemical Screening of Extract

The phytochemical compounds of kencur (aromatic ginger-*Kaempferia galanga*) ethanolic extract, such as phenolic, flavonoid, tannin, alkaloid, and terpene were qualitatively identified following standard procedures describing in the Harborne and Indonesian Herb Pharmacopoeia [12].

2.4 Determination of Total Phenolic Level

From the base solution of 100 ppm gallic acid, several concentrations were made, namely 18 ppm, 30 ppm, 42 ppm, 54 ppm, and 66 ppm. From each concentration of the standard solution of gallic acid, 300 µl was pipetted and then 1.5 ml of *Folin-Ciocalteu* reagent (1:10) was added. After being allowed to stand for 3 minutes, each solution was added with 1.2 ml of 7.5% Na₂CO₃ solution, shaken homogeneously, and allowed to stand in the operating time range at room temperature. All solutions were measured for absorbance at the maximum absorbance wavelength obtained, which was 756.5 nm, then a calibration curve was made for the relationship between gallic acid concentrations. The extract solution obtained was pipette 300 l and added 1.5 ml of *Folin-Ciocalteu* reagent and shaken. It was allowed to stand for 3 minutes, added 1.2 ml of 7.5% Na₂CO₃ solution and allowed to stand again in the operating time range at room temperature. The absorbance of the extract solution was measured by UV-Vis spectrophotometer at the maximum absorbance wavelength. The absorbance of the extract solution was measured by UV-Vis spectrophotometer at a wavelength of 756.5 nm. Performed 3 repetitions [13].

2.5 Determination of Total Flavonoid Level

From 1000 ppm quercetin mother liquor, several concentrations were made, namely 33, 57, 81 and 105 ppm. A total of 0.5 ml of quercetin solution was made with several concentrations of pipette then added with 1.5 ml of methanol and added 0.1 ml of 10% AlCl₃ reagent 0.1 ml of sodium acetate (1M) and 2.8 ml of distilled water. Then the solution was shaken and left for an operating time of 60 minutes at room temperature. Measure the absorbance at a wavelength of 434 nm against the standard. Each concentration of 0.5 ml of the test solution was added with 1.5 ml of methanol and added 0.1 ml of 10% AlCl₃ reagent 0.1 ml of sodium acetate (1M) and 2.8 ml of distilled water, then allowed to react during the operating time at room temperature. Measure the absorption at the maximum wavelength against the standard [14].

2.6 Antioxidant Activity Test

Weigh 100.0 mg of ethanol extract of katuk leaves from each extraction time dissolved with methanol in a volumetric flask, then diluted to obtain concentrations of 20, 40, 60, 80 and 100 ppm. A total of 0.2 mL of sample solution of each concentration was added with 1 mL of 0.5 mM DPPH and 5 mL of methanol, allowed to stand according to the operating time for 30 minutes and then measured at a maximum wavelength of 515.5 nm. The percentage of radical scavenging activity was calculated by the formula [15].

2.7 Sunscreen Potency Test

The ethanol extract of Arabica coffee leaves was diluted with ethanol to a concentration of 100 ppm. The SPF value was determined by measuring the absorbance of the solution from the extract using a UV-Vis spectrophotometer at a wavelength of 290-320 nm. The sample was dissolved in 5 mL of ethanol p.a. The determination of the SPF value was carried out three times for each sample. The absorbance data obtained were processed by the equation of Mansur (1986) [16].

$$\text{SPF} = \text{CF} \times \text{EE} \times \text{I}(\lambda) \times \text{abs}(\lambda)$$

The SPF value can be calculated by multiplying the correction factor (CF), the erythema effect spectrum (EE), the intensity spectrum from the sun (I), and also the absorbance (Abs) of the cream sample of coffee leaf ethanol extract. The SPF assessment refers to the provisions of the FDA (Food and Drug Administration), namely minimal protection if SPF 2-4, moderate protection 4-6, extra protection 6-8, maximum protection 8-15, and ultra protection >15 [17].

2.8 Data Analysis

The observed data from the dispersion test and color stability test were then analyzed using one-way ANOVA. If there is a significant difference between the formulas, then it is continued with the Tukey HSD test with a 95% confidence level to see if the formula has a significant difference or not.

3. Results and Discussion

The purpose of using three solvents with different polarities is to obtain active compounds from katuk leaves based on their polarity [10]. In this study, repetition was carried out with the aim of providing sufficient extract weight and being able to attract residual compounds that were still left in the previous iteration. Extraction using solvents with different polarities will produce different polyphenol components so that the antioxidant properties or characteristics of each compound obtained from the extraction are also different [18].

Table 1. Katuk Leaf Extraction Results

Extract type	Average extract weight (g) ± SD	Average extract yield (%) ± SD
<i>n</i> -Hexane	18,05 ± 0,73	7,22 ± 0,29
Ethyl acetate	14,87 ± 1,15	5,95 ± 0,46
Ethanol 70 %	50,27 ± 1,90	20,11 ± 0,76

Table 2. Characteristics of Katuk Leaf Extract

Parameters	Extract type					
	<i>n</i> -Hexane		Ethyl acetate		Ethanol 70%	
Organoleptic						
Form	Thick	extract	Thick	extract	Thick	extract
Smell	Typical		Typical		Typical	
Favor	Bitter		Bitter		Rather	Bitter
Color	Green		Green		Chocolate	
Ash Level	2,59 % ± 0,22		0,55 % ± 0,05		6,88 % ± 0,93	
Drying Shrinkage	1,08 % ± 0,48		3,04 ± 0,58		3,48 ± 2,41	

The yield (Table 1) of the extract obtained from the multilevel extraction resulted in the lowest yield of the ethyl acetate extract, followed by the *n*-hexane extract. The yield obtained in the multistage extraction with 70% ethanol showed the highest yield. Based on the principle of mass transfer of solute into a solvent that is in accordance with characteristics *like dissolves like*, where there is a transfer at the interface layer (solvent and solute) then the solute diffuses into the solvent. The results are different because the ability to attract compounds in each solvent is different too. 70% ethanol extract has the highest yield because 70% ethanol solvent is universal. It can attract non-polar and polar compounds so that 70% ethanol extract is attracted to more secondary metabolites [8].

Organoleptic test on leaf extract was conducted to determine the characteristics of shape, smell, taste, and color. Of the three katuk leaf extracts the difference was only in the extract, the *n*-hexane, and ethyl acetate extracts had a green color while the 70% ethanol had a brown color.

The ash content obtained from this research is *n*-hexane 2.59%, ethyl acetate 0.55%, and ethanol 70% 6.88% as seen in (Table 2). The three extracts had simplicia criteria that were in accordance with the guidelines for a good extract ash content, namely the ash content of not more than 12%. The purpose of ash content is to show the content of organic and inorganic compounds from plants such as alkali metals, alkaline earth metals and heavy metals, as well as to provide an overview of internal and external mineral content until they become extracts.

Drying shrinkage has a goal to give the maximum amount of compounds lost in the drying process. In this study, a moisture balance was used at a temperature of 105°C from the three extracts of *n*-hexane, ethyl acetate, and 70% ethanol. The three results (Table 2) fall into the drying shrinkage limit of no more than 10%.

Table 3. Phytochemical Screening

Detected compounds	Extract type		
	<i>n</i> -Hexane	Ethyl acetate	Ethanol 70%
Alkaloids			
- Dragendor	-	-	+
- Bouchardat	-	-	+
- Mayer	+	+	+
Phenolic	+	+	+
Flavonoids	+	+	+
Saponins	+	+	+
Tannins	+	+	+
Terpenoids	-	-	-
Steroids	+	+	+

Note: (+) There is a compound (-) There is no compound

The results of phytochemical screening can be seen in (Table 3). Phytochemical screening was carried out to identify the compounds contained in *n*-hexane extract, ethyl acetate extract, and 70% ethanol extract of katuk leaves. Screening is an important step in an effort to reveal the potential of plant resources [1]. The 70% ethanol extract showed positive results for alkaloids, while the ethyl acetate and *n*-hexane extracts were negative.

Determination of total phenolic levels uses the *Folin-Ciocalteu* method because it is easy, cheap, fast to do, and can be done routinely in the laboratory. Polyphenols in plant extracts react with the reagent used, namely *Folin-Ciocalteu* (Blainski et al., 2013). Phenolic compounds react with *Folin-Ciocalteu* reagent only in an alkaline environment so that proton dissociation occurs in phenolic compounds into phenolic ions, hence 7.5% Na₂CO₃ is used to make an alkaline environment. The hydroxyl group in the phenolic compound reacts with the *Folin-Ciocalteu* reagent to form a blue *molybdenum-tungsten* complex which can be detected by a spectrophotometer [19]. Gallic acid (3,4,5-*Trihydroxybenzoic acid*) was used as standard. Gallic acid was chosen as the standard because it represents the general properties of phenolic compounds. In this case gallic acid is a natural phenolic compound that has a strong antioxidant effect [20].

Determination of the maximum wavelength of gallic acid aims to determine at what wavelength the compound reaches the highest absorbance value. The result of measuring the maximum wavelength is 765.50 nm with an absorbance of 0.3445 as the maximum absorbance value. The operating time of the results obtained is stable absorbance at 60 minutes so that it can be concluded that the purpose of determining the operating time is to get the measurement time when the reaction has been running optimally which is characterized by a stable absorbance, so as to maximize the measurement [21].

Various concentrations of gallic acid standards were made because the assay used a calibration curve equation (Table 4), so to make a calibration curve, several concentrations were made in order to obtain a linear equation. The results obtained were at concentrations of 18, 30, 42, 54, and 66 ppm, the data met the requirements of *Lambert Beer's* law. The calibration curve obtained by the linear regression equation $y = 0.0093.x + 0.1824$ with a correlation coefficient (r) = 0.9944.

In the determination of the total phenolic content of katuk leaf extract, as seen in (Table 5) that the 70% ethanol extract had the highest concentration of 16.71 mgGAE/g. The extract was followed by ethyl acetate and *n*-hexane with a concentration of 2.95 mgGAE/g and 0.57 mgGAE/g extract, respectively. Phenolic compounds are extracted well in 70% ethanol because the phenolic content will increase in the extract as the polarity of the solvent increases. Phenolic also tends to be polar so it can dissolve in polar solvents well. These results are similar to other studies which stated polar extracts have the highest phenolic content followed by semi-polar and non-polar extracts. Methanol, a polar solvent, has a higher phenolic content than acetone which is a semi polar solvent [22].

Table 4. Results of Total Phenolic Levels

Extract type	Average Phenolic content (mgGAE/g Extract \pm SD)
<i>n</i> -Hexane	0,57 \pm 0,02
Ethyl Acetate	2,95 \pm 0,07
Ethanol 70%	16,71 \pm 0,43

Table 5. Results of Total Flavonoid Levels

Extract type	Average levels of Flavonoids (mgQE/g Extract \pm SD)
<i>n</i> -Hexane	55,57 \pm 0,11
Ethyl Acetate	88,79 \pm 0,73
Ethanol 70%	6,23 \pm 0,05

Determination of total flavonoid levels uses quercetin standards. Quercetin is a flavonoid class of the flavonol group [23]. Meanwhile, Quantitative Analysis is conducted by applying the UV-Vis spectrophotometric method (Hanani, 2015). The maximum wavelength resulted from this study was 434 nm. This was due to the reaction of the $AlCl_3$ complex which caused the shift of the wave to become visible, which was indicated by the solution becoming more yellow, and the addition of sodium acetate to maintain the visible wavelength [24]. $AlCl_3$ reagents with flavonoids form complexes between neighboring hydroxyl groups and ketones or with neighboring hydroxyl groups. $AlCl_3$ will react with the ketone group at C4 and the OH group at C3 or C5 in flavone or flavonoid compounds to form a yellow stable complex compound.

The results of the absorbance measurement of the quercetin standard solution to obtain a linear calibration curve obtained results such as (Table 6). These results were entered into the regression equation which then produced linear $y = 0.0024x + 0.1897$, $r = 0.9931$. The flavonoid content was expressed in mgQE/g Extract.

After reading the levels of flavonoids (Table 7) of the graded extract of katuk leaves, the results obtained were *n*-hexane 55.57 mgQE/g, ethyl acetate 88.79 mgQE/g, and 70% ethanol 6.23 mgQE/g. In this study, ethyl acetate had the largest concentration, this is because the content of flavonoid compounds contained in katuk leaves has a low polarity level, namely in the form of aglycones or commonly called aglycone flavonoids. Aglycone flavonoids have less polar properties and tend to be more soluble in chloroform and ether. Flavonoids generally bind to sugars to form glycosides which cause these compounds to be easily soluble in polar solvents, such as methanol, butanol, and ethyl acetate.

In this study, the free radical compound DPPH (2,2-diphenyl-1-picrylhydrazyl) was used. The results obtained on the measurement of the maximum wavelength is 515.5 nm, the wavelength can be used because the wavelength of the maximum absorbance for measurements with the DPPH method is 515-520 nm [25].

Quercetin was chosen as the standard for comparison because quercetin is a natural secondary antioxidant that has been shown to have free radical scavenging activity. Quercetin, which is a flavonoid group, has several biological activities. These activities can be attributed to the antioxidant properties of quercetin, including its ability to scavenge free radicals [23].

The results of antioxidant activity (Table 8) show that 70% ethanol extract has a lower IC_{50} value than ethyl acetate and *n*-hexane, the smaller the % inhibition obtained, the stronger the antioxidant activity produced [7]. The compounds contained in the ethanol extract are an accumulation of polar, semi-polar, and non-polar compounds. When the extract is macerated in stages, the synergistic function

between the compounds will reduce because the components contained in the extract have been separated. The non-polar chemical components were extracted in the *n*-hexane solvent, the semi-polar chemical components were extracted in ethyl acetate, and chemical components that are polar were extracted in 70% ethanol solvent. This causes the antioxidant activity of 70% ethanol extract to be the strongest and at the same time has the least % inhibition, compared to others.

Table 6. Antioxidant Activity Test Results and SPF

Sample Type	Average antioxidant activity (IC ₅₀ ± SD)	SPF
Quercetin	8,83	-
<i>n</i> -Hexane	88,43 ± 1,20	2,13
Ethyl Acetate	77,65 ± 1,78	5,23
Ethanol 70%	70,33 ± 1,64	2,13

Ethyl acetate extract gave the highest SPF results compared to ethanol and *n*-hexane extracts. The three extracts belonged to the minimum SPF capability (range 2 to 4). Phytochemical compounds in the extract that may play a role in the potential for absorption of ultraviolet light are flavonoids and phenolic compounds [26]. Several groups of active compounds derived from natural ingredients such as flavonoids, tannins, anthraquinones, cinnamates, and glycosides are reported to have the ability to protect against UV rays [27].

4. Conclusions

Based on the results obtained in this study, the determination of total phenolic content had the highest concentration in 70% ethanol extract, which was then followed by ethyl acetate extract and *n*-hexane extract. In the determination of total flavonoid content, the highest yield was in the ethyl acetate extract, then *n*-hexane content, while the 70% ethanol had the lowest content. The results of the antioxidant activity test used the DPPH method. IC₅₀ of *n*-hexane extract had the highest concentration in this activity test, followed by ethyl acetate and 70% ethanol extract. The smaller the IC₅₀ value, the higher the antioxidant activity produced. Multilevel extraction greatly affects the levels of each extract used. Besides, it will also look for different compounds in the solvent used.

References

- [1] P. Tiwari, M. Kaur, and H. Kaur, "Phytochemical screening and Extraction: A Review," 2011.
- [2] Z. Zuraida, S. Sulistiyani, D. Sajuthi, and I. H. Suparto, "FENOL, FLAVONOID, DAN AKTIVITAS ANTIOKSIDAN PADA EKSTRAK KULIT BATANG PULAI (*Alstonia scholaris* R.Br)," *J. Penelit. Has. Hutan*, vol. 35, no. 3, pp. 211–219, 2017, doi: 10.20886/jphh.2017.35.3.211-219.
- [3] S. Handani and A. Indyah Sulisty, "Synthesis of Hydroxyl Radical Scavengers from Benzalacetone and its Derivatives," *J. Phys. Sci.*, vol. 19, no. 2, pp. 61–68, 2008, [Online]. Available: [http://web.usm.my/jps/19-2-08/article 19-2-7.pdf](http://web.usm.my/jps/19-2-08/article%2019-2-7.pdf).
- [4] Y. D. Putri, H. Kartamihardja, and I. Lisna, "Yola et al 2019," *Formulasi dan Eval. Losion Tabir Surya Ekstrak Daun Stevia (Stevia rebaudiana Bertoni M)*, vol. 6, no. 1, pp. 32–36, 2019.
- [5] D. Andini, "Potential Of Katuk Leaf (*Sauropus androgynus* L. Merr) As Aphrodisiac," *J. Major.*, vol. 3, no. 7, pp. 16–21, 2014.
- [6] N. Andarwulan, R. Batari, D. A. Sandrasari, B. Bolling, and H. Wijaya, "Flavonoid content and antioxidant activity of vegetables from Indonesia," *Food Chem.*, vol. 121, no. 4, pp. 1231–1235, 2010, doi: 10.1016/j.foodchem.2010.01.033.
- [7] C. F. Zuhra, J. B. Tarigan, and H. Sihotang, "Aktivitas Antioksidan Senyawa Flavonoid Dari Daun Katuk (*Sauropus Androgynus* (L) Merr.)," 2010.
- [8] N. P. E. Hikmawanti, S. Fatmawati, and A. W. Asri, "The effect of ethanol concentrations as the

- extraction solvent on antioxidant activity of Katuk (*Sauropus androgynus* (L.) Merr.) leaves extracts,” *IOP Conf. Ser. Earth Environ. Sci.*, vol. 755, no. 1, 2021, doi: 10.1088/1755-1315/755/1/012060.
- [9] N. P. E. Hikmawanti, S. Fatmawati, Z. Arifin, and . V., “Pengaruh Variasi Metode Ekstraksi Terhadap Perolehan Senyawa Antioksidan Pada Daun Katuk (*Sauropus androgynus* (L.) Merr),” *J. Farm. Udayana*, vol. 10, no. 1, p. 1, 2021, doi: 10.24843/jfu.2021.v10.i01.p01.
- [10] E. C. Jeyaseelan, S. Jenothiny, M. K. Pathmanathan, and J. P. Jeyadevan, “Antibacterial activity of sequentially extracted organic solvent extracts of fruits, flowers and leaves of *Lawsonia inermis* L. from Jaffna,” *Asian Pac. J. Trop. Biomed.*, vol. 2, no. 10, pp. 798–802, 2012, doi: 10.1016/S2221-1691(12)60232-9.
- [11] M. Arista, “Aktivitas antioksidan ekstrak etanol 80% dan 96% daun katuk (,” *J. Ilm. Mhs. Univ. Surabaya*, vol. 2, no. 2, pp. 1–16, 2013.
- [12] S. Winarti, U. Sarofa, and D. Anggrahini, “EKSTRAKSI DAN STABILITAS WARNA UBI JALAR UNGU (*Ipomoea batatas* L .,) SEBAGAI PEWARNA ALAMI,” *J. Tek. Kim.*, vol. 3, no. 1, pp. 207–214, 2008.
- [13] Y. Y. Lim and J. Murtijaya, “Antioxidant properties of *Phyllanthus amarus* extracts as affected by different drying methods,” *LWT*, vol. 40, no. 9, pp. 1664–1669, 2007, doi: 10.1016/j.lwt.2006.12.013.
- [14] D.-O. Kim, S. W. Jeong, and C. Y. Lee, “Antioxidant capacity of phenolic phytochemicals from various cultivars of plums,” *Food Chem.*, vol. 81, no. 3, pp. 321–326, 2003, doi: [https://doi.org/10.1016/S0308-8146\(02\)00423-5](https://doi.org/10.1016/S0308-8146(02)00423-5).
- [15] M. M. Rahman, M. B. Islam, M. Biswas, and A. H. M. Khurshid Alam, “In vitro antioxidant and free radical scavenging activity of different parts of *Tabebuia pallida* growing in Bangladesh,” *BMC Res. Notes*, vol. 8, no. 1, p. 621, 2015, doi: 10.1186/s13104-015-1618-6.
- [16] W. A. d. S. Almeida *et al.*, “Photoprotective activity and increase of SPF in sunscreen formulation using lyophilized red propolis extracts from Alagoas,” *Rev. Bras. Farmacogn.*, vol. 29, no. 3, pp. 373–380, 2019, doi: 10.1016/j.bjp.2019.02.003.
- [17] V. Damogalad, H. Jaya Edy, and H. Sri Supriati, “Formulasi Krim Tabir Surya Ekstrak Kulit Nanas (*Ananas Comosus* L Merr) Dan Uji in Vitro Nilai Sun Protecting Factor (Spf),” *PHARMACON J. Ilm. Farm. – UNSRAT*, vol. 2, no. 02, pp. 2302–2493, 2013.
- [18] R. Y. Senja, E. Issusilaningtyas, A. K. Nugroho, and E. P. Setyowati, “Perbandingan Metode Ekstraksi dan Variasi Pelarut terhadap Rendemen dan Aktivitas Antioksidan Ekstrak Kubis Ungu (*Brassica oleracea* L. var. *capitata* f. *rubra*),” *Maj. Obat Tradis.*, vol. 19, no. 1, pp. 43–48, 2014.
- [19] R. Alfian and H. Susanti, “PENETAPAN KADAR FENOLIK TOTAL EKSTRAK METANOL KELOPAK BUNGA ROSELLA MERAH (*Hibiscus sabdariffa* Linn) DENGAN VARIASI TEMPAT TUMBUH SECARA SPEKTROFOTOMETRI,” *Pharmaciana*, vol. 2, no. 1, 2012, doi: 10.12928/pharmaciana.v2i1.655.
- [20] Y. M. Huliselan, M. R. J. Runtuwene, and D. S. Wewengkang, “Antioxidant Activity of Ethanol, Ethyl Acetate and n-Hexane Extract from *Sesuvium portulacastrum* Leaves (*Clerodendron squamatum* Vahl.),” *Pharmacogn.*, vol. 4, no. 3, pp. 155–163, 2015, [Online]. Available: <https://ejournal.unsrat.ac.id/index.php/pharmacogn/article/view/8855>.
- [21] T. G. Arikalang, S. Sudewi, and J. A. Rorong, “Penentuan Kandungan Total Fenolik pada Ekstrak Daun Gedi Hijau (*Abelmoschus manihot* L .) yang Diukur dengan Spektrofotometer UV-Vis,” *Pharmacogn J. Ilm. Farm.*, vol. 7, no. 3, pp. 14–21, 2018.
- [22] F. Ulfa, A. Anggo, and R. Romadhon, “Uji Potensi Aktivitas Antioksidan Dengan Metode Ekstraksi Bertingkat Pada Lamun Dugong (*Thalassia Hemprichii*) Dari Perairan Jepara,” *J. Pengolah. dan Bioteknol. Has. Perikan.*, vol. 3, no. 3, pp. 32–39, 2014.
- [23] M. Materska, “QUERCETIN AND ITS DERIVATIVES: CHEMICAL STRUCTURE AND BIOACTIVITY– A REVIEW,” *Polish J. Food Nutr. Sci.*, vol. 58, no. 4, pp. 407–413, 2008, [Online]. Available: <http://journal.pan.olsztyn.pl/QUERCETIN-AND-ITS-DERIVATIVES->

- CHEMICAL-STRUCTURE-AND-BIOACTIVITY-8211-A-REVIEW,98157,0,2.html.
- [24] C. C. Chang, M. H. Yang, H. M. Wen, and J. C. Chern, "Estimation of total flavonoid content in propolis by two complementary colometric methods," *J. Food Drug Anal.*, vol. 10, no. 3, pp. 178–182, 2002, doi: 10.38212/2224-6614.2748.
- [25] P. Molyneux, "The Use of the Stable Free Radical Diphenylpicryl-hydrazyl (DPPH) for Estimating Antioxidant Activity," *Songklanakar J. Sci. Technol.*, vol. 26, no. December 2003, pp. 211–219, 2004, doi: 10.1287/isre.6.2.144.
- [26] C. Malsawmtluangi, D. K. Nath, I. Jamatia, E. Z. Lianhingthangi, and L. Pachuau, "Determination of Sun Protection Factor (SPF) number of some aqueous herbal extracts," *J. Appl. Pharm. Sci.*, vol. 3, no. 9, pp. 150–151, 2013, doi: 10.7324/JAPS.2013.3925.
- [27] M. M. Donglikar and S. L. Deore, "Development and evaluation of herbal sunscreen," *Pharmacogn. J.*, vol. 9, no. 1, pp. 83–97, 2017, doi: 10.5530/pj.2017.1.15.