

PAPER • OPEN ACCESS

Preface International Conference on Environmental, Energy and Earth Science (ICEEES)

To cite this article: 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1041** 011001

View the [article online](#) for updates and enhancements.

You may also like

- [Small and Medium Enterprises in Pekanbaru: Implementing an Integrated Work Program](#)
Hardi
- [Analyzing the critical locations in response of constructed and planned dams on the Mekong River Basin for environmental integrity](#)
Yuan Gao, Shiblu Sarker, Tanni Sarker et al.
- [Nutrient testing of organic liquid fertilizer in tempeh factory in Pekanbaru City](#)
Neng Susi, Vonny Indah Sari, Indra Purnama et al.

ECS Toyota Young Investigator Fellowship



For young professionals and scholars pursuing research in batteries, fuel cells and hydrogen, and future sustainable technologies.

At least one \$50,000 fellowship is available annually.
More than \$1.4 million awarded since 2015!



Application deadline: January 31, 2023

Learn more. Apply today!

Preface International Conference on Environmental, Energy and Earth Science (ICEEES)

Universitas Lancang Kuning Pekanbaru organized with **the International Conference on Environmental, Energy and Earth Science (ICEEES)** on September 22, 2021 in Pekanbaru, Indonesia. The conference is aims to exchange knowledge and research finding among academicians, researchers, professionals, policy makers, and postgraduate students.

The Awareness to increase the number of publications on the results of research that has been done, deserves appreciation by all parties. Because of that, Universitas Lancang Kuning Pekanbaru provide motivation and space for researcher to disseminate their research and accommodate the result of research that has been done. **The International Conference on Environmental, Energy and Earth Science (ICEEES)**, is another International Conference held by Universitas Lancang Kuning Pekanbaru.

The International Conference on Environmental, Energy and Earth Science (ICEEES) was chosen to be implemented virtually, this is because the cov-19 pandemic is still spreading. The conference was perform using zoom. **The International Conference on Environmental, Energy and Earth Science (ICEEES)** event is virtually implemented with a model that all invited speakers are given time to present their material for 30 minutes every invited speaker and after that a question and answer is carried out with the participants with a direct questioning system, through chat forums and Q&A forums provided by the zoom application. Overall, the conference took 5 hours the number of participants who joined the zoom room was recorded at participants. Participants came from few countries, namely Indonesia, Malaysia, India, Egypt and Australia.

The International Conference on Environmental, Energy and Earth Science (ICEEES) is implemented with the support of a stable internet network system and a zoom application. In the implementation there were several technical obstacles encountered by the participants, namely the difficulty of joining the zoom application due to the unstable internet signal. The holding of a virtual conference felt less meaningful, due to the lack of interaction between speakers and participants

The International Conference on Environmental, Energy and Earth Science (ICEEES) committee received 124 manuscripts and a total of 117 papers were presented and discussed. The papers were authored by researchers from Indonesia, Malaysia, India, Egypt, japan and Australia.

All papers have been scrutinized by a panel of reviewers who provide critical comments and corrections, and thereafter contributed to the improvement of the quality of the papers. Based on the reviewer's reports, 92 papers were selected and eligible to be published in the proceeding.

We sincerely express our gratitude to the international/national advisory committee, presenters, organizing committee members, session chairs, all members of organization, participants, contributors and all the members of **The International Conference on Environmental, Energy and Earth Science (ICEEES)**. Last but not the least, we are thankful to IOP EES Conference Series for producing the proceeding.



Organizing Committee

General Chair :

Dr. Anto Ariyanto, M.Si (Universitas Lancang Kuning, Indonesia)

General Co-Chair:

Dr. H. Jeni Wardi, S.E., M.Ak., Ak., CA (Universitas Lancang Kuning, Indonesia)

Secretary :

Dr. David Setiawan, M.T (Universitas Lancang Kuning, Indonesia)

Treasurer and Finance Chair :

Ambar Tri Ratnaningsih, S.Hut., M.Si (Universitas Lancang Kuning, Indonesia)

Publication Chair :

Robbi Rahim, S.Kom., M.Kom (STIM Sukma Medan, Indonesia)

Website and Social Media Chair :

Guntoro, S.T., M.Kom (Universitas Lancang Kuning, Indonesia)

Febrizal Alfarasy Syam, S.Kom., M.Kom (Universitas Lancang Kuning, Indonesia)

Ismar Husnudin, S.T (Universitas Lancang Kuning, Indonesia)

Editors

1. Prof. Dr. Joran (Linkoping University, Swedia)
2. Prof. Dr. Amran bin Hamzah (Universitas Teknologi Malaysia, Malaysia)
3. Prof. Dr. Khaerurijal (Institut Teknologi Bandung, Indonesia)
4. Prof. Dr. Asep Saefuddin (Institut Pertanian Bogor, Indonesia)
5. Prof. Dr. Mauridi (Institut Teknologi Sepuluh November, Indonesia)
6. Prof. Dr. Adi Soeprijanto (Institut Teknologi Sepuluh November, Indonesia)
7. Prof. Dr. Syafrani (Universitas Lancang Kuning, Indonesia)
8. Prof. Dr. Sudi Fahmi (Universitas Lancang, Kuning, Indonesia)
9. Prof. Dr. Hariadi Kartodiharjo (Institut Pertanian Bogor, Indonesia)
10. Prof. Dr. Sumbangan Baja (Universitas Hasanuddin, Indonesia)
11. Prof. Wiendu Nuryanti Ph.D (Universitas Gajah Mada, Indonesia)
12. Prof. Dr. Chairul Saleh (Universitas Islam Indonesia, Indonesia)
13. Prof. Dr. Richardus Eko Indrajit (STMIK Perbanas, Indonesia)

Organizing Institutions

- Universitas Lancang Kuning

PAPER • OPEN ACCESS

Peer Review Statement

To cite this article: 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1041** 011002

View the [article online](#) for updates and enhancements.

You may also like

- [Peer review declaration](#)
- [Peer review declaration](#)
- [Peer Review Declaration Form Guidance](#)

ECS Toyota Young Investigator Fellowship



For young professionals and scholars pursuing research in batteries, fuel cells and hydrogen, and future sustainable technologies.

At least one \$50,000 fellowship is available annually.
More than \$1.4 million awarded since 2015!



Application deadline: January 31, 2023

Learn more. Apply today!

Peer Review Statement

All papers published in this volume have been reviewed through processes administered by the Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

- **Type of peer review:** Single Anonymous
- **Conference submission management system:** <https://iceees.unilak.ac.id/paper-submission/>
- **Number of submissions received:** 129
- **Number of submissions sent for review:** 129
- **Number of submissions accepted:** 87
- **Acceptance Rate (Submissions Accepted / Submissions Received × 100):** 67.4
- **Average number of reviews per paper:** 2
- **Total number of reviewers involved:** 15
- **Contact person for queries:**
Name: Robbi Rahim
Email: robbirahim@ieee.org
Affiliation: Sekolah Tinggi Ilmu Manajemen Sukma



Table of contents

Volume 1041

2022

◀ Previous issue Next issue ▶

International Conference on Environmental, Energy and Earth Science 22/09/2021 - 23/09/2021

Online

Accepted papers received: 27 May 2022

Published online: 23 June 2022

Open all abstracts

Preface

OPEN ACCESS 011001

Preface International Conference on Environmental, Energy and Earth Science (ICEEES)

+ Open abstract  View article  PDF

OPEN ACCESS 011002

Peer Review Statement

+ Open abstract  View article  PDF

Agriculture

OPEN ACCESS 012001

The effect of chitosan addition on physicochemical properties, proximate and antioxidant of functional drinks of red and purple roselle (*Hibiscus sabdariffa* L.) extracts

Kusnadi, Aldi Budi Riyanta and Rizki Febriyanti

+ Open abstract  View article  PDF

OPEN ACCESS 012002

The characteristics of bio-oil hydrothermal pyrolysis empty fruit bunches based on powder size and hydrogen gas pressure

Rina Novia Yanti, Erliza Hambali, Gustan Pari and Ani Suryani

+ Open abstract  View article  PDF

OPEN ACCESS 012003

The effect of proportion of ganyong starch and waste of straw rice on biodegradable foam production as sustainable packaging

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Gresita Fitaloka Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012004

Physicochemical and antioxidant characteristics of black garlic from indigenous Indonesian garlic

K S Sasmitaloka, SM Widayanti, I Mulyawanti and ES Iriani

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012005

Efficiency analysis of vannamei shrimp supply chain performance in PT Sari Indonesia Group, Suak Pandan Village, West Aceh Regency, Aceh Province

Yoga Nugroho, Teuku Athaillah and Muhibbul Qadri

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012006

Nutrient testing of organic liquid fertilizer in tempeh factory in Pekanbaru City

Neng Susi, Vonny Indah Sari, Indra Purnama and Hazra Yuvendus

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012007

Extraction of catfish bone waste (*pangasius hypophtalmus*) by utilizing an organic liquid of pineapple peel waste (*ananas comosus*) into gelatin

Ira Oktaviani Rz, Rahimatul Uthia, Fathul Jannah and Alexsander Yandra

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012008

The effect of orange peel waste addition on ZnO characteristics and its ability to degrade methyl orange

Aulia Dewi Rosanti, Fahmi Hidayat, Yuly Kusumawati, Arif Fadlan, Anggita R.K Wardani and Eva Umi Latifah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012009

Carbon sequestration of city agriculture: between farming and non-farming land

D.R.D. Hastuti, R. Darma, D. Salman, S. Santoso and A. Rahim

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012010

Comparison predicted using nonlinear model and in vivo studies growth production of broiler after fed different levels of fermented tamarind flour as replacing soya bean meal

Onsijofan, D.N. Adlis, MBH Natsim, Y.F. Nuringtias and A.D. Radianto

Use of cookies. To find out more, see our Privacy and Cookies policy.



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012011

Dry matter (DM), crude protein (CP) and water consumption in sheep which supplemented from *Moringa* leaves meal and palm oil in elephant silage-based feed

S Rizqiana, W Suryapratama and F M Suhartati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012012

Effect of various packaging materials for storing ground yellow corn of hybrid C-1 variety on water and amyllum content

Munasik, N Nurbaety, N Hidayat and E Susanti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012013

Exploring the herbaceous plant height and dry matter relationship on a sub-humid rangeland in Zimbabwe

T A Zezai, P H Mugabe and V E Imbayarwo-Chikosi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012014

Potential development based on the agricultural waste in the eastern region of West Java Province

F. D. Perwitasari, A. R. S. Putra, B. Suwignyo and R. Widiati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012015

Farming in Kediri Indonesia: analysis of cluster k-means

N Solikin, B Hartono, Sugiono and Linawati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012016

The Identification and Role pf Mold in Lowering the HCN Levels in Black Cassava Based on the Immersion Duration

Mery Pattipeilohy and Alamanda Pelamonia

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012017

The Effects of Some Additives On Characteristics RVA Profile of Cassava Flour

H Herawati and E. Kamsiati

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Agriculture Engineering and Technology

OPEN ACCESS

012018

A comprehensive review on power tillers usage – bottlenecks and prospects in India

Krishnadas R and Renganathan R

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012019

Android-based feasibility study transformation analysis case study of the RASI development in Cirendeu-Kampung Cimahi

H Herawati, A S Soemantri, E Kamsiati and M. Bachtiar

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012020

Water pump control system in hydroponic plants using the ebb and flow method

David Setiawan, Hamzah, Latifa Siswati, Anto Ariyanto and Guntoro

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012021

Resetting the ground fault relay on a 28 MVA power transformer due to grounding method retrofit

Hamzah Eteruddin, Arlenny, Hadi Siswanto, Yanuar Z. Arief and Fathimah Hasanti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012022

Intelligent ubiquitous technology as a precision agri-food framework: a proposed framework

T A Cahyanto, S Wahjuni, H Sukoco, H Rahmawan and S N Neyman

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012023

Development of internet of things-based monitoring system for application solar energy technology in Bunaken island

M Rumbayan, I Pundoko, D Ruindungan and N V Panjaitan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012024

Technical analysis on household energy consumption: LMDI decomposition index and innovative

Yetty Murni, Eka sudarmaji, Sri Ambarwati and Ismiriati Nasip

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012025

Optimization of electrical energy in the fan using fuzzy logic controllers

Arlenny and David Setiawan

[+ Open abstract](#) [View article](#) [PDF](#)

Animal Biodiversity

OPEN ACCESS

012026

The effect of giving chia seeds on the bodyweight of pregnant mice (*Mus Musculus L.*)

R Uthia and I O Rz

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012027

Supplementing mineral selenium and vitamin e in diets on *in vivo digestibility*, blood glucose, and urea levels of cows

D. Pamungkas, A.S. Putri, F. Firdaus, R. Widiyawati and D.M. Dikman

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012028

Effect of *Saccharomyces cerevisiae* on Probiotic Properties of Goat Milk Kefir

L E Radiati, D L Hati, D Fardiaz and L R H Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012029

Cattle production system in semi-arid area of Timor Island

Y Ngonjo, S Ratnawaty and PR Matitaputty

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012030

Sustainable indigenous cattle production in Brebes Regency, Indonesia: opportunities and threats during pandemic

M Sugiarto, Y. N Wakhidati, O. E. Djatmiko, S. Nur and A Einstein

[+ Open abstract](#) [View article](#) [PDF](#)

Disaster Management

OPEN ACCESS

012031

The role of philanthropy in reducing impact of disaster on economies of ASEAN-9

M S A Majid, R Nurdin, T Azhar and S Sartiyah

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS 012032
Community participation in disaster management in Gondoriyo village administration, Ngaliyan sub-district, Semarang city
Supratiwi, Yuwanto and Kushandajani
[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012033
Assessing and preparedness for earthquake disaster in Salaf-Khalaf Islamic Boarding School
Fatiya Rosyida, Neni Wahyuningtyas, Ardyanto Tanjung and Abdul Kodir
[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012034
Disaster education in disaster-prone schools: a systematic review
R Yusuf, Razali, Sanusi, Maimun, I Fajri and S A Gani
[+ Open abstract](#) [View article](#) [PDF](#)

Environmental Issue and challenge

OPEN ACCESS 012035
LCA methodology for detecting environmental impacts on natural gas drilling process
Masayu Rosyidah, Andiyan Andiyan, Haniek Listyorini, Pranoto Hadi Prayitno, Yuswardi Yuswardi and Yuhanah Yuhanah
[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012036
Community based ecotourism management to strengthen environmental ethics and supports sustainable development in Pelalawan district, Indonesia
Nawari, Thamrin, Nofrizal, Almasdi Syahza, Juandi Muhammad and Nur Islami
[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012037
Community forestry for environmental sustainability and ecotourism: the context and problems in Indonesia
R Sulistiowati, S Wahyuni, M K Yunanto, M Elsera, W E Yudiatmaja and T Samnuzulsari
[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012038
Study on the commitment of oil palm companies to achieve sustainable agriculture in Riau Province from the perspective of pesticide use
S Syafrani, I Purnama, A Mutamima and W N Dewi
[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



OPEN ACCESS

012039

Assessing the Feasibility of Migration Policy from LPG Stoves to Induction Stoves in Indonesia

Tri Anggono, Iyung Ruslan, Chrisnawan Anditya, Dian Galuh Cendrawati and Muhammad Indra al Irsyad

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012040

The innovation of ecological fiscal transfers policy at Siak Regency

Muhammad Ichsan Kabullah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012041

Farmers' perceptions of the performance of agricultural instructors in mangosteen farmers' groups in Segati Village, Langgam District, Pelalawan Regency

Asgami Putri, Latifa Siswati and Hamdan Yasid

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012042

Environment Conservation Responsibility in Community Development

N Sutisna, H Herdiansyah and P Sanjatmiko

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012043

Environmental Conservation in Ciletuh Geopark, West Java

R A Zuvara, H Herdiansyah and D Asteria

[+ Open abstract](#) [View article](#) [PDF](#)

Food Science and Technology

OPEN ACCESS

012044

Effectiveness of silica gel from palm kernel shell ash as a moisture absorber of bottle packaging medicine

Lidya Novita and Iswadi Idris

[+ Open abstract](#) [View article](#) [PDF](#)

Land Use Management

OPEN ACCESS

012045

Comparative study of village community participation in peat restoration in the Peat Swamp Forest Management Unit (PSFMU) Tebing Tinggi Island, Indonesia

Elhi Malik, Agus Iko Skripriyana, Sindangdji, Jc. Ws Satriyasa, Igr Witayonarko and M. Nasirul find out more, see our Privacy and Cookies policy.



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012046

Swot analysis of utilization of palm oil mill effluent to improve soil quality and crop productivity (case study at PT X, Lampung, Indonesia)

Dela Febriana, Sylvia Madusari and Vira Irma Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012047

Groundwater level fluctuations in peatlands as responses to canal block construction in Sungai Tohor, Tebing Tinggi Island, Indonesia

H. Malik, Ig. L. S. Purnama, Sudarmadji, L. W. Santosa and M. Naufal

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012048

Identification of land use and land cover using the image Landsat 8 in upstream Lematang sub-watershed by support vector machine and random trees methods

A Dinata, F Dhiniati and L E Diansari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012049

Conservation and utilization on Sawahlunto ex mining area for cultural landscape

Rika Cheris and Amanda Rosetia

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012050

Measuring cross-sector collaboration an effective solution for solving unlicensed mining problems in Kuantan Singingi, Indonesia

MY Tiyas Tinov, Aras Mulyadi, Sujianto, Zulkarnain, Ardi Putra, Tika Mutia and Tito Handoko

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012051

Helminthiasis of rabbits on the upland and lowland areas and the risk factors

D Indrasanti, M Indradji, Sufiriyanto, M Samsi and E Yuwono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012052

Water Quality Analysis Based on Land Use in Sekampung River, Lampung, Indonesia

K Fitriani, T K Nufutomo and R Putra

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



Oil Plantation and Management

OPEN ACCESS

012053

Application of oil palm empty bunches as organic mulch in oil palm plantation (*Elaeis guineensis* Jacq.): an evaluation and SWOT analysis

Lala Novita Sari, Sylvia Madusari and Vira Irma Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012054

Oil palm trunk replanting as brown sugar raw materials

Latifa Siswati, Enny Insusanty, Neng Susi and Nopryanti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012055

Prediction of palm oil production in Riau Province using the single exponential smoothing method

Guntoro, Lisnawita, Zamzami and David Setiawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012056

Study of biogas utilization as waste-to-energy plant and transport modelling of iron (Fe), lead (Pb) and copper (CU) in leachate at Muara Fajar landfill Pekanbaru

H Yuvendius, E Zondra, Zainuri and V I Sari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012057

Utilization of oil palm waste as animal feed for Bali cattle in Margomulyo Village, Central Bengkulu

E Ramon, Z Efendi, H Kusnadi, S Yuliasari, A Ishak and WA Wulandari

[+ Open abstract](#) [View article](#) [PDF](#)

Plant Biodiversity

OPEN ACCESS

012058

Etnobotany of medicine plants in the community of Rawang Kao Village, Siak, Indonesia

Sri Wahyuni and Mar'atul Afidah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012059

Can mangrove intensive policies in Indonesia benefit more coastal communities? environmental policy analysis

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012060

Validation of UV-VIS spectrophotometric method for the determination of Calcium in Maman (cleome gynandra l)

Fathul Jannah, S Meri Agritubella, Ira Oktaviana and Lily Restusari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012061

Utilization of bamboo charcoal to decrease tannin content of lindur fruit flour (Bruguiera gymnorrhiza)

Niken Pujirahayu, Zakiah Uslinawaty, Nuhayati Hadjar and Muhtadir

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012062

Supplementing safflower oil (*Carthamus Tinctori L*) and inocytol in feed and its effect

Ning Iriyanti, Ismoyowati, Efka Aris Rimbawanto and Modawy Abdelgader Albasheer Altayb

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012063

The effects of nano encapsulated *Syzygium Polyanthum* leaves extract inclusion in drinking water

A N 'Afifah, Zuprizal and N D Dono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012064

Effects of plant extracts against gastrointestinal nematodes of livestock: a meta-analysis

A Jayanegara, D M Sinaga and E B Laconi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012065

Coconut oil effects on in vitro ruminal fermentation and methanogenesis

A Cusiayuni, RK Nurfatahillah, D Evvyernie, A Jayanegara and KG Wiryawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012066

Effect of palm oil supplementation level on in vitro ruminal fermentability

RK Nurfatahillah, A Cusiayuni, A Jayanegara, KG Wiryawan and D Evvyernie

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

012066

The Composition of Goat Milk Kefir Added with *Moringa oleifera* Leaf Powder

S R N Endah, P D Wulasari, A Nofriyaldi, Nurliyani and E Harmayani

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012068

The effect of safflower oil (*Carthamus Tinctorius* L.) and inositol supplementation on egg production

Ismoyowati, N Iriyanti and M Abdelgader

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012069

The Anti-Inflammatory Activity of Cherry Leaf Extract (*Muntingia Calabura* L.) Balm Stick

Fitria Nugrahaeni, Kriana Efendi and Abdul Kholik Aziz

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012070

The Effect of HPMC Concentration as a Gelling Agent on Color Stability of Copigmented Blush Gel Extract of Purple Sweet (*Ipomoea Batatas* (L.) Lam.)

Fitria Nugrahaeni, Nining and Redina Okvianida

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012071

Sunscreen Factor Formulation and Test of Gel Preparations of 70% Ethanol Extract on Arabica Coffee Leaf (*Coffea arabica* L.)

Sofia Fatmawati, Fitria Nugrahaeni, Fith Khaira Nursal and Ana Fitriana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012072

Antioxidant Activity and Sun Protection Factor (SPF) Graded Extract of Katuk Leaves (*Sauropus androgynus* (L.) Merr.)

Sofia Fatmawati, Ni Putu Ermi Hikmawanti, Akbar Fadillah and Awlia Mustia Putri

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012073

Anti-Inflammatory Activity of Water Extract of *Talinum paniculatum*. (Jact). Gaertn Leaves on Wistar Rat

A Emelda, Sukmawati, A L Dongke and I Marzuki

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



Sustainable Food Systems

OPEN ACCESS

012074

Assessing food safety compliance for food SMEs in Indonesia

Tri Fajarwaty and David Jukes

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012075

In-silico screening of inhibitor on protein epidermal growth factor receptor (EGFR)

Jeremi Ongko, Jesica Viona Setiawan, Alfina Gracia Feronytha, Agnes Juliana, Andy Effraim, Mariana Wahjudi and Yulanda Antonius

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012076

Effect of Various Feed Additives on Carcass and Meat Quality of Two Different Strains of Chickens

E Tugiyanti and E Susanti

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012077

Microbiological profile of concentrated yoghurt manufactured from low and full fat milk during storage

T Setyawardani, J Sumarmono and D R Kusuma

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012078

Effect of dietary supplementation with flying fish (exocoetidae) silage on the egg quality and duck performance

L Ambarwati and S P Syah

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012079

Production profile and microbiological properties of locally produced pork sausage in the Philippines

MM Dealino and CM Bueno

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012080

Farming Sustainability: A Sensitivity Analysis

Y N Wakhidati, M Sugiarto, H Aunurrohman, S Mastuti and J A Suryani

[+ Open abstract](#)[View article](#)[PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



-
- OPEN ACCESS** 012081
Improving genetic quality of local livestock for achieving food security
E Kurnianto
[+ Open abstract](#) [View article](#) [PDF](#)

-
- OPEN ACCESS** 012082
Current goat milk production, characteristics, and utilization in Indonesia
J Sumarmono
[+ Open abstract](#) [View article](#) [PDF](#)

Waste Management

-
- OPEN ACCESS** 012083
The impact of agricultural waste on river water quality of kreo watershed in Semarang city
W Setyaningsih and R S Sanjaya
[+ Open abstract](#) [View article](#) [PDF](#)

-
- OPEN ACCESS** 012084
Utilization of plastic waste as an eco-friendly construction material
Zainuri, G Yanti and S W Megasari
[+ Open abstract](#) [View article](#) [PDF](#)

-
- OPEN ACCESS** 012085
Biogas as methane mitigation path on dairy cattle waste management
A Herliatika and Y Widiawati
[+ Open abstract](#) [View article](#) [PDF](#)

Water, Climate and Emission Science

-
- OPEN ACCESS** 012086
Does rebound effect influence the factors of carbon emission in Indonesia? Kaya index and LMDI decomposition
Eka Sudarmaji, Noer Azam Achسانی, Yandra Arkeman and Idqan Fahmi
[+ Open abstract](#) [View article](#) [PDF](#)

-
- OPEN ACCESS** 012087
Study of phytoplankton biology index and water quality parameters of kali Surabaya River
O H Cahyonugroho, S Hariyanto and G Supriyanto
[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012088

Factor and perception analysis of the impact of boat parking on river pollution

B Rahman and M S Aryanto

[+ Open abstract](#)[View article](#)[PDF](#)

JOURNAL LINKS

[Journal home](#)[Journal scope](#)[Information for organizers](#)[Information for authors](#)[Contact us](#)[Reprint services from Curran Associates](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).



PAPER • OPEN ACCESS

Sunscreen Factor Formulation and Test of Gel Preparations of 70% Ethanol Extract on Arabica Coffee Leaf (*Coffea arabica* L.)

To cite this article: Sofia Fatmawati *et al* 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1041** 012071

View the [article online](#) for updates and enhancements.

You may also like

- [Price transmission of Arabica coffee between North Sumatra with state main export purpose \(America-Europe\)](#)
Z Damanik, Rahmanta and S F Ayu
- [Comparison of Gayo Arabica coffee taste sensory scoring system between Eckenrode and Fuzzy-Eckenrode methods](#)
R Fadhil and D Nurba
- [Strategy to increase selling price of organic Toraja Arabica coffee at farmers levels in International Markets](#)
Y Musa, R Darma and Y B S Panggabean



ECS The Electrochemical Society
Advancing solid state & electrochemical science & technology

242nd ECS Meeting

Oct 9 – 13, 2022 • Atlanta, GA, US

Early hotel & registration pricing ends September 12

Presenting more than 2,400 technical abstracts in 50 symposia

The meeting for industry & researchers in

BATTERIES
ENERGY TECHNOLOGY
SENSORS AND MORE!

 Register now!

 **ECS Plenary Lecture featuring M. Stanley Whittingham,**
Binghamton University
Nobel Laureate –
2019 Nobel Prize in Chemistry



Sunscreen Factor Formulation and Test of Gel Preparations of 70% Ethanol Extract on Arabica Coffee Leaf (*Coffea arabica* L.)

Sofia Fatmawati*, Fitria Nugraheni, Fith Khaira Nursal, Ana Fitriana

Universitas Muhammadiyah Prof. DR. HAMKA, Jakarta, Indonesia

*sofia.fatmawati@uhamka.ac.id

Abstract. Arabica coffee leaf extract (*Coffea arabica* L.) contains phenolic and flavonoid compounds that are potentially active as sunscreen because it might absorb UV (Ultra Violet) A and UV B rays. It is well-known that UV rays exposure causes a negative effect to skin, yet is possible to be avoided by applying sunscreen. This research aims to test the effect of arabica coffee leaf extract on formulation of gel preparations and to discover how its concentration affects the amount of gel preparations' sunscreen factor. There were 4 formulas with different extract concentrations whose characteristics were tested chemically, physically, and they were tested to measure its SPF number. From this test, it was discovered that the gel was qualified and it was plastic thixotropic flow. The measurement of SPF number proposed that F1 could not be categorized as protector because it was only 0.6 while F2, F3, and F4 showed that these formulas could be categorized as minimum protector. The samples were also tested statistically by using One Way ANOVA and Tukey HSD test. The result of the ANOVA test showed the number of sig $0.000 < 0.05$ which meant all four formulas were significantly different. The research concludes that the extract of arabica coffee leaf might affect the formulation of gel preparations and its extract concentration might affect SPF number on gel preparations.

1. Introduction

Skin does an important role to protect the body from outside environments such as physical impact and exposure to free radicals. In order to maximize the function of the skin as a protector, it is necessary to give special treatment on the skin. Amongst information about how dangerous UV rays affect our skin, it is also well-known that sunscreen has been trusted to protect our skin from the damage of UV rays[1].

UV rays are rays that are emitted by the sun that can reach the earth surface beside visible light and infrared rays. The wavelength of UV rays is approximately 200-400 nm. The UV spectrum, based on the wavelength, is categorized into three; UV C (200-290), UV B (290-320) and UV A (320-400). Not all of these three UV rays' radiation can reach earth's surface. UV C rays, the one with the strongest energy, cannot reach the surface because of being absorbed in ozone [2].

It is known that UV radiation from the sun causes some damages to human skin. UV A radiation is discovered to cause damages to skin cells and DNA. It also causes photoaging and photocarcinogenesis. These damages can be caused by long term exposure of the sun's rays, even in a small dose. UV A radiation contributes to the decreasing of skin elasticity, the increasing of wrinkles, and the increasing of free radicals that cause acute to chronic changes to skin. Besides, UV A radiation



also makes other diseases worsen, such as lupus erythematosus. It also contributes as immunosuppressants that increase cancer cell growth [3]. From another resource, it is stated that the exposure of ultraviolet radiation might cause epidermis damages, which is also called sunburn, pigmentation, wrinkling, premature skin aging, and long term exposure of the sun's rays can also cause mutation of lughture tissue in the stratum corneum[4]

On the other hand, the sun's rays stimulate melanin and pigment as natural sunscreen for human skin. The sun's rays also stimulate hormone protection which allows vitamin D synthesizing to increase skin cell regeneration. It is discovered that the shorter the wavelength, the stronger the energy level, which also causes more damage to the skin (Lu et al., 1996). If the sun's rays expose the skin excessively, it might not resist the negative effects. Therefore, it is necessary to give some protection by using sunscreen. Sunscreen is preparations applied on skin to absorb, to scatter, or to reflect UV rays. It is also applied to help natural protection mechanisms to protect skin from dangerous UV radiation of the sun's rays [6]. It is a product that provides protection from the damage of the sun's rays, but this product is commonly made of synthetic chemicals which allow, in long term exposure, some bad effects to human skin. This research is conducted, to test the formula, so that there would be a chance for natural ingredients to be natural protection which causes less bad effect [7]. The grade of sunscreen's photo-protective is universally measured by Sun Protective Factor (SPF) which determines the increase of the sun's rays' exposure dose with photo-protective products applied without eritema [8].

Coffee is categorized into genus *Coffea* in the family of Rubiaceae. Genus *Coffea* L. consists of more than 100 species, but only two species, *Coffea arabica* (arabica coffee) and *Coffea canephora* (robusta coffee), are commercially cultivated on large scales [9]. *Coffea arabica* is a tetraploid species from highland rain forest in the south of Ethiopia. Six of eight chosen species are closely related, including two commercial species *Coffea arabica* dan *Coffea canephora*, as illustrated in the recent phylogeny from genus [10].

Coffee leaf has not been optimally utilized. In Indonesia, coffee leaf is used as a brewed drink named "Aia Kawa" in Sumatera [11]. It is discovered that *Coffea arabica* leaf contains phenolic compounds with a concentration of 17.4% in green leaf and 13.9% in old/pure leaf (Rodrigues et al., 2008). Arabica coffee leaf contains some compounds, such as alkaloid, flavonoid, and phenol which are included into the derivative of hydroxycinnamic acid, caffeine, chlorogenic acid, coumarin, ferulic acid and cinnamic acid. Chlorogenic acid is easily oxidized in aqueous solutions [13] [14].

Phenolic and flavonoid compounds have potential as sunscreen [15]. Flavonoid is commonly able to absorb UV rays, but not able to absorb any radiation on wavelength around 280-315 nm maximally, as other phenylpropanoid compounds. However, compared with other phenylpropanoid compounds, flavonoid compounds will increase significantly, as it is exposed by UV B rays' radiation, including the derivative of hydroxycinnamic acid (for instance: p-Coumaric acid, ferulic acid and caffeine acid). Some research showed that when UV B rays' exposes plants, flavonoid concentration is increased compared with hydroxycinnamic compounds [16].

Sunscreen is applied to help natural protection mechanisms to protect skin from dangerous UV radiation of the sun's rays [6]. It is a product that provides protection from the damage of the sun's rays, but this product is commonly made of synthetic chemicals. In the market, sunscreen is provided in some forms of preparations; salve, cream, gel, lotion, spray, and wax stick [17]. Gel is a semi-solid preparation which consists of suspensions made of inorganic particles in small parts or organic molecules in big parts penetrated by some kinds of fluid [18].

In Indonesia, research related to arabica coffee leaves is limited. Therefore, it is necessary to conduct further research because the resources are overflowing and it should be utilized and developed, for instance, for cosmetic purposes. This research aims to fill the void in the literatures related to arabica coffee leaf by testing how the extract of arabica coffee leaf affects the formulation of gel preparations and discovering how far the extract of arabica coffee leaf concentration affects the grade of Sun Protective Factor of gel preparations.

2. Material and Methods

2.1 Material

The materials used in this study were arabica coffee leaves (*Coffea arabica* L.) as the active ingredient which was obtained by maceration using 70% ethanol. The chemicals used for the gel formulation were HPMC K100M type 2208 (High Viscosity Grade), Propylene Glycol (Pharmaceutical Grade), Methyl Paraben (Pharmaceutical Grade), Propyl Paraben (Pharmaceutical Grade), Aqua Dest, 70% Ethanol and 70% Ethanol for analysis.

2.2 Plant Determination

The determination of the arabica coffee plant used in the research was carried out at the Herbarium Bogoriense, Botany Division, Biological Research Center, LIPI Cibinong.

2.3 Arabica Coffee Leaf Extraction

Coarse powder of arabica coffee leaves that have been obtained from IPB Bogor was re-pollinated using a grinder or blender. It was then sieved using a mesh sieve no. 40. The extraction method used maceration. The maceration extraction method (cold method) is a better method for extracting cherry leaves compared to the soxhlation method (hot method). This is due to the thermolable nature of flavonoids which will be damaged when extracted using a soxhletation [19]. 1:10 fine powder and macerated solvent. A total of 1300 grams of powder was macerated with 9700 ml of 70% ethanol filter in a dark glass jar at room temperature. It was then left for 2 days and stirred every 8 hours. The maceration results were filtered with a flannel cloth and filter paper. Then, the remaining dregs were re-macerated with 3300 ml of sprinkler for 1 day. Afterwards, it was stirred and filtered again. The filtrate obtained was then concentrated using a rotary vacuum evaporator at a temperature of 50 ° C until a thick extract was obtained. Thick extract of arabica coffee leaves, and the yield was obtained. The yield was calculated based on the ratio of the final weight (weight of the extract produced) to the initial weight (weight of cell biomass used) multiplied by 100% [20].

2.4 Organoleptic Determination of Arabica Coffee Leaf Extract and Determination of Ash Content

Organoleptic determination of arabica coffee leaf extract was carried out by observing the shape, color, and odor of the arabica coffee leaf extract. 2 g of extract was weighed carefully and put in a porcelain crucible that had been annealed, tared, and leveled. Krus was slowly annealed until the charcoal ran out. Afterglow was carried out at a temperature of 600°C for 3 hours, which was then cooled and weighed until a fixed weight was obtained. The ash content was calculated against the material which was dried in the air. If in this way the charcoal cannot be removed, hot water can be added and filtered with an ash-free filter paper. The rest of the paper and filter paper was glazed in the same crucible. The filtrate was put into the crucible, and left to evaporate. Afterglow until the weight was fixed, weighed and calculated.

2.5 Phytochemical Screening Test

Extract was identified of alkaloid, flavonoid, phenol, saponin, tannin, steroid, and triterpenoid contents using Harbonne or Material Medica Method.

2.6 Gel Formulation

Arabica coffee leaf extract gel formula (*Coffea arabica* L.) consisted of arabica coffee leaf extract (*Coffea arabica* L.), HPMC K100M type 2208, Propylene Glycol, Methyl Paraben, Propyl Paraben and Aqua Dest. The gel formulation can be seen in table (1).

Table 1. Formulation Gel Arabica Coffee Leaf Extract

| No. | Materials | % (b/v) | | | | Functions |
|-----|------------------|---------|-------|-------|-------|----------------------|
| | | F1 | F2 | F3 | F4 | |
| 1 | Ekstrakt | - | 1.5 | 2 | 2.5 | Active ingredients |
| 2 | HPMC | 1 | 1 | 1 | 1 | <i>Gelling agent</i> |
| 3 | Propylene Glycol | 15 | 15 | 15 | 15 | Humectant |
| 4 | MethylParaben | 0.075 | 0.075 | 0.075 | 0.075 | Preservatives |
| 5 | Propylparaben | 0.025 | 0.025 | 0.025 | 0.025 | Preservatives |
| 6 | Aqua Dest ad | 100 | 100 | 100 | 100 | solvent |

2.7 Procedure for Preparation Arabica Coffee Leaf Extract Gel

HPMC 4.5 g was developed in a mortar with aqua dest and was then expanded for 24 hours and crushed to form a gel mass (M1). 0.3775 g of methyl paraben was dissolved with a portion of propylene glycol (M2). 0.1125 g of propyl paraben was dissolved with a portion of propylene glycol (M3). The arabica coffee leaf extract was also dissolved with the remaining propylene glycol (M4). M1, M2 and M3, M4 were mixed, crushed, and homogeneous. Furthermore, the evaluation of the determination of the FPS value of gel preparations were carried out.

2.8 Organoleptic Test for Gel

Organoleptic testing was carried out by direct observation of the shape, color and smell of the gel made. Gel was usually clear with a semi-solid consistency [18].

2.9 Homogeneity Test

The homogeneity test was carried out by applying 0.1 g of gel preparation to a piece of transparent glass and observing its homogeneity. The test preparation must show a homogeneous arrangement, indicated by the absence of coarse grains on the object glass [21].

2.10 pH test

The pH test was carried out by turning on the pH meter then dipping the pH meter electrode into the gel formula. It was left for a few moments until the pH meter screen showed a stable number [22].

2.11 Spreadability Test

As much as 1 gram of gel preparation was placed carefully on a glass measuring 20x20 cm, and was given a 125 gram weight on it. It was then measured the diameter formed after 1 minute [23].

2.12 Adhesion Test

A total of 1 gram of gel preparation was flattened on one glass object and covered with another glass object until the two plates were joined. The object glass pair was pressed with a load of 1000 g for 5 minutes. It was then installed on the adhesion test equipment, while simultaneously recording the time it took for the two plates to come off each other [24].

2.13 Viscosity and Flow Properties Test

The viscosity test was performed using the Brookfield Viscometer RV type. The gel preparation was put into a 500 mL beaker glass container, then 'the spindle no. 04 was installed. Afterwards, the spindle was lowered into the preparation to the specified limit. The tool was turned on and set at a speed

of 10 rpm, 12 rpm, 20 rpm, 30 rpm and 50 rpm to show a stable number. Then, the results were recorded. To determine the flow properties of the preparation, a curve was made between the shear rate and shear stress [21].

2.14 Centrifugation Test

10 grams of each preparation were put into a centrifugation tube and centrifuged at 3750 rpm for 5 hours. It was then observed whether there was separation or not in the gel [21].

2.15 Freeze Thaw Test

The phase separation cycle using the freeze thaw method on the gel preparation was carried out in 6 cycles for each formula. Each cycle was observed after 48 hours of storage at 4°C and 48 hours after at 45°C for 24 days. Each cycle was observed whether there was phase separation or not on the gel preparation [21].

2.16 Determination of the FPS Value of Arabica Coffee Leaf Extract

The FPS value of the extract in vitro was determined using the Uv-Vis spectrophotometric method developed by Mansur (1986) using equation 3. The extract was diluted with a concentration of 60 ppm, 80 ppm and 100 ppm with 70% pa ethanol and then measured using the Uv-Vis spectrophotometer at a wavelength of 290-320 nm. The determination of the FPS value was carried out in three replications. Then, the data obtained was processed with the Mansur equation (1986). To determine the FPS value of this wavelength, the EE x I value has been described in table 2 [7].

2.17 Determination of the FPS Value of Arabica Coffee Leaf Extract Gel Preparation

The FPS value of the extract in vitro was determined using the Uv-Vis spectrophotometric method developed by Mansur (1986) using equation 3. The preparation was weighed as much as 0.02 grams in 5 mL ethanol 70% pa and then measured using a Uv-Vis spectrophotometer at a wavelength 290-320 nm. The determination of the FPS value was carried out in three replications. Then, the data obtained was processed with the Mansur equation (1986). To determine the FPS value of this wavelength, the EE x I value has been described in table 2 [7].

The equation for determining the FPS value can be seen in below:

$$SPF = CF \times \sum_{290}^{320} EE(\lambda) \times I(\lambda) \times Abs(\lambda)$$

Based on this equation, there was a variable CF (Correction Factor) = 10, EE (Erythemogenic Effect), I is the simulated intensity of sunlight and Abs is the absorbance of the sample.

The value of EE x I can be seen in table 2.

Table 2. The Value of EE x I

| Wave Length (nm) | EE x I |
|------------------|--------|
| 290 | 0.0150 |
| 295 | 0.0817 |
| 300 | 0.2874 |
| 305 | 0.3278 |
| 310 | 0.1864 |
| 315 | 0.0839 |
| 320 | 0.0180 |

According to the FDA (Food Drug Administration), the distribution of sunscreen capabilities is Minimum (SPF is between 2-4), Medium (SPF is between 4-6), Extra (SPF is between 6-8), Maximum (SPF is between 8-15), and Ultra (SPF is more than 15) [25].

2.18 Statistical Analysis

Qualitative testing of physical characteristics and determination of the FPS value of gel preparations were analyzed using One Way ANOVA, with a confidence level of 95% ($p > 0.05$). If there WERE significant differences between formulas, we then proceeded with the Tukey HSD test to see which formulas had significant differences.

3. Results and Discussion

The object of this research was fresh arabica coffee plants (*Coffea arabica* L.). Then, the fresh arabica coffee plants were determined in Herbarium Bogoriense, Botany Department of Biology Research Center, LIPI Cibinong. The determination of the plants in this research meant to find out the clear identity of the research object to avoid mistakes in collecting main material for the research. After determined, it was confirmed that the plants were arabica coffee leaf (*Coffea arabica* L.).

Table 3. The Results of Characterization pf Arabica Coffee Leaf Extract

| Examination | Results |
|------------------|-----------------|
| yield | 9.84% |
| Organoleptic: | |
| a. form | Thick extract |
| b. smell | coffee |
| c. taste | bitter |
| d. color | Chocolate black |
| Drying shrinkage | 6.61% ± 0.43 |
| Ash content | 9.26% ± 0.46 |

Based on the results of phytochemical screening tests, 70% thick ethanol extract of arabica coffee leaves had alkaloid, phenolic, flavonoid, saponin, tannin and steroid compounds. This is in accordance with previous research [26], [27]. By knowing the truth of its benefits, arabica coffee leaf extract can be formulated into gel preparations that contain antioxidants.

Table 4. Results of Phytochemical Screening Test

| Compound | Result |
|--------------|--------|
| Phenolic | + |
| Alkaloids | + |
| Flavonoids | + |
| Saponins | + |
| Tannins | + |
| Steroids | + |
| Triterpenoid | - |

Organoleptic tests included observations on the consistency, color and odor of the gel preparation. The results of organoleptic testing can be seen in table 5. In accordance with the results from the organoleptic test, it was found that all four formulas had semi-solid consistency. For F1, the gel was clear/transparent and odorless because it was not mixed with the extract, while for F2, F3, and F4, they were clear brown and had a typical smell from the extract. The extract concentration affected the color of gel preparations, the higher the concentration, the deeper the brown color.

Homogeneity test aimed to determine the homogeneity of preparations that have been made, homogeneous preparations will produce good quality because it showed the active ingredients of the drug are evenly dispersed in the basic ingredients. Based on the research results, the four formulas produced a homogeneous gel.

Table 5. Results of Organoleptic Tests and Homogeneity Tests

| Formula | Organoleptic | | | |
|---------|--------------|-------------|----------|-------------|
| | Consistency | Color | Smell | Homogeneity |
| 1 | Semisolid | Transparent | Odorless | Homogeneous |
| 2 | Semisolid | Clear brown | Unique | Homogeneous |
| 3 | Semisolid | Clear brown | Unique | Homogeneous |
| 4 | Semisolid | Clear brown | Unique | Homogeneous |

The purpose of this test was to determine that the resulting gel was acceptable for skin pH, because if it is not in accordance with the pH of the skin it can cause irritation. Based on the results of the study, the four formulas met the pH criteria on sunscreen preparations. pH test discovered that all formulas were qualified for pH criterias on sunscreen preparations, which was around 4.5-8.0 (SNI, 1996). The extract concentration affected the acid of the preparations because the extract contained a flavonoid compound which was quite acidic. Therefore, the more the arabica coffee leaf extract concentration, the more the acid it has (Markham, 1988). Arabica coffee leaf extract was 6.16 pH. A product that had too high or too low pH could cause irritation on skin [6]. The graph of the pH test results can be seen in Figure 1.




Figure 1. The Results of pH test

The dispersion test was carried out to ensure that the semisolid preparations are able to spread easily without significant pressure, so that they are easy to apply without causing pain and provide comfort to the user. The graph of the results of the scattering power test can be seen in Figure 2. The greater the spreadability of the preparation, the greater the ability of the active substance to spread and contact with the skin. The spreadability is inversely proportional to viscosity, the greater the viscosity of a preparation, the thicker the consistency, so that the resulting spreadability is smaller [30].




Figure 2. The Results of Spreadability Test

The adhesion test was carried out to determine the ability of the gel to adhere to the skin surface. The graph of the adhesion test results can be seen in Figure 3. Based on the research results, the four formulas met the adhesion criteria. Topical preparations must have sufficient adhesiveness but should not be sticky to the skin as they can reduce comfort during use.

The viscosity test was aimed to determine the thickness of the gel preparation. The higher the viscosity value, the higher the viscosity level of the gel preparation. Viscosity was tested by using *Brookfield* Viscometer RV type. The result revealed that all formulas were qualified for criterias of gel viscosity, which was 2000-4000 *cPs*. [31]. Gel consistency, which was not too liquid or too thick, was a characteristic of the good gel. HPMC concentration in gel preparations was used only 1%, as HPMC K100M is categorized into *High Viscosity Grade* which can be read in *Certificate of Analysis*. Gel viscosity affects the gel itself to disperse and adhere to skin. The higher the viscosity, the more difficult the gel to disperse although it is easier to adhere on skin. The higher the viscosity, the smaller the separation rate of dispersed phase [32]. Of the four formulas, the gel viscosity criteria were met. The graph of the results of the viscosity test can be seen in Figure 4.

All four formulas produced plastic thixotropic flow. It could be seen that the curve was started without getting through point zero (0) but cutting the axis of motion stress (or the outside of the curve was extrapolated cutting the axis) at the certain points named yield grade [33]. It was also shown that the decreased curve was to the left of the increased curve because there was structural change that did not immediately return to the first condition, if the pressure was diminished. Yield grade resulted from F1 was $f = 3.496$ dyne/cm², F2 was $f = 3.097$ dyne/cm², F3 was $f = 2.979$ dyne/cm² and F4 was $f = 3.557$ dyne/cm². Thixotropic is the characteristic of the flow that is expected for semi-solid preparations because it is expected to be in high consistency in a container. Nevertheless, it is easier to pour when it is given force, making it easier to be applied on skin [34]




Figure 3. The Results of Viscosity Test




Figure 4. Flow Properties Formula 1




Figure 5. Flow Properties Formula 2




Figure 6. Flow Properties Formula 3




Figure 7. Flow Properties Formula 4

The centrifugation test aimed to observe whether there is a phase separation of the gel preparation and to see the stability of the gel preparation after very strong shaking. Based on the research results, the four formulas did not undergo phase separation or syneresis. The table of centrifugation test results can be seen in table 6.

Table 6. Centrifugation Test

| Formula | Results | |
|---------|---------|---|
| F1 | 1 | - |
| | 2 | - |
| | 3 | - |
| F2 | 1 | - |
| | 2 | - |
| | 3 | - |
| F3 | 1 | - |
| | 2 | - |
| | 3 | - |
| F4 | 1 | - |
| | 2 | - |
| | 3 | - |

Note : (-) no separation phase

The freeze thaw test is a test that has the purpose of simulating the product during the distribution process in vehicles, which are rarely equipped with temperature control devices. Therefore, this test was carried out at a certain temperature or humidity at a certain time so that the product in its packaging will undergo various changes. If during the six cycles, the preparation is stable or there is no phase that changes, it means that the product is stable during the distribution process [35]. Based on the results of the freeze thaw test, the four formulas were stable as there was no phase separation. The table of freeze thaw test results can be seen in table 7.

Table 7. Results of *Freeze Thaw*

| cycle | F1 | | | F2 | | | F3 | | | F4 | | |
|--------|----|---|---|----|---|---|----|---|---|----|---|---|
| | 1 | 2 | 3 | 1 | 2 | 3 | 1 | 2 | 3 | 1 | 2 | 3 |
| 1 4°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 1 45°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 2 4°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 2 45°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 3 4°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 3 45°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 4 4°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 4 45°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 5 4°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 5 45°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 6 4°C | - | - | - | - | - | - | - | - | - | - | - | - |
| 6 45°C | - | - | - | - | - | - | - | - | - | - | - | - |

Note: (-) no separation phase

The Sun Protection Factor (SPF) value or FPS value is measured as the ability or effectiveness of a material as a sunscreen. The higher the FPS value, the better the sunscreen will protect against UV rays. The purpose of using sunscreen is based on its ability to absorb, reflect or scatter sunlight [36]. The value of FPS was determined in vitro using the UV-vis spectrophotometry method developed by Mansur (1986). It was done by measuring the absorbance of each sample, namely the extract of arabica coffee leaves and the four formulas for the gel preparation for three replications with a wavelength between 290 - 320 nm where the measurements were described for each 5 nm interval. It is known that the arabica coffee leaf extract has these groups found in flavonoids and phenolic compounds, making it able to absorb radiation energy and produce absorbance values on the uv-vis spectrophotometer.

Based on the results of the study, the arabica coffee leaf extract has a concentration of 60 ppm. Of the three extract concentrations, it has a minimal protection category. The F1 gel preparation was a negative control, which did not contain extracts but produced a low FPS value for sunscreen so that it did not fall into the protection category. F2, F3 and F4 have a minimal protection category. The results of determining the FPS value of extracts and gel preparations can be seen in table 8 and table 9. From the second to fourth formulas, all formulas had minimal protection categories. From the results, it was obtained that gel preparations produce low protection on the skin. This can be overcome by adding other active sunscreen ingredients to support the effectiveness of sunscreens to produce more maximally.

On the SPF grade of the gel preparation that was higher than the extract, from the results of the research of [38] Donglikar and Deore (2017), it was revealed that the phytochemical SPF of the formulation was higher, which indicated the synergy and compatibility of the excipients as well. This result indicated that the prepared formulation has good SPF and good sun protection activity. Moreo et al., (2013) explained that this is possible to happen because it is influenced by the combination and concentration of the gel carrier component, the type of gel, the effect and the interaction of the carrier component such as the humectant used in the formulation. This factor can increase or decrease the UV absorption of sunscreens [39].

Therefore, to develop sunscreens with better safety and high SPF, the formulator must understand the physicochemical principles of not only active UV absorbance, but also vehicle components, such as esters, emulsifiers and emulsifiers used in formulations. This is because sunscreens can interact with other components of the vehicle, and this interaction can affect the efficacy of the sunscreen.

Table 8. Results of SPF Value Extract

| Sample | FPS Value | Protection Category |
|------------------|-----------|---------------------|
| Ekstrakt 60 ppm | 2.08 | Minimum protection |
| Ekstrakt 80 ppm | 2.49 | Minimum protection |
| Ekstrakt 100 ppm | 3.68 | Minimum protection |

Table 9. Results of SPF Value Gel Extract

| Sample | Value SPF | Protection Category |
|-----------|-----------|---------------------|
| Formula 1 | 0.61 | - |
| Formula 2 | 2.59 | Minimum protection |
| Formula 3 | 2.95 | Minimum protection |
| Formula 4 | 3.64 | Minimum protection |

4. Conclusions

Based on the research results, arabica coffee leaf extract could affect the gel dosage formulation and meet the specified requirements. The results of the ANOVA test showed a grade significance of $0.000 < 0.05$, where there were significant differences in the four formulas for gel preparations. This showed that the concentration of arabica coffee leaf extract could affect the SPF value of the gel preparation. In this study, a gel preparation formulation that met the pharmaceutical requirements was obtained by providing an SPF grade with minimal protection.

References

- [1] H. Winarsi, *Isoflavon*. Gadjah Mada University, 2005.
- [2] COLIPA, *Sun Protection Test Methods*. COLIPA guidelines international, 2006.
- [3] K. M. Yuliawati, E. R. Sadiyah, R. Solehati, and A. Elgiawan, "Sunscreen Activity Testing of Robusta Coffee (*Coffea canephora* ex Froehner) Leave Extract and Fractions Pengujian Aktivitas Tabir Surya Ekstrak dan Fraksi Daun Kopi Robusta (*Coffea canephora*)," *Indones. J. Pharm. Sci. Technol.*, vol. 1, no. 1, 2019.
- [4] D. RI, *Formularium Kosmetika Indonesia*. Jakarta: Departemen Kesehatan Republik Indonesia, 1985.
- [5] M. R. Lu H, Edwards C, Pearse A, "Melanin content and distribution in the surface corneocyte with phototypes," *Br J Dermatol*, vol. 135, pp. 263–267, 1996.
- [6] A. D. Puspitasari, D. A. K. Mulangsri, and H. Herlina, "Formulasi Krim Tabir Surya Ekstrak Etanol Daun Kersen (*Muntingia calabura* L.) untuk Kesehatan Kulit," *Media Penelit. dan Pengemb. Kesehat.*, vol. 28, no. 4, pp. 263–270, 2018, doi: 10.22435/mpk.v28i4.524.
- [7] W. A. d. S. Almeida *et al.*, "Photoprotective activity and increase of SPF in sunscreen formulation using lyophilized red propolis extracts from Alagoas," *Rev. Bras. Farmacogn.*, vol. 29, no. 3, pp. 373–380, 2019, doi: 10.1016/j.bjp.2019.02.003.
- [8] M. C. Gonçalves *et al.*, "Preliminary studies of the cytotoxicity and photoprotective properties of benzophenone and lactone derivatives," *Rev. Virtual Quim.*, vol. 10, no. 3, pp. 600–608, 2018, doi: 10.21577/1984-6835.20180044.
- [9] A. P. Davis, R. Govaerts, D. M. Bridson, and P. Stoffelen, "An annotated taxonomic conspectus of the genus *Coffea* (Rubiaceae)," *Bot. J. Linn. Soc.*, vol. 152, no. 4, pp. 465–512, 2006, doi: 10.1111/j.1095-8339.2006.00584.x.
- [10] H. L. . Hamon, P.; Grover, C.E.; Davis, A.P.; Rakotomalala, J.-J.; Raharimalala, N.E.; Albert, V.A.; Sreenath and E. Stoffelen, P.; Mitchell, S.E.; Couturon, "Genotyping-by-sequencing provides the first well-resolved phylogeny for coffee (*Coffea*) and insights into the evolution of

- caffeine content in its species: GBS coffee phylogeny and the evolution of caffeine content.," *Mol. Phylogenet. Evol.*, vol. 109., pp. 351–361, 2017.
- [11] D. Ristiana, "Aktivitas Antioksidan Dan Kadar Fenol Berbagai Ekstrak Daun Kopi (*Coffea Sp.*): Potensi Aplikasi Bahan Alami Untuk Fortifikasi Pangan," *J. Apl. Teknol. Pangan*, vol. 6, no. 2, pp. 89–92, 2017, doi: 10.17728/jatp.205.
- [12] O. Rodrigues Salgado, P.; Favarin, J.L.; Leandro, R.A.; Fontão de Lima Filho, "Total phenol concentrations in coffee tree leaves during fruit development," *Sci. Agric.*, vol. 65, pp. 354–359, 2008.
- [13] C. Manach, A. Scalbert, C. Morand, C. Rémésy, and L. Jiménez, "Polyphenols: Food sources and bioavailability," *Am. J. Clin. Nutr.*, vol. 79, no. 5, pp. 727–747, 2004, doi: 10.1093/ajcn/79.5.727.
- [14] I. Tomac and M. Šeruga, "Electrochemical properties of chlorogenic acids and determination of their content in coffee using differential pulse voltammetry," *Int. J. Electrochem. Sci.*, vol. 11, no. 4, pp. 2854–2876, 2016, doi: 10.20964/110402854.
- [15] Shovyana HH; A Karim Zulkarnain, "Physical Stability and Activity of Cream W/O Etanolic Fruit Extract of Mahkota Dewa (*Phaleria macrocarpha* (scheff.) Boerl.) as A a Sunscreen," *YogyakartaFakultas Farm. UGM. Tradit. Med. J.*, vol. 18, no. 2, 2013.
- [16] G. Agati and M. Tattini, "Multiple functional roles of flavonoids in photoprotection," *New Phytol.*, vol. 186, no. 4, pp. 786–793, 2010, doi: 10.1111/j.1469-8137.2010.03269.x.
- [17] Fields, "Sunscreens:Mechanisms of Action, Use, and Excipients," *Int. J. Pharm. Compd.*, 2008.
- [18] A. P. Baskar M, Kiranmathyi B, Sivaraj C, Saraswathi K, "Journal of drug delivery and therapeutics (jddt)," *J. Drug Deliv. Ther.*, vol. 9, no. 3, pp. 661–668, 2019, [Online]. Available: <http://dx.doi.org/10.22270/jddt.v9i3.2678>.
- [19] A. D. Puspitasari and R. L. Wulandari, "Aktivitas Antioksidan dan Penetapan Kadar Flavonoid Total Ekstrak Etil Asetat Daun Kersen (*Muntingia calabura*)," *J. Pharmascience*, vol. 4, no. 2, pp. 167–175, 2017, doi: 10.20527/jps.v4i2.5770.
- [20] R. N. Sani, F. C. Nisa, R. D. Andriani, and J. M. Maligan, "ANALISIS RENDEMEN DAN SKRINING FITOKIMIA EKSTRAK ETANOL MIKROALGA LAUT *Tetraselmis chuii* Yield Analysis and Phytochemical Screening Ethanol Extract of Marine Microalgae *Tetraselmis chuii*," *J. Pangan dan Agroindustri*, vol. 2, no. 2, pp. 121–126, 2014.
- [21] H. C. A. Loyd V. Allen, Nicholas G. Popovich, "Ansel 's Pharmaceutical Dosage Forms and," *Int. J. Pharm. Compd.*, p. 537, 2014, [Online]. Available: https://downloads.lww.com/wolterskluwer_vitalstream_com/sample-content/9780781779340_allen/samples/frontmatter.pdf.
- [22] S. Septiani, N. Wathoni, and S. R. mita Mita, "Formulasi Sediaan Masker gel Antioksidan Dari Ekstrak Etanol Biji Belinjo," *Fak. Farm. Univ. Padjajaran*, pp. 2–4, 2011.
- [23] P. Niyogi, N. Raju, P. Readdy, and B. Rao, "Formulation And Evaluation Of Antiinflammatory Activity Of *Solanum Pubescens* Wild Extracts Gel On Albino Wistar Rats," vol. 2, no. 3, pp. 484–490, 2012.
- [24] L. Allen, *The art, Science, and Technology of Compounding*, Second Edi. Washington DC: American Pharmaceutical, 1998.
- [25] V. Damogalad, H. Jaya Edy, and H. Sri Supriati, "Formulasi Krim Tabir Surya Ekstrak Kulit Nanas (*Ananas Comosus* L Merr) Dan Uji in Vitro Nilai Sun Protecting Factor (Spf)," *PHARMACON J. Ilm. Farm. – UNSRAT*, vol. 2, no. 02, pp. 2302–2493, 2013.
- [26] M. Rahmah Nasution and M. Br Manullang, "AKTIVITAS ANTIOKSIDAN SEDUHAN DAUN KOPI KAWA KERING (*Coffea arabica* L) DENGAN METODE DPPH," *J. Insa. Farm. Indones.*, vol. 3, no. 1, pp. 114–123, 2020, doi: 10.36387/jifi.v3i1.467.
- [27] Y. Retnaningtyas, N. Kristiningrum, H. D. Renggani, and N. P. Narindra, "Karakteristik Simplisia dan Teh Herbal Daun Kopi Arabika (*Coffea arabica*)," *Farm. Jember*, vol. 1, no. 1, pp. 46–54, 2016.

- [28] Badan Standardisasi Nasional, "Sediaan Tabir Surya," *Dewan Stand. Nas.*, vol. 16, no. 4399, pp. 1–3, 1996.
- [29] Markham KR, *Cara Mengidentifikasi Flavonoid*. Bandung: Penerbit ITB, 1988.
- [30] Nutrisia Aquariushinta Sayuti, "Formulasi dan Uji Stabilitas Fisik Sediaan Gel Ekstrak Daun Ketepeng Cina (*Cassia alata* L.)," *J. Kefarmasian Indones.*, vol. 5, no. 2, pp. 74–82, 2015.
- [31] A. Garg, "Spreading of semisolid formulation," USA, pp. 84–102, Sep. 2002.
- [32] N. S. Daud, N. S. Daud, L. Ode, and Z. Al, "FORMULASI LOTION TABIR SURYA EKSTRAK ETANOL BERAS MERAH (*Oryza nivara*)," vol. 1, no. September 2016, pp. 143–150.
- [33] Agoes, *Sediaan Farmasi Likuida-Semisolida*. Bandung: Intitut Teknologi Bandung, 2012.
- [34] U. Shinde, S. Pokharkar, and S. Modani, "Design and evaluation of microemulsion gel system of nadifloxacin," *Indian J. Pharm. Sci.*, vol. 74, no. 3, pp. 237–247, 2012, doi: 10.4103/0250-474X.106066.
- [35] S. Bajaj, D. Singla, and N. Sakhuja, "Stability testing of pharmaceutical products," *J. Appl. Pharm. Sci.*, vol. 2, no. 3, pp. 129–138, 2012, doi: 10.7324/JAPS.2012.2322.
- [36] N. Sharon, S. Anam, and Yuliet, "Formulasi Krim Antioksidan Ekstrak Etanol Bawang Hutan (*Eleutherine palmifolia* L. Merr.)," *Online J. Nat. Sci.*, vol. 2, no. 3, pp. 111–122, 2013.
- [37] B. M. Mansur JDS, "Determinação do fator de proteção solar por espectrofotometria," *Mansur MCDA*, 1986.
- [38] M. M. Donglikar and S. L. Deore, "Development and evaluation of herbal sunscreen," *Pharmacogn. J.*, vol. 9, no. 1, pp. 83–97, 2017, doi: 10.5530/pj.2017.1.15.
- [39] More BH; Sakharwade SN; Tembhone S; Sakarkar DM, "Evaluation of Sunscreen activity of Cream containing Leaves Extract of *Butea monosperma* for Topical application," *Int. J. Res. Cosmet. Sci.*, vol. 3, no. 1, pp. 1–6, 2013.