

HAYATI

p-ISSN: 1978-3019
e-ISSN: 2086-4094

Journal of Biosciences

[ABOUT](#) [CURRENT](#) [ARCHIVES](#) [EDITORS](#) [SUBMISSIONS](#) [REGISTER](#) [CONTACT](#) [SEARCH](#) [Search](#)

HAYATI Journal of Biosciences (HAYATI J Biosci; **p-ISSN: 1978-3019; e-ISSN: 2086-4094**) is an international peer-reviewed and open access journal that publishes significant and important research from all area of biosciences fields such as biodiversity, biosystematics, ecology, physiology, behavior, genetics and biotechnology. All life forms, ranging from microbes, fungi, plants, animals, and human, including virus, are covered by HAYATI J Biosci.

HAYATI J Biosci published by Department of Biology, Bogor Agricultural University, Indonesia and the Indonesian Society for Biology. We accept submission from all over the world. Our Editorial Board members are prominent and active international researchers in biosciences fields who ensure efficient, fair, and constructive peer-review process. All accepted articles will be published on payment of an article-processing charge, and will be freely available to all readers with worldwide visibility and coverage.

HAYATI J Biosci has been also indexed/registered in SCOPUS, Crossref, DOAJ, CABI, EBSCO, Agricola and ProQuest. From October 2015-2017, HAYATI was hosted by Elsevier and available free of charge through Science Direct.

HAYATI J Biosci Indexed by Scopus

2017-04-14



Starting May 2017, HAYATI J Biosci is included in SCOPUS database/index. [Read More](#) >

Current Issue

Vol. 30 No. 2 (2023): March 2023



Published: 2023-01-06

Short Communication

Population Estimate of Long-Tailed Macaques (*Macaca fascicularis*) on Tinjil Island

Dyah Perwitasari-Farajallah, Entang Iskandar, Hana Intishar Sawitri, Tommy Langgeng Abimanyu, Vallen Sakti Maulana, Annisa Devi Rachmawati, Indra Purnama, Huda S. Darusman
193-197



PDF

First Record of The Thecate Marine Dinoflagellate *Protoperidinium abei* (Peridiniales, Dinophyceae) in Indonesia

Danang Ambar Prabowo, Debora Christin Purbani, Diah Radini Noerdjito
207-215



PDF

Articles

Increasing The Stocking Density of Grouper Nurseries for Aquabusiness Efficiency in Recirculating Aquaculture System (RAS) with Bioremediation

Belinda Astari, Tatag Budiardi, Suko Ismi, Irzal Effendi, Yani Hadiroseyani
198-206



PDF

In Silico Analysis of Protein of Milk, Soybean, and Kefir as Anti- Thrombotic Bioactive Peptide

Muhammad Ilham Fahri, Fadilla Sherlyna, Mochamad Nurcholis
216-223



PDF

Evaluation of Nutrient Content and Antioxidant Activity of Wood Ear Mushroom (*Auricularia auricula-Judae*) in the Addition of Reeds (*Imperata cylindrica* (L.) Beauv) as a Cultivation Medium

Sotya Rawiningtyas, Adi Setyo Purnomo, Sri Fatmawati
224-231



PDF

Revisit Study of Freshwater Sponges *Eunapius carteri* (Bowerbank, 1863) and a New Record of *Oncosclera asiatica* Manconi and Ruengsawang, 2012 (Porifera: Spongillida) in Porong River, East Java, Indonesia

Setiawan Edwin, Yanuar Ahmad, Michael Einstein Hermanto, Riani Catur , Prayogo Fitra Adi , Budiharjo Anto
232-245



PDF

Metabolites Profiling of *Penicillium citrinum* Recovered from Endophytic of Ramie (*Boehmeria nivea*) as a Potential Biocontrol Against Pathogenic Fungi

Kartika Sari, Vira Kusuma Dewi, Asri Peni Wulandari, Nia Rossiana, Tati Herlina, Fitri Widiyanti
246-255



PDF

Reef Fish Biodiversity at Different Depths in Tunda Island, Banten Province, Indonesia

Mujiyanto Mujiyanto, Amran Ronny Syam, Sasanti R. Suharti, Yayuk Sugianti, Saras Sharma
256-270



PDF

Paederia foetida Linn Leaves-Derived Extract Showed Antioxidant and Cytotoxic Properties Against Breast Carcinoma Cell

Jepri Agung Priyanto, Muhammad Eka Prastya, Gian Primahana, Ahmad Randy, Diyah Tri Utami
271-280



PDF

In Silico Analysis of Gene Expression Location and Single Nucleotide Polymorphisms (SNPs) of The Glucagon Like Peptide 1 Receptor (GLP-1R)

Rita Maliza, Putra Santoso, Bramadi Arya, Alimuddin Tofrizal, Rarastoeti Pratiwi

281-291



Genotype-Environment Interaction and Yield Stability of Upland Rice in Intercropping Cultivation

Aris Hairmansis, Supartopo, Yullianida, Nafisah, Rini Hermanasari, Angelita Puji Lestari, Suwarno

292-301



Growth of Red Amaranth (*Amaranthus cruentus* L.) Cultivated on Soil-Based Substrate Amended with a Residue of the Black Soldier Fly Larvae Containing Heavy Metals

Hana Haruna Putri, Benyamin Lakitan, Zaidan Panji Negara, Strayker Ali Muda

302-312



Biodefluorination of Perfluorooctanesulphonate by *Ensifer adhaerens* M1

Sharipov Danil, Starikov Sergey, Chetverikov Sergey

313-320



Morphological and Anatomical Comparison between Tetraploid *Stevia rebaudiana* (Bertoni) Bertoni and its Parental Diploid

Rifatul Adabiyah, Diah Ratnadewi, Tri Muji Ermayanti, Erwin Al Hafizh, Elsa May Susanti

321-335



The Changes in Seed Germination Capacity, Seedling Growth, and Leaf Morphology of *Ficus variegata* Blume Influenced by Gamma-Ray Irradiation

Muhammad Zanzibar, Agus Astho Pramono, Yulianti, Kurniawati Purwaka Putri, Dede J. Sudrajat, Naning Yuniarti, Nina Mindawati, Dida Syamsuwida, Danu, Ratna Uli Damayati Sianturi, Y. M.M. Anita Nugraheni, Nurhasybi, Junaedi Ahmad

336-346



Protein Hydrolysate of Green Peas Bromelain Attenuates Kidney Fibrosis in Cisplatin-Induced Nephrotoxicity Rats: Emphasis on Anti-Inflammatory Activities

Meilinah Hidayat; Retno Pangastuti, Sijani Prahastuti; Khomaini Hasan

347-357



A Meta-Analysis of Latex Physiology Studies Reveals Limited Adoption and Difficulties to Interpret Some Latex Diagnosis Parameters in *Hevea brasiliensis*

Junaidi, Anne Clément-Vidal, Tri Rini Nuringtyas, Eric Gohet, Siti Subandiyah, Pascal Montoro

358-371



A Possible Mixing Genetic Pool of Eastern Little Tuna *Euthynnus affinis* Cantor (1849) from the Western and Southern Part of Indonesian Waters

Bram Setyadji, Maya Agustina, Raymon Rahmanov Zedta, Ririk Kartika Sulistyaningsih, Putu Viby Indriani, Daniar Kusumawati

372-379



A Study of Chemical Constituents in Platinum Fast-Grown Teak Wood (*Tectona grandis*) with Age Differences Using Py-GCMS Coupled with Interdependence Multivariate Analysis

Maya Ismayati, Dwi Ajas Pramasari, Wahyu Dwianto, Danang Sudarwoko Adi, Nyndia Tri Muliawati, Ratih Damayanti, Narita Ayu Putri Pramesti, Syahrul Ramadhan, Ari Hardianto, Nadia Nuraniya Kamaluddin

380-391




Effect of Different Enriched Crickets (*Gryllus sigilatus*) on Growth and Pigmentation of Asian Arowana (*Scleropages formosus*) Var. Super Red

Sukarman Sukarman, Rendy Ginanjar, Mochammad Zamroni, Idil Ardi, Ahmad Musa, Melta Rini Fahmi, Agus Priyadi, Muhamad Yamin
392-403



[View All Issues >](#)



JOURNAL INFORMATION

[About The Journal](#)

[Editorial Team](#)

[Focus and Scope](#)

[Publication Ethics](#)

[Privacy Statement](#)

[Author Guidelines](#)

[Register New Account](#)

[Submit or Check Status](#)

[Contact](#)



INDEXING





PUBLISHER



Department of
Biology



GOOGLE SCHOLAR CITATIONS

HAYATI J.Biosci : GS citation		
	All	Since 2018
Citations	8186	5879
h-index	41	34
i10-index	255	193
<div> <div>GS citation</div> <div>></div> <div><</div> </div>		

VISITOR STATISTICS

00576932

[View Hayati J. Biosci. Stats](#)



HJB Most Read Papers

[Propolis Components and Biological Activities from Stingless Bees Collected on South Sulawesi, Indonesia](#)

84

1 January 2020

Ryo Miyata, Muhamad Sahla...

[Photosynthesis and Transpiration Rates of Rice Cultivated Under the System of Rice Intensification and the Effects on Growth and Yield](#)

76

10 November 2016

Nurul Hidayati, . Triadiati...

[Localization of Alkaloid and Other Secondary Metabolites in Cinchona ledgeriana Moens: Anatomical and Histochemical Studies on Fresh Tissues and Cultured Cells](#)

75

1 January 2020

Dian Rahma Pratiwi, Yohana...

[The Development of In Vitro Culture Sterilization Method of Gametophyte Explant Lopholejeunea sp.](#)

60

1 April 2021

Anna Widyastuti, Afiatry ...

[Antibacterial and Antioxidant Activities of Fungal Endophytes Isolated from Medicinal Plants in Simeulue Island, Aceh](#)

60

28 June 2022

Ahmad Fathoni, Muhammad Ily...



This journal is published under the terms of the <https://creativecommons.org/licenses/by-nc/4.0/>



IPB University
— Bogor Indonesia —

HAYATI Journal of Biosciences (p-ISSN: 1978-3019; e-ISSN: 2086-4094)

Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Dramaga Campus, Bogor 16680, Indonesia

Phone/Fax: +62-251-8421258; E-mail: hayati.jbiosci@apps.ipb.ac.id; URL : <http://journal.ipb.ac.id/index.php/hayati>

Platform &
workflow by
OJS / PKP

HAYATI


p-ISSN: 1978-3019

e-ISSN: 2086-4094

Journal of Biosciences

[ABOUT](#)[CURRENT](#)[ARCHIVES](#)[EDITORS](#)[SUBMISSIONS](#)[REGISTER](#)[CONTACT](#)[SEARCH](#) [Home](#) / [Archives](#) / Vol. 29 No. 4 (2022): July 2022

Vol. 29 No. 4 (2022): July 2022

**Published:** 2022-05-11

Articles

Decolorization of Synthetic Dyes by Tropical Fungi Isolated from Taman Eden 100, Toba Samosir, North Sumatra, Indonesia

Sita Heris Anita, Fenny Clara Ardiati, Kharisma Panji Ramadhan, Raden Permana Budi Laksana, Fahriya Puspita Sari, Oktan Dwi Nurhayat, Dede Heri Yuli Yanto

417-427

[PDF](#)

Evaluation of Cardiovascular Biomarkers and Lipid Regulation in Lactation Friesian Holstein at Different Altitude in West Java, Indonesia

Ujang Hidayat Tanuwiria, Iin Susilawati, Didin Tasripin, Lia Budimulyati Salman, Andi Mushawwir

428-434

[PDF](#)

Ethanol Productivity of Ethanol-Tolerant Mutant Strain *Pichia kudriavzevii* R-T3 in Monoculture and Co-culture Fermentation with *Saccharomyces cerevisiae*

Nurfadhilla Rahmadhani, Rika Indri Astuti, Anja Meryandini

435-444

[PDF](#)

Adaptive Mangrove Ecosystem Rehabilitation Plan based on Coastal Typology and Ecological Dynamics Approach

Aswin Rahadian, Cecep Kusmana, Yudi Setiawan, Lilik Budi Prasetyo

445-458



PDF

Correlation of Heavy Metals Cd, Cr, Cu, Hg, Pb, and Zn with Intestinal Bacteria in *Anas platyrhynchos* L. Duck

R Susanti, Ari Yuniastuti, Muchamad Dafip, Fidria Fibriana

459-467



PDF

Production, Characterization, and Molecular Phylogenetic Analysis of Phytase from *Aspergillus niger* Isolates of an Indonesia Origin

Baraka Stewart Mkumbe, Sajidan Sajidan, Artini Pangastuti, Ari Susilowati, Ralf Greiner

468-478



PDF

***Gynura procumbens* Adventitious Root Extract Altered Expression of Antioxidant Genes and Exert Hepatoprotective Effects Against Cadmium-Induced Oxidative Stress in Mice**

Sugiharto, Dwi Winarni, Anjar Tri Wibowo, Ufairanisa Islamatasya, Idqa Nurtri Bhakti, Nabilatun Nisa, Boon Chin Tan, Yosephine Sri Wulan Manuhara

479-486



PDF

Morphology and Anatomy of Endemic Fish *Leptobarbus melanopterus* (Cyprinidae) in Danau Sentarum National Park Kapuas Hulu Regency

Ari Hepi Yanti, Tri Rima Setyawati, Barrata Barrata

487-498



PDF

Overexpression of Heat Shock Cognate Protein 71 kDa and Pyruvate Dehydrogenase in the Brain Tissue at the Early Stage of High Fat Diet Consumption

Yan-Fen Lee, Xuan-Yi Sim, Mohd Nazri Ismail, Lay-Harn Gam

499-506



PDF

Genetic Diversity and Population Structure of Bullet Tuna (*Auxis rochei*) from Bali and Its Adjacent Waters

Maya Agustina, Bram Setyadji, Made Pharmawati, I ketut Junitha

507-514



PDF

***Nicotiana tabacum* var. Virginia Bio Oil-based Pyrolysis Extraction Have Prominence Antimicrobial Potential Compared to Ethanol Heat Reflux Extraction (EHRE)**

Andri Pramesyanti Pramono, Basra Ahmad Amru, Halimah Anggi Rahmani, Sheila Azelya Fernanda, Yudhi Nugraha, Muhammad Yusuf Arya Ramadhan, Nure Fahriz Perdana Harahap, Ahmad Fauzantoro, Nasihin Saud Irsyad, Meiskha Bahar, Oktania Sandra Puspita, Fajriati Zulfa, Kori Yati, Mahdi Jufri, Misri Gozan

515-525



PDF

Component Analysis of Propolis from Papua New Guinea

Ryo Miyata, Mikina Matsui, Shigenori Kumazawa

526-530



PDF

Molecular Detection of Hemagglutinin Gene Fragment of Avian Influenza Virus H9 Subtype Obtained from Poultry Commercial Farm with Prominent Symptom of Decreased Egg Production

Niken Respati Maharani, Heru Susetya, Michael Haryadi Wibowo

531-539



PDF

Genetic Background of β -Lactamase Genes in Extraintestinal Pathogenic *Escherichia coli* ST131 in Indonesia Based on Whole-Genome Sequencing (WGS) Sequences


Noor Andryan Ilsan, Sipriyadi Sipriyadi, Melda Yunita, Siti Nurfajriah, Maulin Inggraini

540-548

**Molecular Identification, GC-MS Analysis of Bioactive Compounds and Antimicrobial Activity of Thermophilic Bacteria Derived from West Sumatra Hot-Spring Indonesia**

Zona Octarya, Titania Tjandrawati Nugroho, Yuana Nurulita, Saryono

549-561

**JOURNAL INFORMATION**[About The Journal](#)[Editorial Team](#)[Focus and Scope](#)[Publication Ethics](#)[Privacy Statement](#)[Author Guidelines](#)[Register New Account](#)[Submit or Check Status](#)[Contact](#)**INDEXING**



PUBLISHER



Department of
Biology



GOOGLE SCHOLAR CITATIONS

HAYATI J.Biosci : GS citation

	All	Since :
Citations	8186	587

GS citation

VISITOR STATISTICS

00576936

[View Hayati J. Biosci. Stats](#)




HJB Most Read Papers

[Propolis Components and Biological Activities from Stingless Bees Collected on South Sulawesi, Indonesia](#)

 84


1 January 2020

 Ryo Miyata, Muhamad Sahla...

[Photosynthesis and Transpiration Rates of Rice Cultivated Under the System of Rice Intensification and the Effects on Growth and Yield](#)

 76

10 November 2016

 Nurul Hidayati, . Triadiati...

[Localization of Alkaloid and Other Secondary Metabolites in Cinchona ledgeriana Moens: Anatomical and Histochemical Studies on Fresh Tissues and Cultured Cells](#)

 75


1 January 2020

 Dian Rahma Pratiwi, Yohana...

[The Development of In Vitro Culture Sterilization Method of Gametophyte Explant Lopholejeunea sp.](#)

 60


1 April 2021

 Anna Widyastuti, Afiatry ...

[Antibacterial and Antioxidant Activities of Fungal Endophytes Isolated from Medicinal Plants in Simeulue Island, Aceh](#)

 60

28 June 2022

 Ahmad Fathoni, Muhammad Ily...



This journal is published under the terms of the <https://creativecommons.org/licenses/by-nc/4.0/>



IPB University
— Bogor Indonesia —

HAYATI Journal of Biosciences (p-ISSN: 1978-3019; e-ISSN: 2086-4094)

Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Dramaga Campus, Bogor 16680, Indonesia

Phone/Fax: +62-251-8421258; E-mail: hayati.jbiosci@apps.ipb.ac.id; URL : <http://journal.ipb.ac.id/index.php/hayati>

Platform &
workflow by
OJS / PKP

HAYATI

p-ISSN: 1978-3019
e-ISSN: 2086-4094

Journal of Biosciences

[ABOUT](#)[CURRENT](#)[ARCHIVES](#)[EDITORS](#)[SUBMISSIONS](#)[REGISTER](#)[CONTACT](#)[SEARCH](#)
[Home](#) / [Editorial Team](#)

Editorial Team

Chief Editor

[Dr. Iman Rusmana](#), Bogor Agricultural University, Indonesia

Managing Editor

[Dr. Kanthi Arum Widayati](#), Bogor Agricultural University, Indonesia

Editorial Board

[Prof. Koji Nakamura](#), Kanazawa University, Japan

[Prof. Yasumasa Bessho](#), Nara Institute of Science and Technology, Japan

[Prof. Wei Zhang](#), Flinders University, Australia

[Prof. Lisdar Idwan Sudirman](#), Bogor Agricultural University, Indonesia

[Prof. Michel Raymond](#), Institut des Sciences de l'Evolution, France

[Prof. Dr. dr. Sri Budiarti](#), Bogor Agricultural University, Indonesia

[Prof. Youji Nitta](#), Fukushima University, Japan

[Prof. Dave B. Nedwell](#), University of Essex, United Kingdom

[Prof. Aris Tri Wahyudi](#), Bogor Agricultural University, Indonesia

[Assoc. Prof. Dr. Yaya Rukayadi](#), Department of Food Science. Faculty of Food Science and Technology. Universiti Putra Malaysia

[Dr. Patrick O'Connor](#), The University of Adelaide, Australia

[Dr. Ardiansyah](#), Universitas Bakrie, Indonesia

[Dr. Rika Indri Astuti](#), Bogor Agricultural University, Indonesia

[Dr. Mashuri Waite](#), University of Hawaii, USA

[Dr. Atsushi Ido](#), Ehime University, Japan

[Dr. Sastia Prama Putri](#), Osaka University, Japan

[Dr. Bambang Retnoaji](#), Gadjah Mada University, Indonesia

[Dr. Malcolm Wegener](#), The University of Queensland, Australia

[Dr. Miftahudin](#), Bogor Agricultural University, Indonesia

[Dr. Dodi Safari](#), Eijkman Institute, Indonesia

[Dr. Emmanuel Paradis](#), Institut des Sciences de l'Evolution de Montpellier, France

[Dr. Erin Phelps Riley](#), San Diego State University, USA

[Dr. Rebecca Johnson](#), Columbia University, USA

[Dr. Takuya Imamura](#), Kyushu University, Japan

[Prof. Dr. Dyah Perwitasari](#), Bogor Agricultural University, Indonesia

[Dr. Tsukasa Sanosaka](#), Keio University, Japan

[Dr. Ocky Karna Radjasa](#), Diponegoro University, Indonesia

[Dr. Tukirin Partomihardjo](#), Indonesian Research Center for Biology (LIPI), Indonesia

[Dr. Yamato Tsuji](#), Kyoto University, Japan

[Dr. Darjono](#), Indonesian Research Center for Biology (LIPI), Indonesia

[Prof. Dr. Hamim](#), Bogor Agricultural University, Indonesia


Technical Editor

[Dr. Puji Rianti](#), Bogor Agricultural University, Indonesia

[Mafrikhul Muttaqin](#), MSi, Bogor Agricultural University, Indonesia

Online Manager

[Jepri Agung Priyanto](#), MSi, Bogor Agricultural University, Indonesia



JOURNAL INFORMATION


[About The Journal](#)

[Editorial Team](#)

[Focus and Scope](#)

[Publication Ethics](#)[Privacy Statement](#)[Author Guidelines](#)[Register New Account](#)[Submit or Check Status](#)[Contact](#)

INDEXING



PUBLISHER



GOOGLE SCHOLAR CITATIONS

HAYATI J.Biosci : GS citation		
	All	Since :
Citations	8186	587
GS citation		

VISITOR STATISTICS

00576939

View Hayati J. Biosci. Stats



HJB Most Read Papers

[Propolis Components and Biological Activities from Stingless Bees Collected on South Sulawesi, Indonesia](#)

84

1 January 2020

Ryo Miyata, Muhamad Sahla...

[Photosynthesis and Transpiration Rates of Rice Cultivated Under the System of Rice Intensification and the Effects on Growth and Yield](#)

76

10 November 2016

Nurul Hidayati, . Triadiati...

[Localization of Alkaloid and Other Secondary Metabolites in Cinchona ledgeriana Moens: Anatomical and Histochemical Studies on Fresh Tissues and Cultured Cells](#)

75

1 January 2020

Dian Rahma Pratiwi, Yohana...

[The Development of In Vitro Culture Sterilization Method of Gametophyte Explant Lopholejeunea sp.](#)

60

1 April 2021

Anna Widyastuti, Afiatry ...

[Antibacterial and Antioxidant Activities of Fungal Endophytes Isolated from Medicinal Plants in Simeulue Island, Aceh](#)

60

28 June 2022

Ahmad Fathoni, Muhammad Ily...



This journal is published under the terms of the <https://creativecommons.org/licenses/by-nc/4.0/>



IPB University
— Bogor Indonesia —

HAYATI Journal of Biosciences (p-ISSN: 1978-3019; e-ISSN: 2086-4094)

Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Dramaga Campus, Bogor 16680, Indonesia

Phone/Fax: +62-251-8421258; E-mail: hayati.jbiosci@apps.ipb.ac.id; URL : <http://journal.ipb.ac.id/index.php/hayati>

Platform &
workflow by
OJS / PKP

Nicotiana tabacum var. Virginia Bio Oil-based Pyrolysis Extraction Have Prominence Antimicrobial Potential Compared to Ethanol Heat Reflux Extraction (EHRE)

Andri Pramesyanti Pramono¹, Basra Ahmad Amru¹, Halimah Anggi Rahmani¹, Sheila Azelya Fernanda¹, Yudhi Nugraha³, Muhammad Yusuf Arya Ramadhan², Andre Fahriz Perdana Harahap², Ahmad Fauzantoro³, Nasihin Saud Irsyad¹, Meiskha Bahar¹, Oktania Sandra Puspita¹, Fajriati Zulfa¹, Kori Yati^{4,5}, Mahdi Jufri⁴, Misri Gozan^{2*}

¹Faculty of Medicine, Universitas Pembangunan Nasional Veteran Jakarta, Jakarta, Indonesia

²Chemical Engineering Department, Faculty of Engineering, Universitas Indonesia, Depok, Indonesia

³National Research and Innovation Agency Republic of Indonesia, Jakarta, Indonesia

⁴Faculty of Pharmacy and Science, Universitas Muhammadiyah Prof. DR. HAMKA, Jakarta, Indonesia

⁵Faculty of Pharmacy, Universitas Indonesia, Depok, Indonesia

ARTICLE INFO

Article history:

Received March 2, 2021

Received in revised form September 24, 2021

Accepted October 5, 2021

KEYWORDS:

bacterial growth inhibition,
Enterococcus faecalis,
Escherichia coli,
Pseudomonas aeruginosa,
Staphylococcus aureus,
tobacco leaves from Ponorogo Indonesia

ABSTRACT

Tobacco leaf contains antibacterial secondary metabolite compounds, such as phenol, alkaloids, and essential oils. This study compares the potential antibacterial effects of Indonesian tobacco leaf extracted using the heat reflux method (producing an extract) and pyrolysis method (providing a bio-oil). The tobacco leaf extract was challenged against *Staphylococcus aureus* ATCC 25923, *Enterococcus faecalis* ATCC 29212, *Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853. The bio-oil from the pyrolysis method showed inhibitory Kirby Bauer zones higher than those of the extract from heat reflux method, with the maximum results in the pyrolysis method indicating zones of 6.35 mm (*S. aureus*), 5.90 mm (*E. faecalis*), 3.97 mm (*E. coli*), and 5.025 mm (*P. aeruginosa*). Further study analyzed the effectiveness of the disc and well diffusion antibacterial test methods for measuring the antibacterial effect of bio-oils against *Escherichia coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC 27853. The bio-oil used in the well diffusion test method showed the most significant antibacterial effectiveness. It showed the biggest inhibition zone, with a maximum of 11.65 mm and 8.90 mm for *E. coli* and *P. aeruginosa*. Our results showed *Nicotiana tabacum* var. Virginia Bio Oil from Ponorogo (Indonesia) is a strong potential antimicrobial, especially using well diffusion test.

1. Introduction

The risk of nosocomial infection is highest approximately 48 hours after admission to a hospital environment and 30 days after hospital care ends (Suleyman *et al.* 2018). Such infections can be of exogenous origins, such as from epidemic diseases, person-to-person contact, and contact with devices (Suleyman *et al.* 2018). Antibiotic resistance caused by inappropriate treatments and therapies can increase the risk of nosocomial infections in patients in hospitals (Gentile *et al.* 2020). *Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, and

Pseudomonas aeruginosa are the most common bacteria causing nosocomial infections (Kudo *et al.* 2014; Iseppi *et al.* 2020; Jean *et al.* 2020; Kolar *et al.* 2020).

The use of antimicrobials from plants for nosocomial infection can reduce the side effects of excessive antibiotics and antibiotic resistance (Cheesman *et al.* 2017). The Indonesian government's limitation of cigarette production makes it important to look for alternative products, one of which is health products. In addition to social and economic needs, tobacco contains many antibacterial substances and can be used as drugs and antimicrobial health products. One of Indonesia's most common tobacco variants is *Nicotiana tabacum* var Virginia, which includes 63% of tobacco types in this country (Ahsan

* Corresponding Author

E-mail Address: mgozan@che.ui.ac.id

et al. 2014). *Nicotiana tabacum* Var Virginia originated from Virginia (America), brought to Indonesia in 1928, and planted in Bojonegoro for the first time to meet the demand for cigarette tobacco in Indonesia (Pertanian RI 2011). Besides containing high levels of nicotine, pyridine, indole, and d-limonene (Gozan *et al.* 2014; Andhika Priotomo *et al.* 2018). Tobacco can be used as an antimicrobial agent because its leaves contain secondary metabolite compounds, including alkaloids, flavonoids, phenols, saponins, and steroids (Sokunvary Oeung *et al.* 2016; Fauzantoro *et al.* 2017; Fathi *et al.* 2018), which can inhibit the growth of bacteria (Pramono *et al.* 2018).

A common extraction method in extracting antimicrobial compounds of the *Nicotiana tabacum* is Ethanolic Heat Reflux Extraction (EHRE) methods (Fauzantoro *et al.* 2017; Pramono *et al.* 2018; Banožić *et al.* 2020). Some of the studies used the pyrolysis extraction method to extract the antimicrobial compound of the *Nicotiana tabacum* (Ario Putra *et al.* 2019; Fernanda *et al.* 2021). The pyrolysis method involves heating a substance to a high temperature without the oxygen that can cause the disintegration of chemical compounds, this process results in bio-oil compounds (Gozan *et al.* 2014). The advantage of the pyrolysis extraction method is that it can attract active compounds in large amounts, with a high temperature (Andhika Priotomo *et al.* 2018), but the antimicrobial potential from bio oil product still needs to be evaluated. Here we compared the antimicrobial potential of *Nicotiana tabacum* L. var. Virginia extracts produced using the heat reflux and pyrolysis methods from Ponorogo, Indonesia. The comparison with another study is also being discussed.

2. Materials and Methods

2.1. Plant Origin and Sample Collection

Tobacco plants (*Nicotiana tabacum* L. var. Virginia) as raw material for the extracts were obtained from Ponorogo, East Java Province (Indonesia). Fresh leaves were collected and immediately processed. The tobacco leaves used are tobacco leaves that are approximately three months old and are ready to be harvested. The tobacco plant is a subtropical plant but can grow in low pliers wide climate (29–33°C). *Nicotiana tabacum* L. var. Virginia is grown in Ponorogo, Indonesia, at the end of the rainy season, and harvested in the dry season, April-Mei in Ponorogo, East Java, Indonesia (Pertanian RI 2011).

2.2. Preparation of Leaf Extracts Using the Heat-Reflux Extraction Method (Extract)

Several steps are needed to make an extract using the heat reflux method. First, the tobacco leaves are cleaned with water. Second, the tobacco leaves are dried in an oven at 120°C for 2 hours. After the tobacco leaves are dried, they are ground into a dried leaf powder. Up to 50 g of dried leaf powder is then put into an Erlenmeyer glass and dissolved with 150 ml of ethanol solvent. Then, the solution is heated to a temperature of 80°C. This heating process is a vapour, which is put into a condenser until it forms a concentrated tobacco leaf extract (briefly named extract) (Fauzantoro *et al.* 2017).

2.3. Preparation of Leaf Extracts Using the Pyrolysis Method (Bio-oil)

Another extraction method used is the pyrolysis method. First, the tobacco leaves are cleaned and then dried. After that, the tobacco leaves are chopped into smaller pieces, and 250 g of chopped leaves are converted to a powder form. This is done to improve the efficiency of the pyrolysis process. The pyrolysis reaction is carried out inside a closed pyrolysis reactor so that no external gas can enter the reactor. The tobacco leaves are heated to a temperature of 500°C. The heating process produces gas and char. The gas and char are separated using a cyclone separator. Finally, the gas is cooled until it forms a bio-oil (Gozan *et al.* 2014).

2.4. Bacterial Strains and Growth Conditions

The bacteria used in this study were obtained from the American Type Culture Collection (ATCC). They included the following strains: *Staphylococcus aureus* ATCC 25923, *Enterococcus faecalis* ATCC 29212, *Escherichia Coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853. All bacteria were cultured overnight with agitation at 150 rpm, in flasks containing Mueller-Hinton broth (Merck, Darmstadt, Germany), before being used in the experiment.

2.5. Antibacterial Activity Assessment

The samples used in this research are divided into several groups: 20%, 40%, 60%, 80%, and 100%. The percentage is obtained by diluting the extract with aquadest or bio-oil with propylene glycol. One hundred percent is the initial concentration obtained from direct extraction results. Then 80%, 60%, 40%, and 20% dilutions are obtained by adding a certain amount. The bio-oils and extracts were then analyzed using

the Kirby Bauer antibacterial test (Hudzicki 2009; Balouiri *et al.* 2016). Each experiment used both negative (only with aquadest for EHRE experiment, only with propylene glycol for the pyrolysis extraction experiment) and positive (Ciprofloxacin) control groups. The repetitive amount used for each test was determined using the Federer formula. Based on the Federer analysis, 4 samples are required for each concentration test for the *Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa* experiments. Furthermore, to assess the Kirby Bauer antibacterial analysis test's strength, we performed a well diffusion antibacterial analysis test on *Escherichia coli* and *Pseudomonas aeruginosa*.

2.5.1. Disc Diffusion Method

The Kirby-Bauer method was performed using a modification of the disc diffusion method. Bacteria were grown on a solid medium. A 0.5 McFarland standard bacterial suspension was prepared in a sterile saline solution (NaCl 0.85% (w/v); Merck, Darmstadt, Germany) via overnight culturing in a flask containing Mueller-Hinton Broth. Respective bacterial suspensions were seeded on Mueller-Hinton agar (MHA; Merck, Germany) plates, and 6 mm sterile blank discs, which had already been soaked with respective concentrations of extract, were impregnated on top. The prepared plates were incubated at 37°C for 24 hours. The inhibitory zone was identified and recorded carefully (Hudzicki 2009).

2.5.2. Well Diffusion Method

Bacteria were grown on a solid medium. 0.5 McFarland standard bacterial suspensions (*Staphylococcus aureus* ATCC 25923, *Enterococcus faecalis* ATCC 29212, *Escherichia coli* ATCC 25922, and *Pseudomonas aeruginosa* ATCC 27853) were prepared in sterile saline solution (NaCl 0.85% (w/v); Merck, Darmstadt, Germany) via overnight culturing in a flask containing Mueller-Hinton Broth. The well diffusion method involved making a 6 mm diameter hole in the MHA media containing the respective bacteria. The extract was then placed inside the hole carefully, and the prepared agars were incubated at 37°C for approximately 24 hours. The inhibitory zone was identified and recorded carefully (Balouiri *et al.* 2016).

2.6. Qualitative Phytochemistry

Phytochemical testing was carried out on extract and bio-oil. Phytochemical testing aims to determine the qualitative content of secondary metabolites contained in a sample.

2.6.1. Alkaloid Test

The extract or bio-oil was added 1 ml of 2N hydrochloric acid and 9 ml of water, heated in a water bath for 2 minutes, cooled, and filtered. Then, 3 ml of the filtrate was transferred to the watch glass, and two drops of Dragendorff reagent were added. If brown sediment occurred, the extract or bio-oil contained alkaloids. If a white or yellow clot residue is dissolved in methanol with the Mayer reagent, there is a positive for alkaloid substances.

2.6.2. Flavonoid Test

1 ml of the extract or bio-oil is evaporated, and the rest is dissolved in 1-2 ml of ethanol (95%) P. Then, 500 mg of zinc powder and 2 ml of 2N hydrochloric acid are added, the solution is allowed to stand for 1 minute, and then ten drops of concentrated hydrochloric acid are added. If the solution turned red in 2-5 minutes, it meant it contained flavonoids.

2.6.3. Tanin Test

Simplicia extract was added 50 ml of aquadest, boiled for 5 minutes, and then cooled. Then, 5 ml of the filtrate was transferred to a test tube, and FeCl₃ reagent was dropped in. If there was a greenish-black colour, that indicated the presence of tannin compounds.

2.6.4. Saponin Test

Extract or bio-oil was put into a test tube, 10 ml of hot water was added, and the mixture was cooled and then shaken vigorously for 10 seconds to form a stable white foam as high as 1-10 cm for not less than 10 minutes. If, with the addition of 1 drop of 2N hydrochloric acid, the foam did not disappear, that indicated that the extract or bio-oil contained saponins.

2.6.5. Steroid/Triterpenoid Test

The extract or bio-oil was added to 20 ml ether and macerated for 2 hours. Then, three drops of the filtrate were transferred to watch glass and dripped with a Lieberman-Burchard reagent (glacial acetic acid-concentrated sulfuric acid). If it turned red, that indicated the presence of steroid compounds, green indicated triterpenoid compounds.

2.7. Quantitative Phytochemistry

Quantitative Phytochemical testing was carried out on extract and bio-oil to measure the total alkaloids and polyphenols in the extract and bio-oil.

2.7.1. Total Flavonoid

A total of 10.0 mg of quercetin was put into a 10 ml volumetric flask and dissolved with ethanol until the limit mark to obtain a 1,000 µg/ml concentration. The standard solution of quercetin with a concentration of 1,000 µg/ml is used for the manufacture of series (3 µg/ml, 5 µg/ml, 7 µg/ml, 9 µg/ml, 11 µg/ml). Then from each concentration, 3 ml of ethanol, 0.2 ml of 10% AlCl₃, 0.2 ml of 1M of Na acetate were added and distilled water to a volume of 10 ml. Then the standard solution at which concentration has been determined is left to stand for operating time, then the absorbance is read at a wavelength of 434.5 nm on a spectrophotometer.

2.7.2. Total Polyphenol

A total of 300 µl of a gallic acid solution with a 50 µg/ml concentration added with 1.5 ml of Folin-Ciocalteu reagent (1:10), then shaken and let stand for 3 minutes. The solution was added with 1.2 ml of 7.5% Na₂CO₃ solution, shaken until homogeneous, and allowed to stand at room temperature in 0-90 minute. The absorbance was measured at a wavelength of 600-850 nm on a spectrophotometer.

2.8. Data Analysis

This experiment was performed in duplicate with a minimum of four repeats. Data were analyzed using a one-way parametric test with the statistical program SPSS 21.0 (Statistical Package for the Social Sciences). A parametric test can be done if the data variance is homogeneous and normal. An independent T-test was performed to equalize the bacterial tests with different concentrations and analyze the differences between the two extraction methods and two antibacterial activity assessments. The significance level for the difference was set at p<0.05.

3. Results

We have previously performed EHRE extraction methods (resulting extract) with ethanol solvent for *Nicotiana tabacum* Var Virginia leaves from Indonesia. This study performed the pyrolysis extraction method (resulting bio-oil) with propylene glycol solvent. The yields produced using the EHRE method are lower (23wt%) than those produced using pyrolysis extraction methods (47.6wt%). Higher nicotine yields of up to 31.1% were obtained using pyrolysis extraction compared to the EHRE method (6.3%). The EHRE method's extraction procedure was done in 6 hours

with 1 atm and 150 rpm centrifuge speed (heated at a temperature of 80°C). In the pyrolysis method, the extraction is done in 50 minutes with 1 atm and heated at 500°C.

3.1. Qualitative Phytochemistry, Quantitative Phytochemistry, and Yield Extract Efficiency

Qualitative phytochemical tests are performed to determine the active compounds in tobacco leaf extracts, which can be useful as antibacterial agents. Based on the phytochemical test results of the tobacco leaf extracts examined in this study, it is confirmed that the active antimicrobial compounds in tobacco leaf extracts are alkaloids, flavonoids, saponin, tannin, phenolic, glycoside, and triterpenoid (Table 1). The number of qualitative total alkaloids is greater in bio-oil, but total polyphenol was still higher in the tobacco extract (Table 2).

Although the polyphenol compound was found higher in the EHRE methods, on the contrary, nicotine yields showed the opposite around. The higher nicotine yields were obtained from pyrolysis extraction until 31.1% of yields (Table 3). The yields produced from

Table 1. Qualitative phytochemical test results for *Nicotiana tabacum* L. var Virginia leaf extracted using the heat reflux method (extract) and the pyrolysis method (bio-oil)

Compound	Qualitative results	
	Heat reflux extraction (extract)	Pyrolysis extraction (bio-oil)
Alkaloid	+	+
Saponin	+	+
Tanin	+	+
Phenolic	+	+
Flavonoid	+	+
Triterpenoid	+	+
Glycoside	+	+
Steroid	-	-

Table 2. Quantitative phytochemical test results for *Nicotiana tabacum* L. var Virginia leaf extracted using the heat reflux method (extract) and the pyrolysis method (bio-oil)

	Qualitative results	
	Heat reflux extraction (Extract)	Pyrolysis extraction (Bio-oil)
Total flavonoid	2,125.60 ppm	2,579.36 ppm
Total polyphenol	73,273.73 ppm	15,800.00 ppm

EHRE methods (23%) are less than those from pyrolysis extraction (47.6%).

3.2. Antimicrobial Activity

3.2.1. Antibacterial Potential of Extracts Produced using Pyrolysis and Heat Reflux Methods against *Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, and *Pseudomonas aeruginosa*

The first analysis compared the antibacterial agents' effectiveness using different extraction methods, namely heat reflux (EHRE) and pyrolysis extraction,

on several bacterial tests. The study results showed that the pyrolysis extraction method produced higher antibacterial potential than the heat reflux method (Figure 1). A significant statistically difference was only seen in *Enterococcus faecalis*, but not in other bacteria. Even so, pyrolysis is still better in inhibiting other bacteria even though it does not show significance in statistical tests.

The data also showed a significant difference in extracts' effectiveness ranging from 40 to 100% against *Enterococcus faecalis* compared to their effectiveness

Table 3. Comparison of Ethanolic Heat Reflux Extraction (EHRE) and pyrolysis extraction yield

Methods	Solvent	Operation conditions	Yields (wt%)	Nicotine yields (wt%)	Reference
Ethanolic Heat Reflux Extraction (EHRE)	Ethanol	Ethanol bp, 1 atm, 6 h, 150 rpm, heated at a temperature of 80°C	23.0	6.3	This research
Pyrolysis extraction	Propylene glycol	Propylene glycol, 1 atm, 50 minutes, heated at a temperature of 500°C	47.6	31.1	This research




Figure 1. Comparison of the antibacterial potential of extracts produced using pyrolysis and heat reflux extraction (mm). (A) *Staphylococcus aureus*, (B) *Enterococcus faecalis*, (C) *Escherichia coli*, (D) *Pseudomonas aeruginosa*. *The mean difference is significant at 0.05

against the other three test bacteria (Figure 1). This study showed that *Enterococcus faecalis* is more susceptible to active antimicrobial substances in tobacco extracts and bio-oils. *Enterococcus faecalis* belongs to the susceptible bacteria in analyzing the potential antimicrobial differences between extractions methods, especially in EHRE and pyrolysis extraction methods.

When it comes to the well diffusion test, *Nicotiana tabacum* var Virginia's showed the highest antibacterial activity in bio-oil produced using the pyrolysis extraction method compared to the EHRE method, which is tested against *Escherichia coli* (Table 3). This is because of the pyrolysis extraction method that we have done has reached 47.6wt%, which is very much higher than the EHRE method, which is only 23wt%. An increase in extraction yield could be accompanied by an increase in *Nicotiana tabacum* var Virginia's antimicrobial compound and enhancement of the inhibition diameter. This is also supported by the GC-MS from our bio-oils results which show that dominant compounds in the bio-oils from the pyrolysis process were organic acids and pyridine (besides nicotine), which the antibacterial potential are well known. In contrast, the heat reflux extract was dominated by γ -sitosterol, where the antimicrobial potential is still unclear.

When comparing the extraction methods for the *Nicotiana tabacum* in this study with other previous extraction studies from the same plant species (*Nicotiana tabacum* sp.), it seems that the maceration method produces more significant inhibition than the pyrolysis method in this study (Table 4). The factor that influences these results must be studied further.

When comparing the antibacterial test, *Nicotiana tabacum* antimicrobial test against *Escherichia coli* with well diffusion in this study, with other previous studies against other bacteria, showed that *E. coli* has a higher susceptibility to the antibacterial compound

in *Nicotiana tabacum*, compared to the different bacteria that have been examined (Table 4). It can reach up to 26.66 mm in inhibition diameter. This might be due to the initial concentration used for the experiment is quite high (3 g/ml) compared to the concentration used in the pyrolysis method (0.33 g/ml) in this study. Higher concentrations for increasing the diameter of inhibition in the pyrolysis method could be an alternative for further research. This is also seen in the antimicrobial test of other plant species against *E. coli*, where *E. coli* also show high susceptibility to active substances from other plant extracts as well.

3.2.2. Comparison of Disc Diffusion Antibacterial Test to the Well Diffusion Antibacterial Test against *Escherichia coli* and *Pseudomonas aeruginosa*

A second analysis was carried out to compare the differences in antibacterial activity for bio-oils used with different test methods, namely the disc diffusion test method and the well diffusion test method when deployed against the two test bacteria (*Escherichia coli* and *Pseudomonas aeruginosa*). The analysis results indicate that extracts implemented using the well diffusion test method create greater inhibition diameters with a significant difference (Figure 2).

Our study confirms previous studies' findings, which showed that the well diffusion method is more effective than the disc diffusion method. Is also confirm from previous studies against *Escherichia coli* using different plant extracts, except *Adenium obesum*, *Syzygium aromaticum*, and *Capparis spinosa* plant extracts produced using the maceration extraction method. In testing using *Pseudomonas aeruginosa*, it also showed a similar trend, except for *Adenium obesum* and *Syzygium aromaticum* plant extracts.

Table 4. Antibacterial activity of *Nicotiana tabacum* leaf and other plants (expressed in diameter of inhibition zone (mm))

Plant	Bacteria	Type of antibacterial test	Solvent	Concentration (w/v)	Extraction method	Inhibition diameter (mm)	Reference		
<i>N. tabacum</i> Var Virginia	<i>S. aureus</i>	DD	Ethanol	0.20 g/ml	HR	3.95±0.45	TS		
	<i>E. faecalis</i>					3.20±0.10	TS		
	<i>E. coli</i>					3.54±0.16	TS		
	<i>P. aeruginosa</i>					3.60±0.15	TS		
	<i>S. aureus</i>	DD	PG	0.33 g/ml	Pyrolysis	6.35±2.24	TS		
	<i>E. faecalis</i>					5.90±1.96	TS		
	<i>E. coli</i>					3.97±1.49	TS		
	<i>P. aeruginosa</i>					3.92±0.46	TS		
	<i>E. coli</i>	WD	PG	0.33 g/ml	Pyrolysis	11.65±1.99	TS		
	<i>P. aeruginosa</i>					8.72±0.26			
<i>N. tabacum</i> L	<i>S. aureus</i>	DD	Ethanol	0.10 g/ml	Maceration	7.13±0.25	(Ameya <i>et al.</i> 2018)		
	<i>P. aeruginosa</i>					6.16±0.47			
	<i>K. pneumonia</i>					4.59±0.387			
	<i>S. aureus</i> (CI)					5.57±0.54			
	<i>S. typhi</i> (CI)					6.16±0.47			
<i>N. tabacum</i> L	<i>B. subtilis</i>	DD	Ethanol	0.25 g/ml	Maceration	13	(Akinpelu and Obuotor 2000)		
	<i>C. pyogenes</i>					15			
	<i>P. aeruginosa</i>					11			
	<i>S. marcescent</i>					11			
	<i>S. dysenteriae</i>					15			
	<i>S. aureus</i>					14			
<i>N. tabacum</i> L	<i>B. cereus</i>	DD	EA	3.00 g/ml	Maceration	22.33	(Bakht and Shafi 2012)		
			Butanol			15.33			
	<i>S. aureus</i>		EA			15.33			
			Acetone			12.33			
			Butanol			20.00			
	<i>E. carotavora</i>		EA			22.33			
			Acetone			9.66			
			Butanol			15.66			
	<i>E. coli</i>		EA			26.66			
			Butanol			20.00			
			Water			10.66			
	<i>A.tumefaciens</i>		EA			17.00			
			Butanol			15.66			
	<i>P. aeruginosa</i>		Ethanol			10.33			
			EA			24.66			
			Butanol			18.33			
			Water			12.33			
	<i>S. typhi</i>		Ethanol			14.33			
			EA			18.33			
			Acetone			15.33			
			Butanol			16.66			
	<i>N. tabacum</i> L	<i>S. pyogenes</i>	DD	Methanol	0.2 g/ml	Maceration		13.00	(Ck <i>et al.</i> 2019)
				Water				10.00	
	<i>N. tabacum</i> L	<i>S. aureus</i>	DD	Methanol		Maceration		3.10	(Gebremedhin Romha <i>et al.</i> 2018)
<i>E. coli</i>			Methanol			3.38			
<i>P. aeruginosa</i>			Methanol			2.53			

HR: heat reflux, DD: disc diffusion, WD: well diffusion, EA: ethyl acetate, PG: polyethylene glycol, TS: this study




Figure 2. The average diameter of the Kirby-Bauer and well diffusion methods (expressed in diameter of inhibition zone (mm)) produced by (A) *Escherichia coli* (B) *Pseudomonas aeruginosa*. *The mean difference is significant at 0.05

4. Discussion

4.1. Pyrolysis Extraction Degraded Some Tobacco Active Compounds but Still Have Functional Bacterial Growth Inhibition

Qualitative total alkaloids are greater in bio-oil, but total polyphenol was still higher in the tobacco extract (Table 2). This study also shows that polyphenol compounds were less in bio-oil than tobacco extract (Table 2). This can be caused by the polyphenols themselves, which are easily degraded at high temperatures done in pyrolysis, compared to alkaloids (Hanuka Katz *et al.* 2020). Our study showed that the extraction of pyrolysis degraded some active tobacco compounds with antimicrobial potential. However, pyrolysis still showed higher antimicrobial activity than EHRE. The pyrolysis extraction can isolate more other antimicrobial components (such as nicotine) than EHRE (Table 3). However, also some components are degraded due to the use of hot temperatures. Previous research from Cardoso *et al.* 2017 on tobacco extraction using the pyrolysis method showed that the phenolic compounds found in the bio-oil were a less significant number than other compounds, such as nicotine, furfural, and acetic acid (Cardoso and Ataíde 2013).

4.2. Organic Acids and Pyridine (besides Nicotine) as Bio-oils Dominant Compounds (in Pyrolysis Extract) Showed Higher Antimicrobial Properties

Our study results showed that the pyrolysis extraction method produced higher antibacterial potential than the heat reflux method (Figure 1). This could be due to the higher number of alkaloids in pyrolysis. It may also be caused by differences in the dominance of the active substance produced

by the heat reflux method and the pyrolysis in our study. Through the pyrolysis process, Bio-oil produces products with more varied compounds in higher concentrations and contain many active compounds. Our GC-MS analysis study showed that the bio-oils dominant compounds from the pyrolysis process were organic acids and pyridine, besides nicotine. Organic acid and pyridine have antimicrobial properties (Gozan *et al.* 2014). Unlike the tobacco extract produced by heat reflux, pyridine is not the dominant substance found other than nicotine, but rather γ -sitosterol (Andjani *et al.* 2019). The pyridine ring is a compound with a spectrum of biological activity the antimicrobial potential. Pyridine has antimicrobial potential against several bacteria (Desai *et al.* 2017; Radwan *et al.* 2020). Some plants that contain predominant γ -sitosterol have antimicrobial potential (Zhang *et al.* 2011; Silveira *et al.* 2017). The possibility from our results can be concluded indirectly that pyridine has more antimicrobial properties than gamma sitosterol. But how far gamma sitosterol can be antimicrobial compared to pyridine remains proven further.

Organic acids can inhibit bacterial growth by increasing the entry of transition metals into the bacteria. A strong synergy was found between low concentrations of non-toxic metals and organic acids, with an inhibitory effect on bacterial growth (Zhitnitsky *et al.* 2017). Previous studies have also shown that organic acids have an inhibitory effect on Gram+ and Gram- bacteria (Kovanda *et al.* 2019).

In addition, the dominant antimicrobial action of pyrolysis can also be caused by the relatively higher number of alkaloids. The antibacterial mechanism of flavonoid compounds causes damage to the bacterial cell walls and disrupts cell metabolic processes (Cushnie and Lamb 2011; Khameneh *et al.* 2019).

4.3. *Enterococcus faecalis* is more susceptible to Active Antimicrobial Substances compared to other Bacteria for Both EHRE (extract) and Pyrolysis (Bio-oil) Extraction

Comparing the bacteria used in our test shows that *Enterococcus faecalis* are more susceptible to active microbial substances from *Nicotiana tabacum* var Virginia than the other three test bacteria (Figure 1) in both tobacco extracts and bio-oils. Some bacterial species are more susceptible to antimicrobial compounds, and some are innately resistant. This is due to each bacterial pathogen's unique profile characteristics and antibacterial compound mechanisms (Khameneh *et al.* 2019). This was also shown in previous studies (Chandrappa *et al.* 2015; Noushad *et al.* 2018).

4.4. Well Diffusion Test Method has a more Efficient Diffusion Process compared to the Disc Diffusion Method on Bio-oils against *Escherichia coli* and *Pseudomonas aeruginosa*

A second analysis of bio-oils on different test methods (disc diffusion test method and the well diffusion test method) against the two test bacteria (*Escherichia coli* and *Pseudomonas aeruginosa*) indicate that the well diffusion test method has greater inhibition diameters with a significant difference (Figure 2). This may be due to the more efficient diffusion process of the well method. This result is confirmed by another study on antimicrobial tests (Valgas *et al.* 2007; Mansour Saleh Saif *et al.* 2017) and suggests that the well diffusion test method is more effective than the disc diffusion test method (Figure 2). In another study on a different plant extract, the well diffusion test method produced a greater inhibitory zone diameter than the disc diffusion test method (Valgas *et al.* 2007; Mansour Saleh Saif *et al.* 2017). This may be because the tobacco leaf extract's osmolarity is higher in the well diffusion test method than in the disc diffusion test method. The well diffusion test method involves making a hole in the Muller Hinton Agar (MHA), filled with tobacco leaf extract. The tobacco leaf extract diffusion into the medium will be complete and homogeneous (Balouiri *et al.* 2016). In contrast to the well diffusion test method, in the disc diffusion test method, the tobacco leaf extract is not directly poured into the MHA; instead, the disc paper is first soaked in each component of each tobacco leaf extract before being placed on top of the MHA (Balouiri *et al.* 2016). In a well diffusion test against *Escherichia coli* from different plant extracts besides *Nicotiana tabacum* (Table 1), the disc diffusion test method produced a smaller inhibitory zone diameter when compared to the well diffusion test method because the greater

absorption of the antimicrobial compound to the disc paper in the disc diffusion method made the process of antimicrobial compound diffusion to the media less than optimal (Valgas *et al.* 2007; Mansour Saleh Saif *et al.* 2017).

Moreover, another antibacterial test against *Escherichia coli* using different plant extracts showed the same trend, except the tests that used *Adenium obesum*, *Syzygium aromaticum*, and *Capparis spinosa* plant extracts using the maceration extraction method (Mansour Saleh Saif *et al.* 2017). Thus, it is most likely that the findings in our study on the well diffusion test method are more effective than the disc diffusion test method against *Escherichia coli* are comparable to the findings of other antibacterial test experiments (Elisha *et al.* 2017; Mansour Saleh Saif *et al.* 2017; Valgas *et al.* 2007).

Disc diffusion and well diffusion antibacterial testing with *Nicotiana tabacum* have been conducted against *Pseudomonas aeruginosa*, as shown in Table 1. Our study produced results in line with previous studies, which indicated that the well diffusion test method is more effective than the disc diffusion test method. Moreover, another antibacterial test against *Pseudomonas aeruginosa* using other plant extracts showed the same trend, except for where *Adenium obesum* and *Syzygium aromaticum* plant extracts derived using the maceration extraction method were used (Mansour Saleh Saif *et al.* 2017). Thus, it is most likely that the findings of this study that the well diffusion test method is more effective than the disc diffusion test method are comparable to the findings of other antibacterial test experiments against *Pseudomonas aeruginosa*. This could be due to the different antibacterial compounds that reside in plants. Some of these antibacterial compounds might be more effectively captured in the disc diffusion and inhibit bacteria growth more effectively. In contrast, other antibacterial compounds might work better in well diffusion test methods since they are abundant in liquid form in the well (Balouiri *et al.* 2016).

In conclusions, this study's antibiotic testing comparison indicates that bio-oils produced using the pyrolysis method are better than the extracts produced using the heat reflux method for extracting the antibacterial compounds from *Nicotiana tabacum* var. Virginia from Ponorogo, Indonesia. This is proven by our result, where the pyrolysis method produces a greater diameter of inhibition compared to the EHRE method, against *Staphylococcus aureus* (6.35 mm), *Enterococcus faecalis* (5.90 mm), *Escherichia coli* (3.97 mm), and *Pseudomonas aeruginosa* (5.025 mm). This is related to antimicrobial compounds inside in the bio-oils, where alkaloids are higher in

bio-oils. However, fewer polyphenols are found in bio-oils, but there are still antimicrobial compounds predominantly found in bio-oils (pyridine) besides nicotine.

Moreover, our study showed that antibacterial using the well diffusion test method was more effective for delivering the antimicrobial compound than agar diffusion, indicated by the higher inhibition diameters produced against *Escherichia coli*. This research proves that bio-oil in the well diffusion test method showed the greatest antibacterial effectiveness. It created the biggest inhibition zone, with a maximum of 11.65 mm and 8.90 mm for *E. coli* and *P. aeruginosa* (Table 3). Nevertheless, the initial concentration used in the experiments, the test method and susceptibility of bacteria against the active antimicrobial compound were the factor that should be considered to influence the inhibition diameter enhancement on antibacterial testing.

Funding

The works were financially supported by the Universitas Indonesia PUTI Q1 2020 (Contract Nr.NKB-1415/UN2.RST/HKP.05.00/2020).

Conflicts of Interest

The authors declare no conflict of interest. The founding sponsors had no role in the study's design; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

Acknowledgments


We would like to thank Mrs. Titi Sudjiati (research assistant in our microbiology laboratory) for her support in this study.

References

- Ahsan, A., Wiyono, N., Kiting, A., 2014. Bunga Rampai-Fakta Tembakau dan Permasalahan di Indonesia, fifth ed. Tobacco Control Support Center- IAKMI, Jakarta.
- Akinpelu, D.A., Obuotor, E.M., 2000. Antibacterial activity of *Nicotiana tabacum* leaves. *Fitoterapia*. 71, 199–200. [https://doi.org/10.1016/S0367-326X\(99\)00148-3](https://doi.org/10.1016/S0367-326X(99)00148-3)
- Ameya, G., Manilal, A., Merdekios, B., 2018. *In vitro* antibacterial activity and phytochemical analysis of *Nicotiana tabacum* L. extracted in different organic solvents. *Open Microbiol. J.* 11, 352–359. <https://doi.org/10.2174/1874285801711010352>
- Andhika Priotomo, R., Fauzantoro, A., Gozan, M., 2018. Fractionation and characterization of semi-polar and polar compounds from leaf extract *Nicotiana tabacum* L. reflux ethanol extraction results. *In: AIP Conference Proceedings*. Bali: American Institute of Physics Inc, pp. 1-9. <https://doi.org/10.1063/1.5023969>.
- Andjani, H.N., Sentosa, Y., Yati, K., Jufri, M., Fauzantoro, A., Gozan, M., 2019. Determination of LC₅₀ value of *Nicotiana tabacum* L. extracts against *Gryllus bimaculatus* imago and *Galleria mellonella* larvae. *In: AIP Conference Proceedings*. Padang: American Institute of Physics Inc, pp. 1-7. <https://doi.org/10.1063/1.5139361>
- Ario Putra, D.A., Pramono, A., Anisah, A., Fauzantoro, A., Gozan, M., 2019. The effect of tobacco leaves pyrolysis extract (*Nicotiana tabacum* L. var. Virginia) against the formation of biofilm by *Staphylococcus aureus*: an *in vitro* study. *IOP Conf. Ser. Mater. Sci. Eng.* 508, 012147. <https://doi.org/10.1088/1757-899X/508/1/012147>
- Bakht, J., Shafi, M., 2012. Antimicrobial activity of *Nicotiana tabacum* using different solvent extracts. *Pak. J. Bot.* 44, 459–643.
- Balouiri, M., Sadiki, M., Ibnsouda, S.K., 2016. Methods for *in vitro* evaluating antimicrobial activity: a review. *J. Pharm. Anal.* 6, 71–79. <https://doi.org/10.1016/j.jpha.2015.11.005>
- Banožić, M., Babić, J., Jokić, S., 2020. Recent advances in extraction of bioactive compounds from tobacco industrial waste—a review. *Ind. Crops Prod.* 144, 112009. <https://doi.org/10.1016/j.indcrop.2019.112009>
- Cardoso, C.R., Ataíde, C.H., 2013. Analytical pyrolysis of tobacco residue: effect of temperature and inorganic additives. *J. Anal. Appl. Pyrolysis*, 99, 49–57. <https://doi.org/10.1016/j.jaap.2012.10.029>
- Chandrappa, P.M., Dupper, A., Tripathi, P., Arroju, R., Sharma, P., Sulochana, K., 2015. Antimicrobial activity of herbal medicines (tulsi extract, neem extract) and chlorhexidine against *Enterococcus faecalis* in Endodontics: an *in vitro* study. *J. Int. Soc. Prev. Community Dent.* 5, 89–92.
- Cheesman, M.J., Ilanko, A., Blonk, B., Cock, I.E., 2017. Developing new antimicrobial therapies: are synergistic combinations of plant extracts/compounds with conventional antibiotics the solution? *Pharmacogn. Rev.* 11, 57–72.
- Ck, A., Mi, N., Cc, O., Io, N., Fc, I., 2019. Antimicrobial activities of extracts of tobacco leaf (*Nicotiana tabacum*) and Its Grounded Snuff (Utaba) on *Candida albicans* and *Streptococcus pyogenes*. *J. Trop. Dis.* 7, 1000300. <https://doi.org/10.4172/2329-891X.1000300>
- Cushnie, T.P.T., Lamb, A.J., 2011. Recent advances in understanding the antibacterial properties of flavonoids. *Int. J. Antimicrob. Agents.* 38, 77–107. <https://doi.org/10.1016/j.ijantimicag.2011.02.014>
- Desai, N.C., Patel, B.Y., Dave, B.P., 2017. Synthesis and antimicrobial activity of novel quinoline derivatives bearing pyrazoline and pyridine analogues. *Med. Chem. Res.* 26, 109–119. <https://doi.org/10.1007/s00044-016-1732-6>
- Elisha, I.L., Botha, F.S., McGaw, L.J., Eloff, J.N., 2017. The antibacterial activity of extracts of nine plant species with good activity against *Escherichia coli* against five other bacteria and cytotoxicity of extracts. *BMC Complement. Altern. Med.* 17, 133. <https://doi.org/10.1186/s12906-017-1645-z>
- Fathi, R.M., Fauzantoro, A., Rahman, S.F., Gozan, M., 2018. Column chromatography isolation of nicotine from tobacco leaf extract (*Nicotiana tabacum* L.). *In: AIP Conference Proceedings*. Bali: American Institute of Physics Inc., pp. 1–9. <https://doi.org/10.1063/1.5023958>
- Fauzantoro, A., Muharam, Y., Gozan, M., 2017. Improvement of nicotine yield by ethanolic heat reflux extraction of *Nicotiana tabacum* var. Virginia origin of Ponorogo. *Int. J. Appl. Eng. Res.* 12, 13891–13897.

- Fernanda, S.A., Amru, B.A., Rahmani, H.A., Gozan, M., Irsyad, N.S., Bahar, M., Puspita, O.S., Zulfa, F., Pramono, A., 2021. Antibacterial Potential of *Nicotiana tabacum* L. var Virginia Pyrolysis Extract Against *Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, and *Pseudomonas aeruginosa*. *IOP Conf. Ser.: Earth Environ. Sci.* 755 012013. <https://doi.org/10.1088/1755-1315/755/1/012013>
- Gebremedhin Romha, Birhanu Admasu, Tsegaye Hiwot Gebrekida, Hailelule Aleme, Gebreyohans Gebru, 2018. Antibacterial activities of five medicinal plants in Ethiopia against some human and animal pathogens. Evidence-Based complement. *Altern. Med.* 2018, 1–10. <https://doi.org/10.1155/2018/2950758>
- Gentile, B., Grottola, A., Orlando, G., Serpini, G.F., Venturelli, C., Meschiari, M., Anselmo, A., Fillo, S., Fortunato, A., Lista, F., Pecorari, M., Mussini, C., 2020. A retrospective whole-genome sequencing analysis of carbapenem and colistin-resistant *Klebsiella pneumoniae* nosocomial strains isolated during an MDR surveillance program. *Antibiotics*. 9, 246. <https://doi.org/10.3390/antibiotics9050246>
- Gozan, M., Yasman, Wulan, P.P.D.K., Dawitri, E., 2014. Tobacco leaves pyrolysis for repellent active compound production. *Int. J. Appl. Eng. Res.* 9, 9739–9750.
- Hanuka Katz, I., Eran Nagar, E., Okun, Z., Shpigelman, A., 2020. The link between polyphenol structure, antioxidant capacity and shelf-life stability in the presence of fructose and ascorbic acid. *Molecules*. 25, 225. <https://doi.org/10.3390/molecules25010225>
- Hudzicki, J., 2009. Kirby-Bauer Disk Diffusion Susceptibility Test Protocol. American Society for Microbiology. Available at: <https://asm.org/Protocols/Kirby-Bauer-Disk-Diffusion-Susceptibility-Test-Pro>. [Date accessed: 5 September 2021]
- Iseppi, R., Di Cerbo, A., Aloisi, P., Manelli, M., Pellesi, V., Provenzano, C., Camellini, S., Messi, P., Sabia, C., 2020. *In vitro* activity of essential oils against planktonic and biofilm cells of extended-spectrum β -lactamase (ESBL)/carbapenemase-producing gram-negative bacteria involved in human nosocomial infections. *Antibiotics*. 9, 272. <https://doi.org/10.3390/antibiotics9050272>
- Jean, S.S., Chang, Y.C., Lin, W.C., Lee, W.S., Hsueh, P.R., Hsu, C.W., 2020. Epidemiology, treatment, and prevention of nosocomial bacterial pneumonia. *J. Clin. Med.* 9, 275. <https://doi.org/10.3390/jcm9010275>
- Khameneh, B., Iranshahy, M., Soheili, V., Fazly Bazzaz, B.S., 2019. Review on plant antimicrobials: a mechanistic viewpoint. *Antimicrob. Resist. Infect. Control.* 8, 118. <https://doi.org/10.1186/s13756-019-0559-6>
- Kolar, M., Cermak, P., Hobzova, L., Bogdanova, K., Neradova, K., Mlynarcik, P., Bostik, P., 2020. Antibiotic resistance in nosocomial bacteria isolated from infected wounds of hospitalized patients in the Czech Republic. *Antibiotics*. 9, 342. <https://doi.org/10.3390/antibiotics9060342>
- Kovanda, L., Zhang, W., Wei, X., Luo, J., Wu, X., Atwill, E.R., Vaessen, S., Li, X., Liu, Y., 2019. *In vitro* antimicrobial activities of organic acids and their derivatives on several species of gram-negative and gram-positive bacteria. *Molecules*. 24, 3770. <https://doi.org/10.3390/molecules24203770>
- Kudo, M., Nomura, T., Yomoda, S., Tanimoto, K., Tomita, H., 2014. Nosocomial infection caused by vancomycin-susceptible multidrug-resistant *Enterococcus faecalis* over a long period in a university hospital in Japan. *Microbiol. Immunol.* 58, 607–614. <https://doi.org/10.1111/1348-0421.12190>
- Mansour Saleh Saif, M., Ali Al-Fakih, A., Abdu Hassan, M.M., Mohammed Mansour Saleh Saif, C., 2017. Antibacterial activity of selected plant (aqueous and methanolic) extracts against some pathogenic bacteria. *J. Pharmacogn. Phytochem.* 6, 1929–1935.
- Noushad, M.C., Balan, B., Basheer, S., Usman, S. Bin, Askar, M.K.M., 2018. Antimicrobial efficacy of different natural extracts against persistent root canal pathogens: an *in vitro* study. *Contemp. Clin. Dent.* 9, 177–181. <https://doi.org/10.4103/ccd.ccd.754.17>
- Pertanian RI, K., 2011. Tembakau Virginia. In *Balitbangtan Kementerian Pertanian. Balai Penelitian Tanaman Tembakau dan Serat*, Malang.
- Pramono, A., Fauzantoro, A., Hidayati, I.R., Hygea, A., Puspita, O.S., Muktamiroh, H., Simanjuntak, K., Gozan, M., 2018. *In vitro* assay of ethanolic heat reflux extract of *Nicotiana tabacum* L. var Virginia against nosocomial bacteria pathogen *in vitro* assay of ethanolic heat reflux extract of *Nicotiana tabacum* L. var Virginia against nosocomial bacterial pathogen. *J. Phys. Conf. Ser.* 970, 012021. <https://doi.org/10.1088/1742-6596/970/1/012021>
- Radwan, M.A.A., Alshubramy, M.A., Abdel-Motaal, M., Hemdan, B.A., El-Kady, D.S., 2020. Synthesis, molecular docking and antimicrobial activity of new fused pyrimidine and pyridine derivatives. *Bioorg. Chem.* 96, 103516. <https://doi.org/10.1016/j.bioorg.2019.103516>
- Silveira, D., Maia, V., Camargo Aranha, B., Chaves, F.C., Padilha Da Silva, W., 2017. Antibacterial activity of *Butia odorata* Barb. Rodr. Extracts antibacterial foodborne pathogens synthetic preservatives chemical characterization phytosterols. *Trends Phytochem. Res.* 1, 169–174.
- Sokunvary Oeung, Nov, V., Ung, H., Roum, K., Yin, V., Keo, S., Sin Chea, 2016. Phytochemical analysis of different extracts of leaves of *Nicotiana tabacum* L. of Cambodia. *Asian J. Pharmacogn.* 1, 18–26.
- Suleyman, G., Alangaden, G., Bardossy, A.C., 2018. The role of environmental contamination in the transmission of nosocomial pathogens and healthcare-associated infections. *Curr. Infect. Dis. Rep.* 20, 1–11. <https://doi.org/10.1007/s11908-018-0620-2>
- Valgas, C., De Souza, S.M., Smânia, E.F.A., Smânia, A., 2007. Screening methods to determine antibacterial activity of natural products. *Brazilian J. Microbiol.* 38, 369–380. <https://doi.org/10.1590/S1517-83822007000200034>
- Zhang, W., Abdel-Rahman, F.H., Saleh, M.A., 2011. Natural resistance of rose petals to microbial attack. *J. Environ. Sci. Health, Part B.* 46, 381–393. <https://doi.org/10.1080/03601234.2011.572502>
- Zhitnitsky, D., Rose, J., Lewinson, O., 2017. The highly synergistic, broad-spectrum, antibacterial activity of organic acids and transition metals. *Sci. Rep.* 7, 1–13. <https://doi.org/10.1038/srep44554>

also developed by scimago:

 SCIMAGO INSTITUTIONS RANKINGS

SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

Home

Journal Rankings

Country Rankings

Viz Tools

Help



About Us


Nursing Research Manuscript

Oper

Hindawi

HAYATI Journal of Biosciences

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
<div>Indonesia</div> <div><div> Universities and research institutions in Indonesia</div><div> Media Ranking in Indonesia</div></div>	<div>Agricultural and Biological Sciences</div> <div><div>Agricultural and Biological Sciences (miscellaneous)</div><div>Biochemistry, Genetics and Molecular Biology</div><div><div>Biochemistry, Genetics and Molecular Biology (miscellaneous)</div></div></div>		<div>22</div>
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	19783019, 20864094	2005-2021	<div><div>Homepage</div><div>How to publish in this journal</div><div>bjuliandi@ipb.ac.id</div></div>

 **Sigma-Aldrich.**


Unleash the Impossible



SCOPE

HAYATI Journal of Biosciences (HAYATI J Biosci) is an international peer-reviewed and open access journal that publishes significant and important research from all area of biosciences fields such as biodiversity, biosystematics, ecology, physiology, behavior, genetics and biotechnology. All life forms, ranging from microbes, fungi, plants, animals, and human, including virus, are covered by HAYATI J Biosci. HAYATI J Biosci published by Department of Biology, Bogor Agricultural University, Indonesia and the Indonesian Society for Biology. We accept submission from all over the world. Our Editorial Board members are prominent and active international researchers in biosciences fields who ensure efficient, fair, and constructive peer-review process. All accepted articles will be published on payment of an article-processing charge, and will be freely available to all readers with worldwide visibility and coverage.

Join the conversation about this journal




Quartiles


FIND SIMILAR JOURNALS ?


1 Research Journal of Biotechnology IND	2 Malaysian Journal of Microbiology MYS	3 3 Biotech CHE	4 Asian Journal of Microbiology, Bi IND
35% similarity	35% similarity	35% similarity	34% similarity





Sigma-Aldrich.

We'll help you construct your breakthroughs










SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Sigma-Aldrich.

Building 3D bioprinted tissue models?



MERCK

Metrics based on Scopus® data as of April 2022

E

Eddy Mangopo Angi 3 weeks ago

**Melanie Ortiz** 2 weeks ago

SCImago Team

Dear Eddy,
Thank you for contacting us. Could you please expand a little bit on your comment?
Best Regards, SCImago Team

**lina rizki** 1 year ago

please i want to read for my study

← reply

**Melanie Ortiz** 1 year ago

SCImago Team

Dear Lina,
Thank you for contacting us.
We suggest you visit the journal's homepage.
Best Regards, SCImago Team

**M. Ilham** 2 years ago

Dear Scimago Team

Our journal, HAYATI J Biosci, is published by Bogor Agricultural University (Institut Pertanian Bogor) Indonesia. Please kindly put this information on the "Publisher" Tab. For your reference, please refer to:

<https://www.scopus.com/sourceid/21100824972>

<https://journal.ipb.ac.id/index.php/hayati/about>

Thank you for your kind help.

← reply

**Ling** 2 years ago

May I know if this journal still active?
Submitted manuscript more than 6 months ago and also emailed to the address provided,
but never get a new or reply.

Dear M. Ilham,
Thank you for contacting us.
As you probably already know, SJR is a portal with scientometric indicators of journals indexed in Scopus. All the metadata have been provided by Scopus /Elsevier and SCImago doesn't have the authority over these data which are property of Scopus/Elsevier. SCImago is not allowed to do any changes of the metadata unless Scopus authorizes it expressly.
Therefore, we've contacted Scopus regarding this matter and, after receiving its confirmation, the publisher has been updated.
Best Regards, SCImago Team

L

Lianah 3 years ago

Terimakasih sudah bisa bergabung. Saya sdh mengajukan dan sudah diterima submitnya tetapi sampai sekarang belum ada jawaban lewat email dan statusnya masih dalam peninjauan. Sampai kapan y. tKS SALAM

 reply**Melanie Ortiz** 3 years ago

SCImago Team

Dear Lianah,
thank you for contacting us.
Sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.
Unfortunately, we cannot help you with your request, we suggest you to contact the journal's editorial staff again, so they could inform you more deeply.
Best Regards, SCImago Team

C

Chandra 3 years ago

Q2, awesome

 reply

H

hasim 4 years ago

good

 reply

 reply**Ahmad Herison** 5 years ago

Dear Journal Management

please inform about submission in this journal and also the guide for author.

Thank you

 reply**Elena Corera** 5 years ago

SCImago Team

Dear Ahmad, in the link below you will find the information corresponding to the author's instructions of this journal. Best regards, SCImago Team

<https://www.elsevier.com/authors/journal-authors/submit-your-paper>

Leave a comment

Name

Email

(will not be published)



I'm not a robot

reCAPTCHA

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.



Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1,1,106)

[Edit Cookie Consent](#)





Sources

Title Enter title

Find sources

Title: HAYATI Journal Of Biosciences x

Improved Citescore

We have updated the CiteScore methodology to ensure a more robust, stable and comprehensive metric which provides an indication of research impact, earlier. The updated methodology will be applied to the calculation of CiteScore, as well as retroactively for all previous CiteScore years (ie. 2018, 2017, 2016...). The previous CiteScore values have been removed and are no longer available.

[View CiteScore methodology.](#)



Filter refine list

Apply Clear filters

Display options

☐ Display only Open Access journals

Counts for 4-year timeframe

☐ No minimum selected

☐ Minimum citations

☐ Minimum documents

Citescore highest quartile

☐ Show only titles in top 10 percent

☐ 1st quartile

☐ 2nd quartile

☐ 3rd quartile

☐ 4th quartile

Source type

☐ Journals

☐ Book Series

☐ Conference Proceedings

☐ Trade Publications

Apply Clear filters

1 result

[Download Scopus Source List](#) [Learn more about Scopus Source List](#)

☐ All

[Export to Excel](#)

[Save to source list](#)

View metrics for year: 2021

Source title	CiteScore	Highest percentile	Citations 2018-21	Documents 2018-21	% Cited
1 HAYATI Journal of Biosciences <i>Open Access</i>	1.0	34% 138/211 General Agricultural and Biological Sciences	126	130	42

[Top of page](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.

