



UNIVERSITAS MUHAMMADIYAH PROF. DR. HAMKA FAKULTAS FARMASI DAN SAINS

Islamic Center, Jl. Delima II/IV Klender, Jakarta Timur 13460 Telp. (021) 8611070, Fax. (021) 86603233

www.uhamka.ac.id, www.ffi.uhamka.ac.id, Email: ffi@uhamka.ac.id

SURAT TUGAS

NOMOR: 927 /F.03.01/2022

Pimpinan Fakultas Farmasi dan Sains, Universitas Muhammadiyah Prof. DR. Hamka dengan ini memberi tugas kepada :

- Nama : 1. Ni Putu Ermi Hikmawati, M.Farm.
2. apt. Tuti Wiyati, M.Sc.
- Jabatan : Dosen FFS UHAMKA
- Alamat : Islamic Center Jl. Delima Raya II/ IV, Perumnas Klender – Jakarta Timur
- Tugas : Sebagai Penulis pada Jurnal Pharmaceutical Sciences Asia:
“Protective effect and potential natural antioxidant of *Cayratia trifolia* (L.) Domin. Leaves extracts on nitrobenzene-induced hepatotoxic rats”
- Waktu : Semester GASAL TA. 2022/2023
- Lain-lain : Setelah melaksanakan tugas agar memberikan laporan kepada Dekan atau sama yang memberi tugas.

Demikian surat tugas ini diberikan untuk dilaksanakan dengan sebaik-baiknya sebagai amanah dan ibadah kepada Allah Subhanahu Wata`ala

Jakarta, 05 Oktober 2022

Dekan



Dr. apt. Hadi Sunaryo, M.Si.

Research Article

Protective effect and potential natural antioxidant of *Cayratia trifolia* (L.) Domin. leaves extracts on nitrobenzene-induced hepatotoxic rats

Ni Putu Ermi Hikmawanti^{1*}, Tuti Wiyati², M. Abdul Muis³, Farah Aisyah Nurfaizah³, Windy Septiani³, Ryantama Indrapraja Putra³

¹ Department of Pharmaceutical Biology, Faculty of Pharmacy and Sciences, Universitas Muhammadiyah Prof. DR. HAMKA, DKI Jakarta, East Jakarta, Indonesia

² Department of Pharmacology, Faculty of Pharmacy and Sciences, Universitas Muhammadiyah Prof. DR. HAMKA, DKI Jakarta, East Jakarta, Indonesia

³ Faculty of Pharmacy and Science, Universitas Muhammadiyah Prof. DR. HAMKA, DKI Jakarta, East Jakarta, Indonesia

ABSTRACT

Cayratia trifolia leaves contain polyphenols as a natural antioxidant that can dissolve in polar solvents. This study aims to evaluate the hepatoprotective and antioxidant activity of polar extracts (ethyl acetate and 70%-ethanol) were extracted sequentially from *C. trifolia* leaves on male rats induced by nitrobenzene. The extracts of *C. trifolia* leaves were given at a dose of 200 mg.Kg⁻¹ BW. Silymarin as a reference was given at a dose of 28.78 mg.Kg⁻¹ BW. The extracts and Silymarin were given orally once a day for 14 days. Rats that have received ethanol extract of *C. trifolia* leaves showed a significant decrease in Alkaline Phosphatase (ALP), Aspartate Transaminase (AST), Alanine Transaminase (ALT), total bilirubin, and direct bilirubin. A significant decrease in malondialdehyde (MDA) levels also occurred in the rats. On histopathological examination, the extract also showed a protective effect on rat liver. Thus, it can be concluded that the sequentially extracted ethanolic extract of *C. trifolia* leaves has the potential as a source of natural hepatoprotective and antioxidant activity.

Keywords:

Antioxidant, *Cayratia trifolia* L., Hepatoprotective, Nitrobenzene

1. INTRODUCTION

The liver is one of the vital organs in the body. The liver functions to help metabolism, detoxification, storage, and synthesis of certain substances such as amino acids, cholesterol, etc.¹. The agent that can protect the liver from a hepatotoxic-induced radical substance is called hepatoprotective². Herbal medicine is still used mainly by the world population for primary health care³. Herbal-based therapy for hepatoprotection is related to its phytoconstituent as an antioxidant⁴. One of those potential herbals is *Cayratia trifolia* (L.) Domin. (Vitaceae).

In Indonesia, *C. trifolia* is known as galing or lakum. This plant is called Fox grape⁵. Empirically, the leaves are used as medicine to cure headaches¹, boils, speed drying of the wound⁶, muscle pains, and antidandruff⁷.

C. trifolia leaves have been scientifically researched and proven to have several activities as antitumor⁸, anti-implantation⁹, anti-ulcer¹⁰, larvicidal¹¹, antibacterial¹², antioxidant¹³⁻¹⁴, antimutagenic¹⁵, and anti-hyperuricemia¹⁶. The purified ethanol extract of *C. trifolia* leaves at a dose <5,000 and 50,000 mg.Kg⁻¹ BW shows no liver cell damage¹⁷.

Epifriedelanol is a compound found in ethanol extract of *C. trifolia*^{8,18}. *C. trifolia* leaves contain stilbenes, several flavonoids¹⁹, and linoleic acid²⁰. The total flavonoids content in the ethanolic extract of *C. trifolia* leaves was 27.95±0.62 mgQE/g²¹. *C. trifolia* leaves also contain phytochemical compounds, such as alkaloids, polyphenols, terpenoids, steroids, and saponins. These compounds play a role in their biological activity, such as anticancer, anti-inflammatory, hepatoprotective, and antioxidant²².

*Corresponding author:

*Ni Putu Ermi Hikmawanti Email: ermy0907@uhamka.ac.id



Pharmaceutical Sciences Asia © 2022 by

Faculty of Pharmacy, Mahidol University, Thailand is licensed under CC BY-NC-ND 4.0. To view a copy of this license, visit <https://www.creativecommons.org/licenses/by-nc-nd/4.0/>

The compounds can generally be dissolved using ethanol and ethanol-water²³. Several other organic solvents (benzene, chloroform, dichloromethane, etc.) need to be considered as extraction solvents because of their possible toxic effects as residual solvents in the extracts²⁴.

Nitrobenzene (NB) is a chemical widely used in industry with an almond-like odor. It is a toxic substance and can be evaporated in the air. If exposed to high concentrations, it can cause liver damage. NB has methemoglobinemia, carcinogenic, neurotoxic, and hepatotoxic properties²⁵. It produces several free radicals such as superoxide anion during its reductive metabolism²⁶. NB-induced hepatotoxicity can be used as an *in vivo* model for hepatoprotective studies.

Previously, ethanol extract of *C. trifolia* leaves at a dose of 500 mg.Kg⁻¹ BW has hepatoprotective activity on paracetamol-induced rats²⁷. Guru Kumar et al. (2011) reported that after administration of the *C. trifolia* whole plant extract (200 mg.Kg⁻¹) orally, there was hepatocyte regeneration in rats induced by nitrobenzene²⁸. The present study evaluates the hepatoprotective and antioxidant effect of sequentially extracted polar extracts (ethyl acetate and ethanolic extracts) of *C. trifolia* leaves on nitrobenzene-induced rats. Thus, an extract containing active compounds with antioxidant and liver protection effects will be obtained.

2. MATERIALS AND METHODS

2.1. Plant Material

The *C. trifolia* leaves were collected from the Tuban, East Java, Indonesia. The plant was identified in Conservation Research Center and Botanical Garden, Indonesian Institute of Sciences/Lembaga Ilmu Pengetahuan Indonesia (LIPI), Bogor, Indonesia. The *C. trifolia* fresh leaves were washed under running tap water. Then, the leaves were dried for a few days in indirect sunlight and made into powder.

2.2. Extraction

The dried leaves powder (1.5 Kg) was extracted in various solvents such as *n*-hexane, ethyl acetate, and 70%-ethanol according to their increasing polarity by cold maceration (2 times, 24 hours). The extracts were filtered through filter paper. The residue was further repeated twice with the same solvent. Each filtrate was evaporated using a vacuum rotary-evaporator N-1200 BS series (EYELA, Shanghai, China) at 40°C. The percentage yield of *n*-hexane, ethyl acetate, and ethanolic extracts of *C. trifolia* leaves was calculated²⁹.

2.3. Phytochemical Identification

The presence of secondary metabolites (viz.,

alkaloids, flavonoids, phenolic, etc.) in the extracts were qualitatively identified using the standard procedures described in the²⁹⁻³¹. Tests were carried out using reagents such as Dragendorff, Mayer, and Bouchardat (for alkaloids detection); Shinoda (for flavonoids detection); FeCl₃ 3% (for phenolics detection), 10% gelatine (for tannins detection), etc.

2.4. Total Phenolic Content Estimation

Total phenolic content (TPC) was measured by colorimetry according to the Folin-Ciocalteu method using gallic acid as the standard³². Extract (300 µL) were mixed vigorously with 1.5 mL of the Folin-Ciocalteu reagent. The reagent was diluted ten times in water. After 3 min, 1.2 mL of sodium carbonate solution (7.5%) was added to the mixture. The mixed solution was then incubated at room temperature (25°C) for 30 min. The absorbance was measured at 743.50 nm with a spectrophotometer UV-Vis Seri UV-1601 (Shimadzu, Kyoto, Japan). Gallic acid solution with concentrations ranging from 14-50 mg.L⁻¹ was used for calibration. A dose-response linear regression was generated by using the gallic acid standard absorbance. A linear calibration curve of gallic acid was obtained ($y=0.0129x+0.0545$) with the coefficient of determination (R) value of 0.99. The total phenolic content was expressed as gallic acid equivalent (mg of GAE.g⁻¹ of extract). The estimation was performed in triplicate, and the results were expressed as mean±SD.

2.5. Preparation of rats

Male white rats (*Rattus norvegicus* L.) of Wistar strain of 2-3 months old (bodyweight 140-160 g) were used in this study. The rats were maintained in a well-ventilated room with a natural day-night period. Rats were fed with standard pellets for rodents and water ad libitum throughout the experimental period. Before treatment, the rats were quarantined for seven days.

2.6. Experimental Design

This protocol (no.02/19.10/0230) was approved by the Health Research Ethics Commission of Universitas Muhammadiyah Prof. DR. HAMKA. Determination of the dose of the extracts in this study refers to Kumar et al. (2011), where used a methanol extract of the aerial part of *C. trifolia* at a dose of 200 mg/kg BW rats (28). Whereas the dose of Silymarin refers to its use in humans (280 mg.60 Kg⁻¹) which has been converted at a rat of 28.78 mg.Kg⁻¹ once a day. The extracts and Silymarin were prepared in 0.5% Na-CMC.

Rats were divided into five groups. All groups have received the treatments for 14 days. All groups were induced with nitrobenzene (50 mg.Kg⁻¹BW) orally on

day 1, except the normal control group.

- Normal control group: rats have received standard pelleted rat feed without Na-CMC each day orally.
- Untreated control group: rats have received Na-CMC 0.5% each day orally.
- Silymarin control group: rats have received Silymarin at 28.78 mg.Kg⁻¹ each day orally.
- Ethyl acetate extract group: rats have received ethyl acetate extract at 200 mg.Kg⁻¹BW each day orally.
- Ethanolic extract group: rats have received ethanolic extract at 200 mg.Kg⁻¹BW each day orally.

2.7. Preparation of the Blood Serum

On day 15th, after administration of the extracts, the rats were anesthetized using ketamine intramuscularly. The blood sample was taken through the orbital sinus and then collected in a microtube. Blood was then centrifuged for 15 minutes at 6,000 rpm to obtain the blood serum for biochemical analysis. After blood sampling, the rats were sacrificed by cervical dislocation, and the liver was dissected entirely and put into the 10% buffered formalin for histopathological evaluation.

2.8. Determination of Liver Enzymes Levels

The Alkaline Phosphatase (ALP) level in rat blood serum was determined using a diagnostic reagent (DiaSys, Holzheim, Germany) with a kinetic photometric test method according to the Germany Society of Clinical Chemistry procedure³³. The absorbance was read at 405 nm. The Aspartate Aminotransferase (AST) level in rat blood serum was carried out following the protocol of GOT (ASAT) IFCC mod. liquiUV multipurpose reagent kit (HUMAN, Magdeburg, Germany), whereas the Alanine Aminotransferase (ALT) level in rat blood serum was carried out following the protocol of GPT (ASAT) IFCC mod. liquiUV multipurpose reagent kit (HUMAN, Magdeburg, Germany). The direct (conjugated) bilirubin and total bilirubin in rat blood serum were determined using Bilirubin D+T liquicolor photometric test kit (HUMAN, Magdeburg, Germany) modified from Jendrassik & Gróf, (1938) method³⁴. The absorbance was read at 546 nm with an optical path is 1 cm. The measurement of absorbance from all liver enzyme levels was performed using a clinical spectrophotometer (Microlab-300).

2.9. Determination of Lipid Peroxidation Levels

The procedure was performed following the method reported by Esterbauer & Cheeseman (1990)³⁵. The rat liver was added with phosphate buffer pH 7.4 (5 times liver weight) and homogenized in mortar. The homogenate of the liver (1 ml) was reacted with 0.5 ml of 20% trichloroacetic acid (TCA) solution (Sigma Aldrich,

Darmstadt, Germany) and then centrifuged at 3,000 rpm for 10 minutes. The supernatant was added with 1 ml of 0.67% (w/v) Thiobarbituric acid (TBA) solution (Sigma Aldrich, Darmstadt, Germany), and then homogenized. The mixture was heated at 100°C for 10 minutes. The product of the reaction from two molecules of TBA and one molecule of MDA produces the pink-coloured solution. The absorbance can be measured at 532 nm using a spectrophotometer UV-Vis. The standard solution was used to make the standard curve is tetra ethoxy-propane (TEP) (Sigma Aldrich, Darmstadt, Germany). The blank solution contains the mixture of 1 ml of aqua distillate, 0.5 ml of 20% TCA solution, and 1 ml of 0.67% TBA solution.

2.10. Histopathological Studies

A small piece of the rat liver tissue from each group was prepared using the paraffin blocks method and transferred to a slide, and were stained using Hematoxylin and Eosin stain for histopathological observations using a biological microscope (Olympus CX23, Tokyo, Japan) at magnification 10×40. The liver cells of each rat were observed. The number of pyknotic nuclei was counted. Observations were made with five fields of view randomly for each sample.

2.11. Statistical Analysis

Data were analysed using one-way analysis of variant (ANOVA) and continued with the Tukey test ($\alpha=0.05$). The histopathological evaluation of the rat liver was analyzed descriptively.

3. RESULTS AND DISCUSSION

3.1. Characteristics of Extracts

In this study, the *C. trifolia* leaves were sequentially extracted using different solvent polarities (*n*-hexane, ethyl acetate, and 70%-ethanol). *C. trifolia* leaves are dominated by polar compounds found in the ethanolic extract (21.30 %, w/w) compared to ethyl acetate extract (1.73 %, w/w). Among the extracts, the *n*-hexane extract gave the lowest yield percentage (0.60%, w/w). The low percentage of yield indicates the most minor content of nonpolar compounds in *C. trifolia* leaves extracted in *n*-hexane.

Based on phytochemical screening, ethyl acetate extract of *C. trifolia* leaves contains alkaloids, flavonoids, phenolics, and steroids. Phytochemical compounds in ethanolic extract of *C. trifolia* leaves are alkaloids, flavonoids, phenolics, tannins, and steroids. *n*-Hexane extract contains steroids and triterpenoids. Sowmya *et al.* (2015) reported that phytochemical compounds in sequentially extracted ethyl acetate extract of *C. trifolia*

leaves are amino acids and proteins, carbohydrates, and polyphenols²². Sequentially extracted ethanolic extract of *C. trifolia* leaves contains amino acids and proteins, carbohydrates, polyphenols, flavonoids, cardio-glycosides, and steroids.

In this study, the *n*-hexane extract was not qualitatively detected to contain phenolics. Thus, TPC was only determined on ethyl acetate and ethanol extracts while the *n*-hexane extracts were not. TPC of the extracts was determined using gallic acid as a reference. Folin-Ciocalteu (F-C) was used as reagent. The colour product of reaction was measured at a maximum wavelength of 743.50 nm. Based on this study, sequentially extracted

ethanolic extract of *C. trifolia* leaves has total phenolic content (23.79 ± 0.09 mgGAE.g⁻¹) higher than ethyl acetate extract (18.28 ± 0.20 mgGAE.g⁻¹). Anita et al. (2019) reported that ethanolic (80%) extract of *C. trifolia* leaves contain 3.1 mg.g⁻¹ dry weight of phenolic content higher than its stem (2.45 mg.g⁻¹ dry weight)¹⁹.

3.2. Hepatoprotective activity

Evaluation of hepatoprotective activity in rats was determined by measuring liver enzyme levels parameters in rats' blood serum of each group, such as ALT, AST, ALP, bilirubin direct, and total bilirubin (Table. 1).

Table 1. Liver enzymes levels in each rat serum after 14 days treatment.

Groups	ALP U.L. ⁻¹ ±SD	AST (U.I. ⁻¹)±SD	ALT (U.I. ⁻¹)±SD	Bilirubin Direct (mg.dL. ⁻¹)±SD	Total Bilirubin (mg.dL. ⁻¹)±SD
Normal control	66.00±6.63 ^a	50.60±2.40 ^a	20.20±2.77 ^a	0.17±0.01 ^a	0.24±0.02 ^a
Untreated control	139.80±8.81 ^b	94.40±3.84 ^b	38.20±1.92 ^b	0.37±0.04 ^b	0.60±0.01 ^{a,b}
Positive control	76.00±6.30 ^a	62.40±3.84 ^{a,b}	23.80±2.58 ^a	0.22±0.01 ^{a,b}	0.33±0.03 ^{a,b}
Ethyl acetate extract	105.00±9.40 ^{a,b}	73.00±5.09 ^{a,b}	29.40±0.54 ^{a,b}	0.28±0.01 ^{a,b}	0.50±0.02 ^{a,b}
Ethanolic extract	89.00±7.31 ^{a,b}	65.20±1.92 ^{a,b}	24.20±2.77 ^{a,b}	0.24±0.01 ^{a,b}	0.38±0.04 ^{a,b}

Note: Data are reported as mean±SD, n=5. There was a significant difference compared to the untreated control group^{a)} and the normal control group^{b)} ($\alpha=0.05$) (through Tukey test analysis)

Based on Table 1, shows a decrease (sig 0.000, $\alpha < 0.05$) in all liver enzymes parameters in the blood serum of nitrobenzene-induced rats after being given the test materials (both extracts of *C. trifolia* and Silymarin) for 14 days (through one-way ANOVA analysis). As compared to the Silymarin control group, the ethanolic extract-treated group of rats did not show any significant ($\alpha > 0.05$) difference in all liver enzymes (through Tukey test analysis). It means that ethanolic extract of *C. trifolia* has a good potential as a source of hepatoprotective agent approached the silymarin activity as a positive control.

In the present study, the extracts and Silymarin were administered suspension form prepared in 0.5% Na-CMC. Rats in the normal group did not receive Na-CMC. Soldatova et al. (2020) reported that Na-CMC could affect hemostatic and provide an anti-adhesive effect in male Wistar rats at concentrations ranging from 3% to 6%³⁶. To avoid things that are not expected in the normal control group, this group only receives standard pelleted rat feed and drink.

The liver is an essential organ in the body that plays an essential role in vascular function, metabolism (metabolism of carbohydrates, protein, fat, storage of vitamins, detoxification/excretion of drugs, hormones, and other substances), secretory, and excretory. Liver damage is generally associated with the occurrence of liver cell necrosis. The increased lipid peroxidation in tissues which will then have an impact on the liver function itself³⁷. NB is one of the chemical compounds as an oxidant. It contains a single electrophilic nitro group

attached to benzene²⁵. Nitrobenzene can induce hepatotoxic conditions. It caused a significant increase in the serum liver marker enzymes and bilirubin in rats. As an oxidant, nitrobenzene also causes oxidative stress that damages to hepatic cells (necrosis)³⁸. High levels of liver enzymes such as ALP, ALT, AST, and bilirubin are used to evaluate liver damage.

Silymarin (as a reference) has been used to treat liver disorders. It is a naturally occurring chemical compound obtained from *Silybum marianum* (or Milk thistle). Flavolignans, such as silybin, silydianin, and silychristine make up Silymarin. Not only as a liver protector, it is also capable of providing regenerative action. One of the mechanism action is that it decreases the production of free radicals, which may harm cell membranes³⁹. In this study, it is suspected that the phenolic content in *C. trifolia* leaves extract can protect the liver. Identifying of the type of phenolics in this extract needs to be studied further to determine its mechanism of action clearly on its target of action.

3.3. Antioxidant Activity

Previously, Utami et al. (2013) reported that epifriedelanol (a terpenoid) isolated from nonpolar leaves and stem bark extract of *Elaeocarpus floribundus* did not provide antioxidant activity against DPPH radicals ($IC_{50} > 500$ g/mL)⁴⁰. It also strengthens why the *n*-hexane extract was not studied further in this study. The graph in Figure 1. shows that providing both extracts and Silymarin as test materials can reduce MDA levels in




Figure 1. MDA level in each serum control and experimental group after 14 days of treatment. Data are reported as mean \pm SD, n=5. There was a significant difference ($p<0.05$) compared to the normal control group(a) and the untreated control group(b) rats.




Figure 2. Histopathological of the liver in each control and experimental group after 14 days of treatment.

Note: (A) Normal control group (no nitrobenzene-induced, no drugs); (B) Untreated control group (nitrobenzene-induced+0.5% Na-CMC); (C) Silymarin control group (nitrobenzene-induced+Silymarin); (D) nitrobenzene-induced+ethyl acetate extract of *C. trifolia* leaves; (E) nitrobenzene-induced+ethanolic extract of *C. trifolia* leaves. Magnification at 10 \times 40. The red arrow indicates the presence of pyknotic cells.

nitrobenzene-induced rats compared to the untreated (nitrobenzene-induced) control group after 14 days of treatment. In this study, the extracts treated groups showed a significant decrease ($p < 0.05$) in malondialdehyde (MDA) levels compared to the untreated control group (through Tukey test analysis).

In the body, increased concentrations of free radicals and low enzymatic antioxidants cause oxidative stress, one of which is marked by the formation of malondialdehyde (MDA)⁴¹. Lipid peroxidation produces a small amount of MDA, as a major end degradation product⁴² that causes hepatic cell damage and leads to the release of marker enzyme. The level of MDA can be determined using thiobarbiturate acid (TBA) under acidic conditions³⁵. TBA test was performed by extracting MDA using aqueous TCA and then reacting with TBA⁴². In this study, there was a decrease in liver enzyme levels in blood serum and MDA levels after administration of *C. trifolia* extract on nitrobenzene-induced rats.

Phenolic compounds (such as aglycon flavonoids, anthocyanin, catechins, phytoestrogens, and tannin) are important antioxidants in herbal medicine. In order to decrease the HO• radical, antioxidants attack it by giving an electron. It may function as an inhibitor of the production of MDA and MDA-TBA, as seen in the activity test result⁴³. Thus, these compounds play a role in radical scavenging activity that leads to hepatoprotective activity³⁷. Ethanol and aqueous ethanol are non-toxic extraction solvents that have suitable polarity to extract polar compounds, such as phenolic compounds (one of them is flavonoids). The compounds are effective as antioxidants⁴⁴.

3.4. Histopathological Studies in Liver

The presence or absence of pyknotic nuclei might indicate hepatic damage (hepatic necrosis). The pyknotic nuclei is the dead cells, shrunken in shape, with irregular borders, and dark in color⁴⁵. In this study, the average total pycnotic nuclei in the normal, no-treated, Silymarin, ethyl acetate extract, and ethanol extract groups were 25, 72, 33, 52, and 42, respectively. The rat liver tissue from the normal control group shows normal hepatic cells (Figure 2A.). In the untreated control group (Figure 2B), severe hepatotoxicity was demonstrated by the appearance of severe hepatic necrosis. The rats which received ethyl acetate and ethanolic extracts of *C. trifolia* (Figure 2D. and 2E.) showed normal hepatocytes and more minor degree of inflammation, whereas the Silymarin control group (Figure 2C.) showed normal hepatic cells.

5. CONCLUSION

The ethanol extract of *C. trifolia* leaves (200 mg.Kg⁻¹) has good liver protective activity and antioxidant activity

in nitrobenzene-induced rats compared to untreated rats. The phytochemical component in the extract responsible for its hepatoprotective mechanism of action using bio-assay guide isolation needs to be investigated further.

6. ACKNOWLEDGEMENT

The authors thank Research and Development Institute (Lembaga Penelitian dan Pengembangan) of Universitas Muhammadiyah Prof. DR. HAMKA for Funding Grant Scheme of Science and Technology Development Research Batch II 2019.

Conflict of interest

None to declare

Funding

This research and APC were funded by Research and Development Institute (Lembaga Penelitian dan Pengembangan) of Universitas Muhammadiyah Prof. DR. HAMKA grant number: 261/F.03.07/2020.

Ethics approval

This protocol (no. 02/19.10/0230) was approved by the Health Research Ethics Commission of Universitas Muhammadiyah Prof. DR. HAMKA.

Author contribution

Ni Putu Ermi Hikmawanti: Conceptualization, Writing-original draft preparation, Writing-review and editing, Supervision, Project administration.

Tuti Wiyati: Conceptualization, Writing-review and editing, Supervision.

M. Abdul Muis: Resources, Project Administration, Investigation, Visualization

Farah Aisyah Nurfaizah: Resources, Project Administration, Investigation, Visualization

Windy Septiani: Resources, Project Administration, Investigation, Visualization

Ryantama Indrapraja Putra: Resources, Project Administration, Investigation, Visualization

Article info:

Received March 16, 2022

Received in revised form April 29, 2022

Accepted July 26, 2022

REFERENCES

1. Qadrie ZL, Raj Kapoor B, Kavimani S. Hepatoprotective Medicinal Herbs and Animal Models for Their Screening-A Review. *Int J Pharm Sci Res.* 2015;6(12):5006-28.
2. Delgado-Montemayor C, Cordero-Pérez P, Salazar-Aranda R, Waksman-Minsky N. Models of hepatoprotective activity assessment. *Med Univ.* 2015;17(69):222-8.
3. Majaz Q, Khurshid M. Herbal Medicine: A Comprehensive Review. *Int J Pharm Res.* 2016;8(2):1-5.
4. Soni RK, Dixit V, Irchhaiya R, Alok S. Potential herbal hepatoprotective plants: An overview. *Int J Pharm Sci Res.* 2014;5(3):774-89.

5. Kumar D, Kumar S, Gupta J, Arya R, Gupta A. A review on chemical and biological properties of *Cayratia trifolia* Linn. (Vitaceae). *Pharmacogn Rev.* 2011;5(10):184-8.
6. Prasetyo B, Riza L, Mukarlina. Pemanfaatan Tumbuhan Lakum (*Cayratia trifolia* (L.) Domin.) Oleh Etnis Melayu di Kecamatan Sungai Kunyit Kabupaten Mempawah. *Protobiont.* 2016;5(2):25-33.
7. Siriwatanametanon N, Fiebich BL, Efferth T, Prieto JM, Heinrich M. Traditionally used Thai medicinal plants: *In vitro* anti-inflammatory, anticancer and antioxidant activities. *J Ethnopharmacol.* 2010;130:196-207.
8. Kundu JK, Rouf ASS, Nazmul Hossain M, Hasan CM, Rashid MA. Antitumor activity of epifriedelanol from *Vitis trifolia*. *Fitoterapia.* 2000;71:577-9.
9. Gupta A, Bhardwaj A, Gupta J, Bagchi A. Antiimplantation activity of petroleum ether extract of leaves of *Cayratia trifolia* Linn. on female Albino rat. *Asian Pac J Trop Biomed.* 2012;2(S):197-S199.
10. Gupta J, Kumar D, Gupta A. Evaluation of gastric anti-ulcer activity of methanolic extract of *Cayratia trifolia* in experimental animals. *Asian Pacific J Trop Dis.* 2012;2(2):99-102.
11. Chakraborty S, Singha S, Bhattacharya K, Chandra G. Control of human filarial vector, *Culex quinquefasciatus* Say 1823 (Diptera: Culicidae) through bioactive fraction of *Cayratia trifolia* leaf. *Asian Pac J Trop Biomed.* 2013;3(12):980-4.
12. Cruz CP, Alcantara JC, Cruz JP. Antibacterial Property of *Cayratia trifolia* L. as an Alternative Treatment for Boils. *Int J Res Publ.* 2014;3(12):9-12.
13. Romyati, Idiawati N, Destiarti L. Uji Aktivitas Antioksidan, Total Fenol dan Toksisitas dari Ekstrak Daun dan Batang Lakum (*Cayratia trifolia* (L.) Domin). *J Kim Khatulistiwa.* 2014;3(3):30-5.
14. Rabeta MS, Lin SP. Effects of different drying methods on the antioxidant activities of leaves and berries of *Cayratia trifolia*. *Sains Malays.* 2015;44(2):275-80.
15. Feriadi E, Wahyuni, Yusuf MI. Antimitotic Activity of *Cayratia trifolia* Ethanol Extract on Zygote Cells of *Tripleneustes gratilla*. *Pharmacol Clin Pharm Res.* 2016;1(2):69-75.
16. Yusuf MI, Daud NS, H MAA. Antihiperurisemic Activities Of Galing Leaf Ethanol Extract (*Cayratia trifolia* L. Domin) In BALB/C Mice. *J War Farm.* 2019;8(2):20-30.
17. Yusuf MI, Wulaisfan R, Haswika, Wahyuni. Uji Toksisitas Akut dan Gambaran Histopatologi Hepar Mencit yang Diberi Ekstrak Terpurifikasi Daun Galing (*Cayratia trifolia* L. Domin). *Pharmauho J Farm Sains Kesehat.* 2018;4(1):12-5.
18. Perumal PC, Sowmya S, Velmurugan D, Sivaraman T, Gopalakrishnan VK. Assessment of dual inhibitory activity of epifriedelanol isolated from *Cayratia trifolia* against ovarian cancer. *Bangladesh J Pharmacol.* 2016;11(2):545-51.
19. Anita, Jasuja ND, Mathur M. Evaluation of Primary Metabolites and Antioxidant Potential Activity of *Cayratia trifolia* (Leaf and Stems). *J Drug Deliv Ther.* 2019;9(4-A):367-72.
20. Perumal P, Sowmya S, Pratibha P, Vidya B, Anusooriya P, Starlin T, et al. Isolation, structural characterization and *in silico* drug-like properties prediction of a natural compound from the ethanolic extract of *cayratia trifolia* (L.). *Pharmacognosy Res.* 2015;7(1):121-5.
21. Hikmawanti NPE, Wiyati T, Muis MA, Nurfaizah FA, Septiani W. Total Flavonoids Content of Polar Extracts of *Cayratia trifolia* Leaves. *IOP Conf Ser: Earth Environ Sci.* 2021;819:012056.
22. Sowmya S, Perumal PC, Anusooriya P, Vidya B, Pratibha P, Malarvizhi D, et al. Comparative Preliminary Phytochemical Analysis Various Different Parts (Stem, Leaf, Fruit) of *Cayratia trifolis* (L.). *Indo Am J Pharm Res.* 2015;5(01):218-23.
23. Tiwari P, Kumar B, Kaur M, Kaur G, Kaur H. Phytochemical screening and Extraction: A Review. *Int Pharm Sci.* 2011;1(1):98-106.
24. WHO. Quality control methods for herbal materials [document on the Internet]. Geneva: World Health Organization; 2011. Available from: <https://apps.who.int/iris/handle/10665/44479>.
25. Beauchamp RO, Irons RD, Rickert DE, Couch DB, Hamm TE, Lyon JP. A critical review of the literature on nitrobenzene toxicity. *Crit Rev Toxicol.* 1982;11(1):33-84.
26. Hsu CH, Stedeford T, Okoci-Takada E, Ushijima T, Noguchi H, Muro-Chaco C, et al. Framework Analysis for the Carcinogenic Mode of Action of Nitrobenzene. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev.* 2007;25(2):155-84.
27. Yusuf MI, Muthmainnah A, Parawansah. Efek hepatoprotektor ekstrak etanol daun galing (*Cayratia trifolia* L. Domin) pada Tikus Putih (*Rattus norvegicus*). *War Farm.* 2017;6(1):19-27.
28. Guru Kumar D, Sonumul VM, Rathi MA, Thirumoorthi L, Meenakshi P, Gopalakrishnan VK. Hepatoprotective activity of *Cayratia trifolia* (L.) domin against nitrobenzene induced hepatotoxicity. *Lat Am J Pharm.* 2011;30(3):546-9.
29. Ministry of Health Republic of Indonesia. *Farmakope Herbal Indonesia (Indonesian Herb Pharmacopoeia)*. 1st Ed. Jakarta: Ministry of Health Republic of Indonesia; 2008. p. 221.
30. Harborne J, Padmawinata K, Soediro I, Sofia N. *Metode Fitokimia: Penentuan Cara Modern Menganalisis Tumbuhan*. Bandung: ITB Press; 1987.
31. Hanani E. *Analisis Fitokimia*. Jakarta: Buku Kedokteran EGC; 2015. p. 34-45.
32. Hikmawanti NPE, Hanani E, Sapitri Y, Ningrum W. Total Phenolic Content and Antioxidant Activity of Different Extracts of *Cordia sebestena* L. Leaves. *Pharmacogn J.* 2020;12(6):1311-6.
33. Recommendation of the German Society of Clinical Chemistry. Standardization of methods for measurement of enzymatic activities in biological fluids. *Z Klin Chem Klin Biochem.* 1972;10(6):182-91.
34. Jendrassik L, Gróf P. Vereinfachte photometrische Methoden zur Bestimmung des Blutbilirubins. *Biochem Z.* 1938;279:82-9.
35. Esterbauer H, Cheeseman KH. Determination of aldehydic lipid peroxidation products: Malonaldehyde and 4-hydroxynonenal. In: *Methods in Enzymology*. New York: Academic Press. Inc.; 1990. p. 407-21.
36. Soldatova DS, Bezhin AI, Kudryavtseva TN. Study of the effect of the concentration of sodium carboxymethylcellulose on hemostatic and antiadhesive activity during liver operations in an experiment. *Sechenov Med J.* 2020;11(1):4-14.
37. Abou Seif HS. Physiological changes due to hepatotoxicity and the protective role of some medicinal plants. *Beni-Suef Univ J Basic Appl Sci.* 2016;5(2):134-46.
38. Rathi MA, Meenakshi P, Gopalakrishnan VK. Hepatoprotective Activity of Ethanolic Extract of *Alysicarpus vaginalis* against Nitrobenzene-Induced Hepatic Damage in Rats. *South Indian J Biol Sci.* 2015;1(2):60-5.
39. Vargas-Mendoza N, Madrigal-Santillán E, Morales-González Á, Esquivel-Soto J, Esquivel-Chirino C, García-Luna Y, et al. Hepatoprotective effect of silymarin. *World J Hepatol.* 2014;6(3):144-9.
40. Utami R, Khalid N, Sukari MA, Rahmani M, Abdul AB, Dachriyanus. Phenolic contents, antioxidant and cytotoxic activities of *Elaeocarpus floribundus* Blume. *Pak J Pharm Sci.* 2013;26(2):245-50.
41. Gulcin İ. Antioxidants and antioxidant methods: an updated overview. *Arch Toxicol.* 2020;94(3):651-715.
42. Botsoglou NA, Fletouris DJ, Papageorgiou GE, Vassilopoulos VN, Mantis AJ, Trakatellis AG. Rapid, Sensitive, and Specific Thiobarbituric Acid Method for Measuring Lipid Peroxidation in Animal Tissue, Food, and Feedstuff Samples. *J Agric Food Chem.* 1994;42(9):1931-7.
43. Sadeer NB, Montesano D, Albrizio S, Zengin G, Mahomoodally MF. The versatility of antioxidant assays in food science and safety-chemistry, applications, strengths, and limitations. *Antioxidants (Basel).* 2020;9(8):709.
44. Alam MN, Bristi NJ, Rafiquzzaman M. Review on *in vivo* and *in vitro* methods evaluation of antioxidant activity. *Saudi Pharm J.* 2013;21(2):143-52.
45. Wijayanti TRI, Nopiyanti V, Purwidyaningrum IKA. Hepatoprotective Effect of Sukun Leaf Extract (*Artocarpus altilis* (Park.) Fsb.) on Balb/C Mice Induced by Hepatitis B Vaccine. *J Farm Indones.* 2013;10(2):125-32.