

MALAYSIAN JOURNAL OF **NUTRITION**

**Supplement Issue:
FOOD CHOICE AND BEHAVIOUR IN INDONESIA:
FROM UNDER- TO OVER- NUTRITION**



**Southeast Asian Ministers of Education Organization
Regional Centre for Food and Nutrition
www.seameo-recfon.org**

***Guest Editor: Dr Jane Willcox
Supplement Editor: Prof Khor Geok Lin***




VOL. 25 SUPPLEMENT

AUGUST 2019

Official Publication of the
PERSATUAN PEMAKANAN MALAYSIA
NUTRITION SOCIETY OF MALAYSIA

PP18053/02/2013 (033331)



Malaysian Journal of Nutrition is abstracted/indexed by Google Scholar, the WHO Western Pacific Region Index Medicus, Elsevier databases of the Scopus, EBiology and Ecare, ASEAN Citation Index (ACI) and CABI Global Health database

MALAYSIAN JOURNAL OF NUTRITION

Peer-reviewed Journal of the Nutrition Society of Malaysia
(<http://www.nutriweb.org.my>)

EDITOR-IN-CHIEF

Tee E Siong, PhD
President, Nutrition Society of Malaysia

EDITORIAL BOARD

Dr Imelda Angeles-Agdeppa
*(Food and Nutrition Research Institute,
Philippines)*

Assoc Prof Dr Hamid Jan Bin Mohd Jan
(Universiti Sains Malaysia)

Assoc Prof Dr Hazizi Abu Saad
(Universiti Putra Malaysia)

Emeritus Prof Dr Khor Geok Lin
(Universiti Putra Malaysia)

Prof Dr Moy Foong Ming
(University of Malaya)

Assoc Prof Dr Pattanee Winichagoon
(Mahidol University, Thailand)

Prof Dr Poh Bee Koon
(Universiti Kebangsaan Malaysia)

Dr Sangeetha Shyam
*(International Medical University,
Malaysia)*

Prof Dr Suzana Shahar
(Universiti Kebangsaan Malaysia)

Dr Umi Fahmida
*(SEAMEO Regional Centre for Food and
Nutrition, Indonesia)*

Prof Dr Zalilah Mohd Shariff
(Universiti Putra Malaysia)

ADVISORY PANEL

Dr Azza Gohar
(National Nutrition Institute, Egypt)

Prof Cecilia Florencio
(University of The Philippines, Diliman)

Prof Dr JC Henry
(Singapore Institute for Clinical Sciences)

Dr Le Thi Hop
(National Institute of Nutrition, Vietnam)

Assoc Prof Dr Majid Karandish
*(Ahwaz University of Medical Science,
Iran)*

Prof Reynaldo Martorell
*(Emory University, United States of
America)*

Dr V Prakash
*(Central Food Technological Research
Institute, India)*

Dr Siti Muslimatun
*(Indonesia International Institute for Life
Sciences)*

Prof Mark L Wahlqvist
*(Monash University, Australia & National
Health Research Institute, Taiwan)*

The Journal

- Serves as a forum for the sharing of research findings and information across broad areas in nutrition.
- Publishes original research reports, topical article reviews, book reviews, case reports, short communications, invited editorials and letters to the editor.
- Welcomes articles in nutrition and related fields such as dietetics, food science, biotechnology, public health and anthropology.

Malaysian Journal of Nutrition

Vol. 25 Supplement, 2019

Contents

Invited Editorial	Sv
Guest Editor	
<i>Dr Jane Willcox, La Trobe University, Victoria, Australia</i>	
Dietary diversity beliefs and practices among working mothers in Jakarta: a qualitative study	S1
<i>Aria Kekalih, Judhiastuty Februhartanty, Muchtaruddin Mansyur & Anuraj Shankar</i>	
Comparing intake adequacy and dietary diversity between adolescent schoolgirls with normal nutritional status (NG) and undernutrition (UG) based on BMI-for-age (BAZ) living in urban slums in Central Jakarta	S19
<i>Rika Rachmalina, Helda Khusun, Luluk Basri Salim, Luh Ade Ari Wiradnyani & Drupadi HS Dillon</i>	
Food purchasing behaviour among urban slum women in East Jakarta: a qualitative study	S33
<i>Dian Sufyan, Judhiastuty Februhartanty, Saptawati Bardosono, Helda Khusun, Evi Ermayani, Purnawati Hustina Rachman & Anthony Worsley</i>	
Exposure and approval of food marketing strategies: a mixed methods study among household food providers in Jakarta	S47
<i>Selma Avianty, Helda Khusun, Saptawati Bardosono, Judhiastuty Februhartanty & Anthony Worsley</i>	
Association between food marketing exposure and consumption of confectioneries among pre-school children in Jakarta	S63
<i>Maria Meilan Y Fernandez, Judhiastuty Februhartanty & Saptawati Bardosono</i>	
Eating behaviour of young female workers with low socioeconomic status in Malang City, East Java: a qualitative study	S75
<i>Intan Yusuf Habibie, Inge Brouwer & Judhiastuty Februhartanty</i>	
Eating behaviour of adolescent schoolgirls in Malang, East Java: a qualitative study	S87
<i>Mulia Sondari, Inge Brouwer & Judhiastuty Februhartanty</i>	

Perceptions of the causes of obesity among normal weight, overweight and obese Indonesian women: a mixed methods approach	S97
<i>Ismi Irfiyanti Fachruddin, Judhiastuty Februhartanty, Saptawati Bardosono, Helda Khusun & Anthony Worsley</i>	
A qualitative study comparing the coping strategies between food secure and food insecure households of Kaluppini indigenous people in South Sulawesi	S111
<i>Nurbaya, Aria Kekalih & Judhiastuty Februhartanty</i>	
Early development process of drama and storytelling scripts as media for nutrition education on balanced diets among elementary school children	S123
<i>Suci Destriatania, Fenny Etrawati & Judhiastuty Februhartanty</i>	
Acknowledgements	S139

Comparing intake adequacy and dietary diversity between adolescent schoolgirls with normal nutritional status (NG) and undernutrition (UG) based on BMI-for-age (BAZ) living in urban slums in Central Jakarta

Rika Rachmalina^{1,2}, Helda Khusun^{3*}, Luluk Basri Salim², Luh Ade Ari Wiradnyani³ & Drupadi HS Dillon^{2,3}

¹Center for Public Health Research and Development, National Institute of Health Research and Development, Indonesian Ministry of Health, Jakarta; ²Community Nutrition Study Program, Faculty of Medicine Universitas Indonesia; ³SEAMEO Regional Center for Food and Nutrition (RECFON)/Pusat Kajian Gizi Regional Universitas Indonesia (PKGR), Jakarta

ABSTRACT

Introduction: Undernutrition among adolescent girls is an important concern due to their rapid growth velocity that requires adequate intake of energy and nutrients. This study compared intake adequacy and dietary diversity between adolescent public schoolgirls from slum areas in Central Jakarta who had normal and poor nutritional status. **Methods:** A total of 220 eligible girls aged 14–18 years were recruited, with an equal proportion in the normal group (NG) [-1 to +1 SD body mass index-for-age z-score (BAZ)], and undernutrition group (UG) (BAZ < -1SD). Dietary intake was assessed using two non-consecutive 24-hour recalls. Dietary diversity scores (DDS) were determined with reference to the intake of 13 food groups with a minimum daily intake of 15 gram/food group. Receiver operating curve analysis was performed to obtain the DDS cut-off. The Mann-Whitney test was performed to compare DDS between the NG and UG. Logistic regression analysis was conducted to examine the likelihood of potential factors in predicting nutritional status outcome. **Results:** Overall, almost half of the girls' daily food intake showed low dietary diversity based on DDS cut-off <5, with no significant difference between NG and UG adolescents. Protein intake inadequacy showed significant unlikelihood of a NG outcome (OR=0.4; 95% CI: 0.2-0.8), while low socioeconomic status (SES) showed a strong likelihood of an UG (OR=2.7; 95% CI: 1.3-5.5) compared to high SES. **Conclusions:** Low dietary intake and DDS were common among adolescent schoolgirls in slum areas in Jakarta. Nutrition interventions promoting appropriate dietary intake among adolescent girls are recommended.

Keywords: Adolescent girls, dietary diversity, nutrient adequacy, undernutrition, Jakarta slums

*Corresponding author: Helda Khusun
SEAMEO Regional Center for Food and Nutrition (RECFON)–PKGR Universitas Indonesia,
Jakarta, SEAMEO RECFON Building, Jl. Salemba Raya No. 4, Jakarta
Telephone no.: +622131930205; Fax no.: +62213913933
E-mail: hkhusun@seameo-recfon.org; hkhusun@gmail.com

INTRODUCTION

Adolescence is a critical period of physical growth and development. Undernutrition among adolescents is a public health concern in Asian countries with prevalence of > 20% underweight (Cappa *et al.*, 2012). In Jakarta, the prevalence of undernutrition in the form of thinness among adolescents was approximately 11.2% in 2010 (MOH Indonesia, 2013), indicating that adolescent undernutrition is at an unsatisfactory level (WHO, 2010).

Undernutrition during adolescence is of significance given that 50% of the adult weight and skeletal mass and 20% of the adult height are gained during this period. As the rapid growth velocity requires adequate intake of energy and nutrients, adolescents become vulnerable to nutrient deficiencies (Stang & Story, 2005). Adolescent girls in low and middle income countries (LMIC) are often reported to have a monotonous cereal based diet consisting of low nutrient-dense foods, resulting in an inadequate intake of energy and nutrients (USAID & SPRING, 2015). As a consequence, poor nutrition during adolescence affects the ability to learn, reduces work productivity, results in a failure to attain potential height and gain optimal bone mass in adulthood, and delays the onset and progression of puberty.

Several factors are known to lead to undernutrition. These include household food insecurity, intra-household allocation of food that does not meet dietary needs, livelihood insecurity, and poor knowledge of nutrition (WHO, 2005). Poor diet quality could be due to a lack of dietary diversity, which indicates consumption of a low variety of food. This condition has been shown to be associated with micronutrient inadequacy among children and

adolescents (Korkalo *et al.*, 2017; Zhao *et al.*, 2017). A higher household dietary diversity was reported to be associated with a lower likelihood of child stunting (Lee & Ryu, 2018; Mahmudiono, Sumarmi & Rosenkranz, 2017).

Studies determining dietary diversity among Indonesian adolescents in relation to undernutrition are few. This study was aimed at comparing the dietary intake adequacy and dietary diversity between adolescent schoolgirls with normal and poor nutritional status living in urban slum areas in Central Jakarta, Indonesia.

MATERIALS AND METHODS

Study design and subjects

This cross-sectional study was conducted in Central Jakarta. A list of schools was obtained from the Education Office of Central Jakarta, from which five high schools located in slum areas were randomly selected.

Sample size

As studies on dietary diversity of Indonesian adolescent girls are lacking, the sample size for this study was computed based on the results reported by Jayawardena *et al.* (2013), that underweight female ($BMI \leq 18.5$ kg/m²) had lower mean dietary diversity (5.69 ± 1.52) than those whose nutritional status was normal ($BMI > 18.5 - \leq 22.9$ kg/m²) (6.52 ± 1.47). Based on 80% power and a 95% confidence interval, a minimum sample size of 220 was estimated with an equal number for each group, that is 110 for the undernutrition group and another 110 for the normal nutritional status group.

The study inclusion criteria for selecting the participants were girls aged 14-18 years, who were post menarche and apparently healthy. A total of 1,073 schoolgirls from the public schools who

had met the study criteria were selected, and were invited for anthropometric screening.

Nutritional status

Anthropometric measurements were conducted following standard procedures (Gibson, 2005). Height was measured using the ShorrBoard (Weigh and Measure, LLC, USA), and weight was measured using an electronic SECA no. 876 weighing scale (Seca, Germany). The average of two consecutive measurements was used to calculate the body mass index-for-age z-score (BAZ) using WHO AnthroPlus software (WHO, 2009). The girls were classified into two groups based nutritional status. These were the normal group (NG) ($-1SD \leq BAZ \leq +1SD$) and the undernutrition group (UG) ($BAZ < -1SD$) (Thomaz, *et al.*, 2010). The 110 schoolgirls whose BAZ status was ($BAZ < -1SD$) agreed, on a voluntary basis to participate, and were classified to the UG. The other 110 schoolgirls with BAZ status ($-1SD \leq BAZ \leq +1SD$) were placed in the NG.

Data collection

All the eligible participants were interviewed by five trained enumerators, to collect data on dietary diversity, food consumption, morbidity (history of diarrhoea and upper respiratory tract infection in the previous one month), physical activity (PA), household food security, working and education status of mother, and household socioeconomic status (SES). The questionnaires on food consumption and dietary diversity were pre-tested on several adolescent girls from a public high school located in the study area. After pretesting, the questionnaires were revised to improve clarity.

Food consumption

Dietary intake was assessed using two non-consecutive 24-hour (24-h) food

recall periods, comprising one weekday and one day of a weekend. A four-stage multiple-pass interviewing technique was used in the 24-h recall method (Gibson & Ferguson, 2008). The national standardised food photograph book was used for the estimation of portion size (MOH Indonesia, 2014a). The daily nutrient intake was determined using the Indonesian food composition database and calculated by NutriSurvey for Windows, version 2007 (Erhardt, 2007). Energy and protein requirements were calculated by using an estimated energy requirement and protein requirement to specific body weight (FAO, WHO & UNU, 2001; WHO, FAO & UNU, 2007). Fat and carbohydrate adequacy were taken as meeting at least 77% of the Indonesian recommended dietary allowance (RDA) (MOH Decree, 2013). The estimated average requirement was used to evaluate micronutrient adequacy (WHO & FAO, 2006). A dietary intake that was above these requirements was classified as energy and nutrient adequacy.

Dietary diversity

A standardised individual dietary diversity questionnaire was used to obtain the dietary diversity score (DDS) (Arimond *et al.*, 2008). The DDS consisted of 13 food groups, namely starchy staples, legumes and nuts, dairy products, organ meats, eggs, small fish eaten with bone, meat ("flesh foods") and animal protein, vitamin A-rich deep yellow/orange/red vegetables, vitamin A-rich deep green leafy vegetables, vitamin A-rich fruits, vitamin C-rich fruits, vitamin C-rich vegetables, and lastly other fruits and vegetables.

Morbidity (upper respiratory tract infection and diarrhoea)

Participants with upper respiratory tract infection were identified based on medical diagnosis or reports of fever, sore throat, and cough in the previous one

month. Participants with diarrhoea were identified based on doctor's diagnosis or had experienced passing liquid or loose stools three or more times in the previous one month.

Physical activity

A short form of the international PA questionnaire (IPAQ) was used to determine PA during the past seven days (IPAQ, 2005).

Household food security

Household food security status was determined by using the food security survey module for children aged ≥ 12 years. The nine questions in the module about food situation at home during the past one month were answered by participants. Response to the questions were assumed to be an indication of the food security status of the children, as perceived by the family (Connell *et al.*, 2004).

Socioeconomic status

The SES of the household was determined based on ownership of assets consisting of the sources of drinking water, electricity and cooking fuel, ownership of toilet, type of latrine, final faecal disposal, ownership of a motorcycle, television, air-conditioner, water heater, 12 kg cooking gas cylinder, refrigerator, and car (MOH Indonesia, 2013).

Data analysis

The DDS was computed by assigning a score of one (1) for the consumption of at least 15g/day of a food group, and zero (0) score for intake < 15 g/day. The total score for the entire food groups ranged from 0-13. The receiver operating characteristic (ROC) curve was plotted to obtain the DDS cut-off corresponding to the nutritional status of the schoolgirls. For this purpose, we contrasted the DDS with the composite score for the

intake adequacy of energy and protein. The procedure yielded a DDS of 5 as the cut-off for dietary diversity with the area under the curve (AUC) of 0.65, sensitivity of 60%, and specificity of 64% ($p = 0.002$). This cut-off was also used in examining the relationship between the DDS and nutritional status.

PA was analysed according to the IPAQ guidelines. The subjects were requested to recall the duration of usual their PA in a week. The duration of these activities was then converted into metabolic equivalent (MET) – minutes per week and categorised into high, moderate, and low PA based on the IPAQ guidelines (IPAQ, 2005).

Household food security status was categorised as food-secure or food-insecure, based on the responses to the nine questions in the module. We examined under- and over- reporting of energy intake to check for potential bias in participant's dietary intake (McCrory *et al.*, 2002).

SES was constructed based on 13 variables that were used to indicate the wealth index (MOH Indonesia, 2013). By using principal component analysis (PCA), a reliability analysis was first conducted, yielding ten variables, which were screened providing a Cronbach's alpha of 0.686. The PCA yielded a correlation score of > 0.6 for the SES variables. The scores were then ranked into tertiles, where tertile one was the lowest and tertile three the highest SES.

The independent *t*-test was used to compare the differences in height and body weight between the two nutritional status groups (NG vs UG). The Mann-Whitney test was used to compare the differences in dietary intake and DDS between these two groups. The chi-square test was used to examine the independence between DDS (cut-off < 5 and ≥ 5 food groups) and nutritional status (NG vs UG). Logistic regression

analysis was undertaken to assess the relationship between the categorical potential predictive factors: DDS, intake adequacy (energy and protein adequacy), morbidity status (history of diarrhoea and upper respiratory tract infection in the previous one month), household food security status, and SES, in predicting the categorical outcome: normal nutritional status versus undernutrition. The potential predictors were selected according to the conceptual framework of nutritional problems and causal factors during adolescence; dietary inadequacies and infectious diseases are immediate cause of undernutrition in adolescence (WHO, 2005). The results of the logistic regression were expressed as the odds ratio and 95% confidence interval. Statistical analyses were performed using SPSS (version 20). *P* values <0.05 were considered statistically significant.

Ethical approval

The study protocol was approved by the Research Ethics Committee of the Medical Faculty of Universitas Indonesia in Jakarta (reference number 206/UN.2.F1/ETIK/2015, dated 16 March 2015). The subjects provided written consent prior to data collection.

RESULTS

The study was conducted from March to April in 2015. Out of a total of 220 eligible participants who were selected, 25 were excluded from data analysis due to under- or over-reporting of energy intake. The final analysis was performed on 195 participants, consisting of 100 adolescents in the UG and 95 in the NG.

The mean body weight and height were significantly different between the NG and UG. The mean body weight of the NG (49.3 ± 5.1 kg) was significantly higher than that of the UG (40.8 ± 4.0 kg)

(Table 1). However, the mean height of the NG at 1.53 ± 0.06 m was significantly less than that of the UG at 1.56 ± 0.06 m. This indicates that on average, the UG was thinner but somewhat taller than the NG. In line with this, the proportion of stunting was significantly higher in the NG compared to the UG (24.4% vs 12.0%, respectively; *p*=0.029).

The SES of the UG was worse off than that of the NG, with the former having a significantly higher proportion with low SES (50.0% vs 31.6%, respectively) (Table 1). Overall (i.e. NG and UG), more than half of the adolescent schoolgirls (55.4%) reported having upper respiratory tract infection while a lower proportion (12.3%) had diarrhoea in the previous month. There were no statistically significant differences in the morbidity status between the NG and UG. There was also no significant difference between the two groups in terms of the prevalence of household food insecurity (UG 60.0% vs NG 50.5%), and the proportion of working mothers (UG 25.5% vs 37.6% NG).

The median daily intake of energy, protein, and fat were approximately 1,500 kcal, 49 g, and 62g, respectively; these figures were 72.0%, 83.0% and 87.0%, respectively, of the Indonesian RDA (Table 2). There were no statistically significant differences in the intake of energy, macro- and micro- nutrients between the UG and NG.

Table 3 shows that, overall, <10.0% of the adolescents had adequate energy intake and >60.0% of them attained adequate intake of protein, fat, vitamin A and vitamin B₆. Nonetheless, very low proportions of the adolescents had adequate intakes of folate, calcium and zinc. There were no significant differences in energy and nutrient intake adequacy between the UG and NG, except for protein intake adequacy at 83.0% vs 65.3%, respectively.

Table 1. Anthropometric measurements, morbidity, household food security, and socioeconomic status of the adolescent schoolgirls according to nutritional status

	All (n=195)	Nutritional status		p [†]
		Undernutrition (BAZ<-1SD) (n=100)	Normal (-1SD ≤ BAZ ≤ 1SD) (n=95)	
Height, m (mean±SD)	1.55±0.06	1.56±0.06	1.53±0.06	0.06*
Weight, kg (mean±SD)	44.9±5.8	40.8±4.0	49.3±5.1	<0.001***
Stunting (Height-for-age z-score <-2.0) (%)	17.9	12.0	24.2	0.029*
Had upper respiratory tract infection in the previous month (%)	55.4	56.0	54.7	0.859
Had diarrhoea in the previous month (%)	12.3	10.0	14.7	0.317
Physical activity level [‡] (%)				
Low	33.3	32.0	34.7	0.976
Moderate	64.6	66.0	63.2	
High	2.1	2.0	2.1	
Household food security status [§] (%)				
Food secure	44.6	40.0	49.5	0.184
Food insecure	55.4	60.0	50.5	
Adolescent girls with working mother (n=191) (%)	31.3	25.5	37.6	0.073
Mother educational level (n=191) (%)				
Lower education (graduated from junior high school or less)	26.7	26.5	26.9	0.956
Higher education (attending high school or more)	73.3	73.5	73.1	
Socioeconomic status [¶] (%)				
Low	41.0	50.0	31.6	0.005**
Middle	24.6	24.0	25.3	
High	34.4	26.0	43.2	

[†]T-test for continuous variables and chi-square test for discrete variable

[‡]Based on average MET-minutes per week (IPAQ, 2005)

[§]The sum of affirmative responses to the nine questions in the food security module for children aged ≥12 years (Connell et al., 2004)

[¶]SES was defined based on tertiles of wealth index score (household ownership of assets) (MOH Indonesia, 2013)

*p<0.05, **p<0.01, ***p<0.001

Table 2. Energy and nutrient intake among the adolescent schoolgirls according to nutritional status

	All (n=195)	Nutritional status		p [†]
		Undernutrition (BAZ<-1SD) (n=100)	Normal (-1SD ≤ BAZ ≤ 1SD) (n=95)	
		Median (IQR [‡])		
Energy (kcal/day)	1531.3 (1265.3–1865.1)	1556.6 (1281.2–1853.9)	1478.7 (1260.9–1883.9)	0.594
Protein (g/day)	48.9 (38.3–62.5)	49.0 (37.8–63.3)	48.8 (38.7–62.2)	0.744
Fat (g/day)	61.9 (47.5–79.1)	62.3 (50.6–80.8)	59.6 (46.1–77.6)	0.455
Carbohydrate (g/day)	198.4 (158.0–252.5)	205.2 (162.4–253.2)	197.5 (153.9–247.9)	0.385
Vitamin A (µg/day)	1068.1 (723.6–1627.2)	1059.3 (739.9–1615.9)	1082.5 (684.7–1627.2)	0.823
Thiamine (mg/day)	0.6 (0.4–0.9)	0.7 (0.5–0.9)	0.6 (0.4–0.9)	0.366
Riboflavin (mg/day)	0.8 (0.6–1)	0.8 (0.6–1.0)	0.8 (0.5–1.0)	0.359
Niacin (mg/day)	9.1 (6.9–12.5)	8.9 (6.4–12.6)	9.1 (7.1–12.4)	0.657
Vitamin B ₆ (mg/day)	1.2 (0.8–1.8)	1.2 (0.8–1.7)	1.1 (0.8–1.8)	0.539
Folate (µg/day)	124.0 (83.0–181.0)	130.0 (88.0–191.0)	118.0 (77.0–174.0)	0.080
Vitamin B ₁₂ (µg/day)	1.9 (1.3–3.5)	2.0 (1.4–3.5)	1.9 (1.2–3.5)	0.732
Vitamin C (mg/day)	24.1 (13.1–44.6)	24.9 (13.8–48.7)	21.9 (11.2–44.5)	0.339
Calcium (mg/day)	294.4 (219.5–421.2)	307.1 (229.3–420.3)	281.8 (190.7–426.6)	0.335
Zinc (mg/day)	6.5 (5.2–8.2)	6.3 (5.3–8.3)	6.6 (5.2–8.1)	0.114
Iron (mg/day)	7.7 (5.4–8.2)	8.0 (6.0–10.2)	7.2 (4.9–9.9)	0.971

[†]Mann-Whitney test[‡]IQR=Interquartile range

Table 3. Percentage of the adolescent schoolgirls according to nutritional status meeting energy and nutrient intake adequacy[†]

	Nutritional status			<i>p</i> [‡]
	All (<i>n</i> =195)	Undernutrition (BAZ<-1SD) (<i>n</i> =100)	Normal (-1SD ≤ BAZ ≤ 1SD) (<i>n</i> =95)	
Energy	9.7	12.0	7.4	0.276
Protein	74.4	83.0	65.3	0.005**
Fat	63.1	66.0	60.0	0.386
Carbohydrate	34.9	38.0	31.6	0.347
Vitamin A	92.8	95.0	90.5	0.226
Thiamine	22.6	25.0	20.0	0.404
Riboflavin	17.4	20.0	14.7	0.333
Niacin	47.7	47.0	48.4	0.843
Vitamin B ₆	61.0	63.0	58.9	0.562
Folate	6.2	7.0	5.3	0.614
Vitamin B ₁₂	49.2	50.0	48.4	0.826
Vitamin C	19.0	20.0	17.9	0.708
Calcium	0.5	1.0	0.0	0.328
Zinc	4.6	6.0	3.2	0.344

[†]Energy and protein requirements were calculated by using estimated energy requirement and protein requirement to specific body weight (FAO, WHO & UNU, 2001; WHO, FAO & UNU, 2007). Fat and carbohydrate adequacy were taken as meeting at least 77% of the Indonesian RDA (MOH Decree, 2013). Estimated average requirement was used to evaluate micronutrient adequacy (WHO & FAO, 2006). Dietary intake above these requirements was classified as energy and nutrient adequacy.

[‡]Chi-square test

***p*<0.01

Based on a DDS cut-off of <5, almost half of all the girls (46.2%) daily consumed a low diversity of foods, with no significant difference between the NG and UG (Table 4). The food groups consumed by the majority of all the adolescents on a daily basis were starchy staples (100%), flesh foods and animal protein (85.1%), legumes and nuts (55.9%) and egg (54.4%). They consumed a low percentage of fruits, only about one-third reported taking vegetables, on a daily basis. Significant differences were observed for the daily intake of legumes and nuts (NG 42.1% vs UG 69.0%), and vitamin C-rich vegetables (NG 14.7% vs UG 28.0%). Higher intake of legumes and nuts by the UG may explain the finding of significant

difference of protein intake adequacy between this group and the NG, as shown in Table 3. Daily consumption of the other food groups did not show significant differences between the NG and UG girls.

The logistic regression analysis showed that among the factors studied, two were found to have significant influence on the nutritional status outcome of the adolescent schoolgirls living in the slum areas of Jakarta. Household SES and protein intake adequacy were significantly associated with the nutritional status of the schoolgirls (Table 5). Protein intake inadequacy was associated with a less likelihood of attaining normal nutritional status (OR=0.4; 95% CI: 0.2-0.8;

Table 4. Distribution (%) of dietary diversity scores (DDS) and food groups consumed by the adolescent schoolgirls according to nutritional status

	All (n=195)	Nutritional status		p [†]
		Undernutrition (BAZ<-1SD) (n=100)	Normal (-1SD≤BAZ≤1SD) (n=95)	
Dietary diversity score, median (IQR)	5.0 (4.0–6.0)	5.0 (3.0–6.0)	5.0 (4.0–6.0)	0.113
DDS <5 (%)	46.2	43.0	49.5	0.365
DDS ≥5 (%)	53.8	57.0	50.5	
Food groups (%)				
Starchy staples	100.0	100.0	100.0	1.000
Legumes and nuts	55.9	69.0	42.1	<0.01**
Dairy products	42.1	41.0	43.2	0.761
Organ meats	5.1	6.0	4.2	0.572
Eggs	54.4	57.0	51.6	0.449
Small fish eaten with bone	4.6	3.0	6.3	0.271
Flesh foods and animal protein	85.1	81.0	89.5	0.097
Vitamin A-rich deep yellow/orange/red vegetables	33.8	33.0	34.7	0.798
Vitamin A-rich deep green leafy vegetables	23.1	22.0	24.2	0.715
Vitamin A-rich fruits	6.2	5.0	7.4	0.493
Vitamin C-rich vegetables	21.5	28.0	14.7	0.025*
Vitamin C-rich fruits	15.4	58.0	63.2	0.152
Other fruits and vegetables	39.5	42.0	36.8	0.463

†Chi-square test; Mann-Whitney test for dietary diversity score

*p<0.05

**p<0.01

Table 5. Logistic regression analysis in predicting nutritional status of the adolescent girls (n=195) according to potential factors

<i>Potential factors</i>	<i>Normal nutritional status vs undernutrition[†]</i>		
	<i>OR</i>	<i>95% CI</i>	<i>p</i>
Dietary diversity score (DDS)			
DDS ≥5 (reference)			
DDS <5	1.1	0.6-2.0	0.805
Dietary intake			
Energy intake adequacy (reference)			
Energy intake inadequacy	0.7	0.3-1.9	0.494
Protein intake adequacy (reference)			
Protein intake inadequacy	0.4	0.2-0.8	0.006**
Morbidity status			
Had upper respiratory tract infection			
Without upper respiratory tract infection (reference)	1.0	0.6-1.9	0.899
Had diarrhoea			
Without diarrhoea (reference)	0.6	0.3-1.6	0.328
Household food security			
Food secure (reference)			
Food insecure	1.3	0.7-2.4	0.405
Socioeconomic status			
High SES (reference)			
Middle SES	1.5	0.7-3.2	0.323
Low SES	2.7	1.3-5.5	0.006**

[†]Normal BMI-for-age score or BAZ: (-1SD ≤ BAZ ≤ +1SD) as reference vs undernutrition (BAZ < -1SD)

***p*<0.01

p=0.006). Schoolgirls from households with low SES were 2.7 times more likely to be undernourished than those from households with high SES (OR=2.7; 95% CI: 1.3-5.5; *p*=0.006). The other factors that were studied including DDS, energy intake, morbidity status, and household food security were not found to exert a significant influence on the nutritional status outcome of the adolescent schoolgirls.

DISCUSSION

Data on dietary intake and dietary diversity among adolescents are limited

in Indonesia. Interventions to improve the dietary intake of adolescents are also lacking. The results of this study provide some insights into the quality and variety of food consumed by adolescent schoolgirls from slum areas in Central Jakarta.

On the average, Indonesian adolescents showed a lower DDS (less than five food groups) than some other countries. By comparison, Iranian adolescent girls consumed an average of approximately six food groups (range 5–14 food groups) (Akbari & Azadbakht, 2014), and the mean DDS was 5.76 for

urban adolescent schoolgirls in Ethiopia with 76% of them having adequate dietary diversity (Birru, Tariku & Belew, 2018).

The diet of the adolescent schoolgirls in this study was predominantly based on starchy staples. More than half of the girls reported taking animal protein foods and legumes and nuts. A low percentage of them consumed fruit and vegetables on a daily basis, when compared to the finding of the national survey in Indonesia, which reported vegetable consumption among Indonesian adolescents aged 13–18 years at 94.7% (Hermina & Prihartini, 2016). In studies on adolescents conducted in Africa and Canada, low energy and nutrient intake were generally found in adolescents from households with low SES and food insecurity (Kirkpatrick & Tarasuk, 2008; Dapi *et al.*, 2010). In low income households, common barriers to low fruit and vegetable intake were the unavailability and poor access to affordable types of fruit and vegetables, a lack of knowledge about healthy foods, the poor quality of the produce, and budgetary constraints (Huang, Edirisinghe & Burton-Freeman, 2016). A poor knowledge of nutrition among adolescent school girls was reported by a study conducted in 12 districts of Indonesia, which showed that less than half of adolescents aged 10–19 years were aware of health benefits of fruits (43.7%) and vegetables (36.2%) (Sudirman & Jahari, 2012). Therefore, disseminating knowledge of the health benefits of fruits and vegetables is essential. The public sector should enable people to have better access to reduce retail prices so that a wider variety of foods is affordable to all socioeconomic strata in a community (Nair, Augustine & Konapur, 2016).

Several studies have revealed that dietary diversity is consistently

associated with micronutrient adequacy in children and adolescents (Korkalo *et al.*, 2017; Zhao *et al.*, 2017). Women of reproductive age in five developing countries reported that dietary diversity consistently predicts micronutrient adequacy (Arimond *et al.*, 2010).

This study showed that the adolescents had low nutrient intake, particularly of vitamin B₁, vitamin B₂, folate, vitamin C, calcium, and zinc. Macronutrient intake, except for fat, was below the national requirements for Indonesian adolescents aged 14–18 (MOH Indonesia, 2014b). The latest Indonesian dietary survey revealed that the highest proportion of energy inadequacy was among adolescents aged 13–18 (MOH Indonesia, 2014b). This is consistent with the literature regarding dietary intake among adolescents in developing countries, which has highlighted the poor diet quality in this age group (Ochola & Masibo, 2014). Their diets are known to be limited in diversity, particularly in the fruit and vegetable food groups (Zhao *et al.*, 2017). Further, energy and micronutrient intake were found to be inadequate in the majority of adolescents in developing countries (Ochola & Masibo, 2014). These findings indicate that nutrition policies and programmes are important to improve the food intake of adolescents for growth, cognition, and educational achievements (Ochola & Masibo, 2014).

This study found that low household SES had a strong influence on the nutritional status of the adolescents. This is consistent with findings elsewhere in other groups such as that of a study of pregnant women in Kenya, which reported that socioeconomic factors including employment status, household assets, and land ownership influenced dietary diversity in pregnant women (Kiboi, Kimiywe & Chege, 2017).

CONCLUSION

The diet of adolescent schoolgirls living in slum areas in Jakarta was inadequate in terms of dietary adequacy and diversity. The Indonesian Ministry of Health issued dietary guidelines in the 2014 on principles of a balanced diet. However, these guidelines may not be well disseminated, and key elements of it, including “consume a variety of foods”, are not widely known or understood. Interventions directed at promoting good diets through dietary diversification among adolescent schoolgirls, are recommended for schools and the community at large.

Acknowledgements

The authors are grateful to all the investigators and adolescent schoolgirls who took part in this study.

Authors' contributions

RR, designed research, carried out data collection, analysed and interpreted the data, and developed manuscript; HK, designed research, interpreted the data and critically reviewed the manuscript; LBS, carried out data collection; LAAW, critically reviewed the manuscript; DHSD, designed research and critically reviewed the manuscript. All authors have seen and approved the final manuscript.

Conflict of interest

The authors declare no conflict of interest.

References

- Akbari F & Azadbakht L (2014). A systematic review on diet quality among Iranian Youth: Focusing on reports from Tehran and Isfahan. *Archives of Iranian Medicine* 17(8):574–584. <https://doi.org/014178/aim.0010>.
- Arimond M, Elin L, Wiesmann D, Joseph M & Carriquiry A (2008). *Dietary Diversity as a Measure of Women's Diet Quality in Resource-Poor Areas: Results from rural Bangladesh site*. Food and Nutrition Technical Assistance (FANTA) Project/Academy for Educational Development (AED), Washington DC.
- Arimond M, Wiesmann D, Becquey E, Carriquiry A, Daniels MC, Deitchler M, Fanou-Fognny N, Joseph ML, Kennedy G, Martin-Prevel Y & Torheim LE (2010). Simple Food Group Diversity Indicators Predict Micronutrient Adequacy of Women's Diets in, 2059–2069. *J Nutr* 140(11):2059S–2069S. <https://doi.org/10.3945/jn.110.123414.2059S>.
- Birru SM, Tariku A & Belew AK (2018). Improved dietary diversity of school adolescent girls in the context of urban Northwest Ethiopia: 2017. *Italian Journal of Pediatrics* 44(1):48. <https://doi.org/10.1186/s13052-018-0490-0>.
- Cappa C, Wardlaw T, Langevin-Falcon C & Diers J (2012). *Progress for Children - A report card on adolescents*. United Nations publication, New York.
- Connell CL, Nord M, Lofton KL & Yadrick K (2004). Food security of older children can be assessed using a standardized survey instrument. *The Journal of Nutrition* 134(10):2566–2572.
- Dapi LN, Hörnell A, Janlert U, Stenlund H & Larsson C (2010). Energy and nutrient intakes in relation to sex and socio-economic status among school adolescents in urban Cameroon, Africa. *Public Health Nutrition* 14(5): 904–913. <https://doi.org/10.1017/S1368980010003150>.
- Erhardt J (2007). *Nutrition surveys and calculations*. From <http://www.nutrisurvey.de/>. [Retrieved February 4 2015].
- FAO, WHO & UNU (2001). *Human energy requirements. Scientific background papers from the Joint FAO/WHO/UNU Expert Consultation*. From <http://www.ncbi.nlm.nih.gov/pubmed/16277811>. [Retrieved January 12 2015].
- Gibson RS (2005). *Principles of nutritional assessment*. Oxford University Press, New York.
- Gibson RS & Ferguson EL (2008). *An interactive 24-hour recall for assessing the adequacy of iron and zinc intakes in developing countries* (Vol. 8). HarvestPlus Technical Monograph 8, Washington DC. <https://doi.org/10.1007/BF02927624>.
- Hermiana & Prihartini S (2016). *Fruits and vegetables consumption of Indonesian population in the context of balanced nutrition: a further analysis of Individual Food Consumption Survey (SKMI) 2014*. From <https://media.neliti.com/media/publications/67991-ID-gambaran-konsumsi-sayur-dan-buah-pendudu.pdf>. [Retrieved January 26 2019].

- Huang Y, Edirisinghe I & Burton-Freeman BM (2016). Low-income shoppers and fruit and vegetables: What do they think? *Nutrition Today* 51(5):242–250. <https://doi.org/10.1097/NT.000000000000176>.
- IPAQ (2005). *Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ) – Short and Long Forms Questionnaire*. From <https://sites.google.com/site/theipaq/scoring-protocol>. [Retrieved May 28 2015].
- Jayawardena R, Byrne NM, Soares MJ, Katulanda P, Yadav B & Hills AP (2013). High dietary diversity is associated with obesity in Sri Lankan adults: an evaluation of three dietary scores. *BMC Public Health* 13:314. <https://doi.org/10.1186/1471-2458-13-314>.
- Kiboi W, Kimiywe J & Chege P (2017). Determinants of dietary diversity among pregnant women in Laikipia County, Kenya: a cross-sectional study. *BMC Nutrition* 3(1):12. <https://doi.org/10.1186/s40795-017-0126-6>.
- Kirkpatrick SI & Tarasuk V (2008). Food insecurity is associated with nutrient inadequacies among Canadian adults and adolescents. *The Journal of Nutrition* 138(3):604–612.
- Korkalo L, Erkkola M, Heinonen AE, Freese R, Selvester K & Mutanen M (2017). Associations of dietary diversity scores and micronutrient status in adolescent Mozambican girls. *European Journal of Nutrition* 56(3):1179–1189. <https://doi.org/10.1007/s00394-016-1167-3>.
- Lee SJ & Ryu HK (2018). Relationship between dietary intakes and the double burden of malnutrition in adults of Malang, Indonesia: An exploratory study. *Nutrition Research and Practice* 12(5):426–435. <https://doi.org/10.4162/nrp.2018.12.5.426>.
- Mahmudiono T, Sumarmi S & Rosenkranz RR (2017). Household dietary diversity and child stunting in East Java, Indonesia. *Asia Pacific Journal of Clinical Nutrition* 26(2):317–325. <https://doi.org/10.6133/apjcn.012016.01>.
- McCrory MA, McCrory MA, Hajduk CL & Roberts SB (2002). Procedures for screening out inaccurate reports of dietary energy intake. *Public Health Nutrition* 5(6A): 873–882. <https://doi.org/10.1079/PHN2002387>.
- MOH Decree (2013). *Angka Kecukupan Gizi Yang Dianjurkan Bagi Bangsa Indonesia, Pub. L. No. Nomor 75 Tahun 2013*. Menteri Kesehatan Republik Indonesia, Jakarta.
- MOH Indonesia (2013). *Basic Health Research 2013*. NIHRD Press, Jakarta.
- MOH Indonesia (2014a). *Food photograph book: Individual food consumption survey*. NIHRD Press, Jakarta.
- MOH Indonesia (2014b). *Individual food consumption survey Indonesia 2014*. NIHRD Press, Jakarta.
- Nair MK, Augustine LF & Konapur A (2016). Food-Based Interventions to Modify Diet Quality and Diversity to Address Multiple Micronutrient Deficiency. *Frontiers in Public Health* 3(January):1–14. <https://doi.org/10.3389/fpubh.2015.00277>.
- Ochola S & Masibo PK (2014). Dietary Intake of Schoolchildren and Adolescents in Developing Countries. *Annals of Nutrition and Metabolism* 64(s2):24–40. <https://doi.org/10.1159/000365125>.
- Stang J & Story M (2005). *Guidelines for Adolescent Nutrition Services*. University of Minnesota, Minneapolis.
- Sudirman H & Jahari AB (2012). Pengetahuan, sikap dan perilaku remaja tentang keluarga sadar gizi (Kadarzi): dengan perhatian khusus terhadap pantauan berat badan dan mengonsumsi makanan beragam. *Media Penelitian dan Pengembangan Kesehatan* 22(2):93–105. From <http://ejournal.litbang.depkes.go.id/index.php/MPK/article/view/2632/615>. [Retrieved January 26 2019].
- Thomaz EBAF, Cangussu MCT, da Silva AAM & Assis AMO (2010). Is malnutrition associated with crowding in permanent dentition? *International Journal of Environmental Research and Public Health* 7(9):3531–3544. <https://doi.org/10.3390/ijerph7093531>.
- USAID & SPRING (2015). *Nutrition of Adolescent Girls and Women of Reproductive Age in Low- and Middle-Income Countries: Current Context and Scientific Basis for Moving Forward*. From www.spring-nutrition.org. [Retrieved September 27 2018].
- WHO (2005). *Nutrition in adolescence – Issues and Challenges for the Health Sector*. WHO Library Cataloguing, Geneva.
- WHO (2009). *World Health Organization. AnthroPlus for Personal Computers. Manual: Software for assessing growth of the world's children*. From <http://www.who.int/growthref/tools/en/>. [Retrieved November 15 2016].

- WHO (2010). *Nutritional Landscape Information System: Country Profile Indicators: Interpretation Guide*. WHO Press, Geneva.
- WHO & FAO (2006). *Guidelines on food fortification with micronutrients*. In L Allen, B De Benoist, O Dary & R Hurrell (Eds.). WHO, Geneva.
- WHO, FAO & UNU (2007). *Protein and amino acid requirements in human nutrition*. From <http://www.ncbi.nlm.nih.gov/pubmed/18330140>. [Retrieved January 12 2015].
- Zhao W, Yu K, Tan S, Zheng Y, Zhao A, Wang P & Zhang Y (2017). Dietary diversity scores: An indicator of micronutrient inadequacy instead of obesity for Chinese children. *BMC Public Health* 17(1):1–11. <https://doi.org/10.1186/s12889-017-4381-x>.