




Search by any title, abstract, authors...

# Research.Publish.Connect.



Search by any title, abstract, authors...

## Subjects (#papers)

<b>Artificial Intelligence</b> (8064)	<b>Biomedical Engineering</b> (5015)	<b>Cloud Computing</b> (1275)
<b>Computational Intelligence</b> (2009)	<b>Computer Vision, Visualization and Computer Graphics</b> (4939)	<b>Computer-Supported Education</b> (1959)
<b>Data Engineering</b> (3054)	<b>Energy and Economy</b> (377)	<b>Enterprise Information Systems</b> (8769)
<b>Health Engineering and Technology Applications</b> (2029)	<b>Human-Computer Interaction</b> (2058)	<b>Informatics in Control, Automation and Robotics</b> (4330)

**Information and Systems Security**  
(1205)

**Knowledge-Based  
Systems (4384)**

**Operational Research**  
(914)

**Pattern Recognition**  
(3999)

**Photonics, Optics and Laser  
Technology (269)**

**Sensor Networks**  
(2467)

**Simulation and Modeling**  
(1798)

**Software Engineering**  
(5132)

**Sport Science Research and  
Technology (241)**

**Telecommunications**  
(2235)

**Web Information Systems and  
Technologies (3424)**

## Partners



**Proceedings will be submitted for  
indexation by:**



## RESOURCES

Proceedings

Papers

Authors

Ontology

## CONTACTS

Science and Technology Publications, Lda  
Avenida de S. Francisco Xavier, Lote 7 Cv. C,  
2900-616 Setúbal, Portugal.

Phone: +351 265 520 185

Fax: +351 265 520 186

Email: [info@scitepress.org](mailto:info@scitepress.org)

## EXTERNAL LINKS

PRIMORIS

INSTICC

SCITEVENTS

CROSSREF

## PROCEEDINGS SUBMITTED FOR INDEXATION BY:

dblp

Ei Compendex

SCOPUS

Semantic Scholar

Google Scholar

Microsoft Academic



Search by any title, abstract, authors...



*Research.Publish.Connect.*

## Proceedings

### Proceedings of the 1st Muhammadiyah International Conference on Health and Pharmaceutical Development

August 10-11, 2018, in East Jakarta, Indonesia



**Editors:** Endang Hanani ; Etin Diah Permanasari ; Landyyun Rahmawan Sjahid ; Lusi Putri Dwita ; Daniek Viviandhari ; Rindita ; Nindya Herdianti ; Numlil Khaira Rusdi ; Nuriza Rahmadini and Wati Sukmawati

**Affiliation:** Universitas Muhammadiyah Prof. DR. HAMKA, Indonesia

**ISBN:** 978-989-758-349-0


**Foreword:** This book contains the proceedings of the first Muhammadiyah International Conference on Health and Pharmaceutical Development (MICH-PhD). MICH-PhD was held on 10-11 Augustus 2018 in Jakarta, Indonesia. The conference organized by Faculty of Pharmacy and Science, Universitas Muhammadiyah Prof. DR. HAMKA (UHAMKA) in collaboration with Association of Muhammadiyah Pharmacy Higher Education (Universitas Muhammadiyah Surakarta, Universitas Ahmad Dahlan, Universitas Muhammadiyah Malang, Universitas Muhammadiyah Banjarmasin, Universitas Muhammadiyah Purwokerto, Universitas Muhammadiyah Yogyakarta, Sekolah Tinggi Farmasi Muhammadiyah Tangerang, Sekolah Tinggi Farmasi Muhammadiyah Cirebon, STIKES Muhammadiyah Pekajangan, STIKES Muhammadiyah Gombong and Universitas Muhammadiyah Magelang). The aim of this conference is to provide opportunities for participants to present their research and to expose participants to the development and innovation made in the field of health and various pharmaceutical disciplines. The technical program in the conference includes several scientific talks by the experts, oral presentations and poster presentations. During two-day conference, researchers presented the latest development in pharmacy as well as established networking for possible collaborating research among participant. MICH-PhD received 114 abstracts which 70% presented as oral and poster presentations. The range of topics covered by in the 1st MICH-PhD Proceeding includes Pharmacology, Toxicology, Immunology, Histology, Molecular biology, Natural products, Microbiology, Biochemistry, Molecular docking, Medicinal chemistry, Clinical research, Pharmacoeconomics, Pharmacovigilance, Drug design, Drug delivery vehicle, Pharma-nanotechnology, Biopharmaceuticals and Drug formulation. To evaluate each submission, a blind review was performed by the Scientific Committee, whose members are highly qualified researchers in MICH-PhD topic areas. Not all papers presented at the conference included in this book of proceeding, only the papers stated accepted after the review process. In addition, the papers selected in this proceeding were also passed through the proofreading and editing process. This international conference can be carried out with the help of various parties, and we are very thankful for that. First of all: We would like to thank the UHAMKA leaders and Association of Muhammadiyah Pharmacy Higher Education for fully supporting this event. Secondly: to the authors whose quality of work is at the core of this conference, without them there will be no conference. Last but not the least, to all members of International Advisory board, Scientific committee and Steering Committee who helped us with their expertise. We would also like to thank the invited speakers Prof. Dr. B. (Bob) Wilffert, Prof. Patrick Ball, Dr. Hana Morrissey, Dr. Arthorn Riewpaiboon, Dr. Nining Sugihartini, M.Si., Apt. and dr. Rizki Edmi Edison, PhD. for accepting our invitation to share their expert knowledge. Finally, thanks to all the committees that have worked hard for the past year to organize MICH-PhD. May the blessings of Allah SWT be upon all of us. We hope that this collaboration can continue and hopefully we can meet again at the second MICH-PhD. **(Less)**

**Volumes:**

Vol. 1 - 978-989-758-349-0

Papers

Authors

Show   papers

**The Impact of Short Message Service (SMS) Reminder and Home Monitoring on Blood Pressure Control in Hypertension Patients: A Case Study of Two Primary Health Care Facilities in Banyumas Regency**

P. 5 - 10

Laksmi Maharani , Hening Pratiwi and Ika Mustikaningtias

**DOI:**10.5220/0008238400050010

**Effect of Acute Administration of Eel (*Anguilla bicolor bicolor*) Oil to Hematological Parameters in Mice** P. 11 - 14

Heru Sasongko

DOI:10.5220/0008238500110014

---

**The Preservation of Liposomes During Air Drying Using a Matrix Containing Maltodextrin and HPMC** P. 15 - 18

Raditya Weka Nugraheni , Helmy Yusuf and Dwi Setyawan

DOI:10.5220/0008238600150018

---

**Molecular Docking Study of Lemon (*Citrus limon* (Linn) Burm. f) Flavonoid Derivatives Compound in Receptor Cyclooxygenase-1 (COX-1) as Antiplatelet in Ischaemic Stroke Disease** P. 19 - 25

Rizky Arcinthya Rachmania , Hariyanti and Nurul Rochmah

DOI:10.5220/0008238700190025

---

**Post-therapeutic Response Evaluation of Patients Receiving Percutaneous Coronary Intervention at the Regional Hospital in Bantul, Yogyakarta** P. 26 - 32

Pramitha Esha Nirmala Dewi

DOI:10.5220/0008238800260032

---

**Evaluation of CA 125, BUN, and Creatinine Serum in Ovarian Cancer Patients Receiving Paclitaxel-Cisplatin Chemotherapy Treatment** P. 33 - 38

Rini Noviyani , P. A. Indrayathi , I. N. G. Budiana , Rasmaya Niruri , K. Tunas and Tamara

Candra Paramitha

DOI:10.5220/0008238900330038

---

**The Effect of Increased Level of Avocado (*Persea americana* Mill.) Seed Starch as Binding Agent on Physical Properties of the Liquorice Extract (*Glycyrrhiza glabra* Linn.) Lozenges** P. 39 - 43

Landyyun Rahmawan Sjahid , Anneke Lionie Kumala Satki and Inding Gusmayadi

DOI:10.5220/0008239000390043

---

**The Antioxidant Activity Analysis of the Ethanolic Extract of Banana Peel (*Musa paradisiaca* forma typica) with DPPH Method** P. 44 - 47

Novia Ariani and Laela Hayu Nurani

DOI:10.5220/0008239100440047

---

**Empirical Antibiotics Study on Pneumonia in Intensive Care Unit** P. 48 - 53

Yeni Farida , Katarina Puspita and Zahra Yusvida

DOI:10.5220/0008239200480053

---

**Formulation of Sechium edule Extract Effervescent Granule with the Variation of Citric Acid, Tartrate Acid and Sodium Bicarbonate** P. 54 - 60

Ina Ba'dia Grajang and Iis Wahyuningsih

DOI:10.5220/0008239300540060

---

**Anti-Hyperlipidemic Effect of 70% Ethanol Extract from Mesona palustris Blume Leaves on Male Hamsters** P. 61 - 65

Dwitiyanti , Ni Putu Ermi Hikmawanti , Kriana Efendi , Riana Puspa Dewi and Fernita Afriyani

DOI:10.5220/0008239400610065

---

**Hepatoprotective Effect of Mountain Papaya (Vasconcellea pubescens A.DC.) Fruit Extract against Acetaminophen-Induced Acute Liver Damage** P. 66 - 70

Heru Sasongko , Diah Pratiwi , Trias Amartawi , Nur Rohman Efendi and Sugiyarto

DOI:10.5220/0008239500660070

---

**Public Counseling: An Educational Model to Improve Medication Adherence in Type 2 Diabetes Mellitus Patients** P. 71 - 76

Daniek Vivianhari , Nora Wulandari and Francyska Putri Puspita

DOI:10.5220/0008239600710076

---

**The Effects of Liposomale Methylprednisolone Palmitate on the Production of TNF $\alpha$  in Mice** P. 77 - 83

Aprilita Rina Yanti Eff , F. D. Suyatna and Erni H. Purwaningsih

DOI:10.5220/0008239700770083

---

**Potential Risk Factors of End Stage Renal Disease in Patients of Hemodialysis** P. 84 - 89

Diana Laila Ramatillah , Syed Azhar Syed Sulaiman , Amer Hayat Khan and Ihsanil Husna

DOI:10.5220/0008239800840089

---

**Analysis of Eugenol Content in Ethanolic Extract of Galangal Rhizome (Alpinia galanga L. Willd) Ointment Using UV-VIS Spectrophotometry Method** P. 90 - 95

Adi Yugatama , Sholichah Rohmani and Rizky Apriliani

DOI:10.5220/0008239900900095

---

**Steamed Watermelon (Citrullus lanatus Thunb.) Juice Improves Spatial Memory in Dementia Rat Model** P. 96 - 101

Maifitrianti , Hadi Sunaryo and Dedi Suryadi

DOI:10.5220/000824000960101

---

**Anti-hyperuricemia Effect of Water Fraction Cinnamon (*Cinnamomum burmannii* (Ness & T. Ness) Blume) on White Male Rats** P. 102 - 106

Dwitiyanti , Ema Dewanti and Rizky Arcintha Rachmania

DOI:10.5220/0008240101020106

---

**The Alpha-Amylase Inhibition Potential of Endophytic Fungi from Indonesian Bay Leaves (*Eugenia polyantha* WIGHT.)** P. 107 - 111

Wahyu Hidayati , Ade Nur Padillah , Maharadingga , Ni Putu Ermi Hikmawanti , Rini Prastiwi , Ani Pratiwi , Lady Farahmayuni , Rezza Syahputra and Muhammad Fahrul

DOI:10.5220/0008240201070111

---

**Identification of Potentially Inappropriate Prescribing in Outpatient Geriatric using STOPP/START Criteria at X Hospital Jakarta** P. 112 - 116

Numlil Khaira Rusdi , Dini Indah Komariah , Nora Wulandari and Arya Govinda Roosheroe

DOI:10.5220/0008240301120116

---

**Study in Activity Combination of *Physalis angulata* and *Hibiscus sabdariffa* in 70% Ethanol Extract to Decrease Blood Sugar Levels and Histopathology of Pancreas Langerhans Island in Alloxan Induced Diabetic Rats** P. 117 - 122

Hadi Sunaryo , Ni Putu Ermi Hikmawanti and Hesty Awanis Listyaningrum

DOI:10.5220/0008240401170122

---

**In Silico Analysis of the Phytochemical Compounds in *Carica papaya* Seeds for Optimizing the Inhibitors of HMG-CoA Reductase** P. 123 - 132

Hariyanti , Rizky Arcintha Rachmania , Mutia Karinah and Hadi Sunaryo

DOI:10.5220/0008240501230132

---

**Antihyperglycemic Activity of Ethanolic Herb Extract of Ceplukan (*Physalis angulata* L.) in Diabetic Hypercholesterolemia in Male Hamsters** P. 133 - 137

Elly Wardani , Dwitiyanti , Sediarto and Dwina Puspandiyah

DOI:10.5220/0008240601330137

---

**Extraction, Identification, and Gel Formulation of Mangiferin from Mango (*Mangifera indica* L.) Leaves Extract** P. 138 - 142



Rudi Afrinanda , Yusa Ristiawati , Muhammad Shoufi Islami and Deasy Vanda Pertiwi

DOI:10.5220/0008240701380142

---

**The Potency of Binahong Leaves (*Anredera cordifolia* (Ten.) Steenis) Subfraction with Ethanol 70% as an Antihyperuricemic Agent** P. 143 - 146

Vera Ladeska , Ani Pahriyani and Monika Silviani Gunawijaya

DOI:10.5220/0008240801430146

---

**The Effect of Concentration Ratio of Gelatine and Polyvinylpyrrolidone as Binders on the Physical Properties of Red Ginger (*Zingiber officinale* Rosc.) Extract Lozenges** P. 147 - 153

Inding Gusmayadi and Priyanto

DOI:10.5220/0008240901470153

---

**Formulation of Moringa oleifera Leaf Extract in Lotion and Gel as Sunscreen** P. 154 - 158

Nining Sugihartini , M. Alif Fajri and Desty Restia Rahmawati

DOI:10.5220/0008241001540158

---

**Quality Control of Turmeric Rhizome (*Curcuma domestica* Val) as Traditional Medicine from Wonogiri, Central Java** P. 159 - 168

Fatimah Nisma , Ema Dewanti , Rini Prastiwi , Alexander , Wanda Puspita Sari and Wido Artanto

DOI:10.5220/0008241101590168

---

**Screening of Antibacterial Potency and Molecular Identification of Endophytic Bacteria from Soursop Leaf (*Annona muricata* L.)** P. 169 - 175

Fitri Yuniarti , Wahyu Hidayati and Lulu Shofaya

DOI:10.5220/0008241201690175

---

Papers  
Authors  
Ontology

Science and Technology Publications, Lda  
Avenida de S. Francisco Xavier, Lote 7 Cv. C,  
2900-616 Setúbal, Portugal.

Phone: +351 265 520 185  
Fax: +351 265 520 186  
Email: [info@scitepress.org](mailto:info@scitepress.org)

INSTICC  
SCITEVENTS  
CROSSREF

dblp  
Ei Compendex  
SCOPUS  
Semantic Scholar  
Google Scholar  
Microsoft Academic

## Steamed Watermelon (*Citrullus lanatus* Thunb.) Juice Improves Spatial Memory in Dementia Rat Model

In Proceedings of the 1st Muhammadiyah International Conference on Health and Pharmaceutical Development - MICH-PhD, 96-101, 2018 , East Jakarta, Indonesia

### Steamed Watermelon (*Citrullus lanatus* Thunb.) Juice Improves Spatial Memory in Dementia Rat Model

Maifitrianti, Hadi Sunaryo and Dedi Suryadi

Faculty of Pharmacy and Science, Universitas Muhammadiyah Prof. DR. HAMKA, Jakarta, Indonesia

**Keywords:** Watermelon, Lycopene, Dementia, Spatial Memory

**Abstract:** Watermelon (*Citrullus lanatus* Thunb.) is one of the fruits that containing lycopene. Lycopene is able to reduce oxidative damage that causes dementia. This research was aimed to determine the activity of steamed watermelon juice in improving the spatial memory in dementia rats model induced by Electroconvulsive Shock (ECS). The animal test was divided into 6 groups, namely normal control group, negative control group, positive control group (vitamin E, 20.7 mg/kg BW), and 3 treatment groups that received steamed watermelon juice with different dose, i.e. 0.92 g/kg BW, 1.85 g/kg BW, and 3.7 g/kg BW. All groups were induced with ECS, except for the normal control group. The spatial memory was measured with the Radial Arm Maze (RAM) test and the observed results were analyzed with ANOVA and a post-hoc Tukey's HSD test. The ANOVA test results showed significant differences between all treatment groups. Meanwhile, the post-hoc Tukey's HSD test at 95% confidence level resulted in a significant difference between the treatment groups with the doses of 0.92 g/kg BW and 1.85 g/kg BW and the negative control group. However, this test indicated an insignificant difference between the positive control group and the normal control group. This finding confirms that the administration of steamed watermelon juice at doses of 0.92 g/kg BW and 1.85 g/kg BW improves the spatial memory in a rat model of dementia comparable to the positive control. The treatment groups receiving these two doses showed corresponding results, meaning that both doses of steamed watermelon juice have an equivalent spatial memory enhancement activity.

SCIENCE AND TECHNOLOGY PUBLICATIONS

#### 1 INTRODUCTION

Dementia is a syndrome of decreased intellectual function that is severe enough to interfere with social and economic activities in daily life, and is, usually followed by behavioral changes not attributable to delirium or major psychiatric disorders. The prevalence of dementia increases rapidly with life expectancy. Even though there has been no national research data regarding the prevalence of dementia the growing number of elderly population predictably increases the incidence of dementia in Indonesia. The Stroke Registry Indonesia 2013 support this assumption. It reports that stroke become more prevalent at a younger age, and 60.59% of stroke patients experience impaired cognition after returning home from hospital treatment. This registry also explains that the stroke risk factors in the country are fairly high (Perhimpunan Dokter Spesialis Saraf Indonesia, 2015).

Oxidative stress is significantly involved in the pathology of dementia even at the early stage of the disease (Mao, 2013). The brain and nerve cells are susceptible to damages by free radicals because the membrane lipids contain many side chains of polyunsaturated fatty acids. Oxidative stress also depends on other factors, namely the quantity of brain oxygen consumption, iron and other metals transition that responsible for the formation of Reactive Oxygen Species (ROS), and antioxidants in the brain. Antioxidants have been widely studied in relation to neurodegenerative diseases (Rao *et al.*, 2003). Research in Parkinson's patients and vascular dementia indicates low serum lycopene levels (Fay, *et al.*, 1999), which, according to a stroke prevention study in Austria, are associated with a high risk of microangiopathy (Schmidt *et al.*, 1997). In elderly patients, a high level of lycopene is positively connected to their functional capacity, as indicated by their ability to perform self-care (Snowdon *et al.*, 1996). Given the importance of

96

Maifitrianti, Sunaryo, H. and Suryadi, D.

Steamed Watermelon (*Citrullus lanatus* Thunb.) Juice Improves Spatial Memory in Dementia Rat Model.

DOI: 10.30605/STP.2018.96.96-101

In Proceedings of the 1st Muhammadiyah International Conference on Health and Pharmaceutical Development (MICH-PhD 2018), pages 96-101

ISBN: 978-602-750-048-0

Copyright © 2018 by SCITEPRESS - Science and Technology Publications, Ltd. All rights reserved.

**Affiliation:** Universitas Muhammadiyah , Indonesia

**Keyword(s):** Watermelon, Lycopene, Dementia, Spatial Memory

**Abstract:** Watermelon (*Citrullus lanatus* Thunb.) is one of the fruits that containing lycopene. Lycopene is able to reduce oxidative damage that causes dementia. This research was aimed to determine the activity of steamed watermelon juice in improving the spatial memory in dementia rats model induced by Electroconvulsive Shock (ECS). The animal test was divided into 6 groups, namely normal control group, negative control group, positive control group (vitamin E, 20.7 mg/kg BW), and 3 treatment groups that received steamed watermelon juice with different dose, i.e 0.92 g/kg BW, 1.85 g/kg BW, and 3.7 g/kg BW. All groups were induced with ECS, except for the normal control group. The spatial memory was measured with the Radial Arm Maze (RAM) test and the observed results were analyzed with ANOVA and a post-hoc Tukey's HSD test. The ANOVA test results showed significant differences between all treatment groups. Meanwhile, the posthoc Tukey's HSD test at 95% confidence level resulted in a significant (More)



CC BY-NC-ND 4.0



Guest: Register as new SciTePress user now for free.



SciTePress user: please login.

You are not signed in, therefore limits apply to your IP address  
139.193.140.151



My Papers

In the current month:

Recent papers: 100 available of 100 total

2+ years older papers: 200 available of 200 total

### Paper citation in several formats:

Harvard

Bibtex

EndNote

Maifitrianti.; Sunaryo, H. and Suryadi, D. (2019). **Steamed Watermelon (*Citrullus lanatus* Thunb.) Juice Improves Spatial Memory in Dementia Rat Model.** In *Proceedings of the 1st Muhammadiyah International Conference on Health and Pharmaceutical Development - MICH-PhD*, ISBN 978-989-758-349-0, pages 96-101. DOI: 10.5220/0008240000960101

Authors	Science and Technology Publications, Lda	SCITEVENTS	dblp
Ontology	Avenida de S. Francisco Xavier, Lote 7 Cv. C, 2900-616 Setúbal, Portugal.	CROSSREF	Ei Compendex
	Phone: +351 265 520 185		SCOPUS
	Fax: +351 265 520 186		Semantic Scholar
	Email: info@scitepress.org		Google Scholar
			Microsoft Academic

# Steamed Watermelon (*Citrullus lanatus* Thunb.) Juice Improves Spatial Memory in Dementia Rat Model

Maifitrianti, Hadi Sunaryo and Dedi Suryadi

Faculty of Pharmacy and Science, Universitas Muhammadiyah Prof. DR. HAMKA, Jakarta, Indonesia

**Keywords:** Watermelon, Lycopene, Dementia, Spatial Memory

**Abstract:** Watermelon (*Citrullus lanatus* Thunb.) is one of the fruits that containing lycopene. Lycopene is able to reduce oxidative damage that causes dementia. This research was aimed to determine the activity of steamed watermelon juice in improving the spatial memory in dementia rats model induced by Electroconvulsive Shock (ECS). The animal test was divided into 6 groups, namely normal control group, negative control group, positive control group (vitamin E, 20.7 mg/kg BW), and 3 treatment groups that received steamed watermelon juice with different dose, i.e 0.92 g/kg BW, 1.85 g/kg BW, and 3.7 g/kg BW. All groups were induced with ECS, except for the normal control group. The spatial memory was measured with the Radial Arm Maze (RAM) test and the observed results were analyzed with ANOVA and a post-hoc Tukey's HSD test. The ANOVA test results showed significant differences between all treatment groups. Meanwhile, the post-hoc Tukey's HSD test at 95% confidence level resulted in a significant difference between the the treatment groups with the doses of 0.92 g/kg BW and 1.85 g/kg BW and the negative control group. However, this test indicated an insignificant difference between the positive control group and the normal control group. This finding confirms that the administration of steamed watermelon juice at doses of 0.92 g/kg BW and 1.85 g/kg BW improves the spatial memory in a rat mode of dementia comparable to the positive control. The treatment groups receiving these two doses showed corresponding results, meaning that both doses of steamed watermelon juice have an equivalent spatial memory enhancement activity.

## 1 INTRODUCTION

Dementia is a syndrome of decreased intellectual function that is severe enough to interfere with social and economic activities in daily life, and is, usually followed by behavioral changes not attributable to delirium or major psychiatric disorders. The prevalence of dementia increases rapidly with life expectancy. Even though there has been no national research data regarding the prevalence of dementia the growing number of elderly population predictably increases the incidence of dementia in Indonesia. The Stroke Registry Indonesia 2013 support this assumption. It reports that stroke become more prevalent at a younger age, and 60.59% of stroke patients experience impaired cognition after returning home from hospital treatment. This registry also explains that the stroke risk factors in the country are fairly high (Perhimpunan Dokter Spesialis Saraf Indonesia, 2015).

Oxidative stress is significantly involved in the pathology of dementia even at the early stage of the disease (Mao, 2013). The brain and nerve cells are susceptible to damages by free radicals because the membrane lipids contain many side chains of polyunsaturated fatty acids. Oxidative stress also depends on other factors, namely the quantity of brain oxygen consumption, iron and other metals transition that responsible for the formation of Reactive Oxygen Species (ROS), and antioxidants in the brain. Antioxidants have been widely studied in relation to neurogenerative diseases (Rao *et al.*, 2003. Research in Parkinson's patients and vascular dementia indicates low serum lycopene levels (Foy, *et al.*, 1999), which, according to a stroke prevention study in Austria, are associated with a high risk of microangiopathy (Schmidt *et al.*, 1997). In elderly patients, a high level of lycopene is positively connected to their functional capacity, as indicated by their ability to perform self-care (Snowdon *et al.*, 1996). Given the importance of

attention to neurogenerative diseases, studies related to the effectiveness of lycopene in the management of dementia are necessary.

Lycopene is fat-soluble antioxidant synthesized by many plants and microorganisms. It is responsible for giving the red color in various vegetables and fruits. Earlier, only tomato and its products were considered as the potential sources of lycopene, but nowadays there are scientifically proven facts that watermelon also contains an appreciable amount of *cis*-configured lycopene. Therefore, people gradually shift their preference towards watermelon and its allied products to deal with their health concerns. Nevertheless, the quantity of lycopene varies depending upon the variety and growing conditions (Fish and Davis, 2003). Overall, in the cells, lycopene is present in the form of crystalline with a range of 2.30-7.20 mg/100 g fresh weight basis (Huh *et al.*, 2008; Artes-Henandez *et al.*, 2010). In general, the human body only absorbs 10-30% of fresh lycopene in fresh lycopene in fruits or vegetables. Lycopene is more effectively absorbed when consumed from processed foods, such as sauce and juice. The increased absorption of lycopene in tomato products is due to the presence of *cis*-isomers of lycopene. Lycopene naturally appears as 'trans', which is thermodynamically stable, soluble in non-polar solvents, and identifiable at a wavelength of 446-506 nm. Lighting and heating can potentially change the isomer form 'trans' to 'cis' (Rao & Rao, 2007). Maulida & Zulkarnaen (2010) conclude that the optimum temperature for watermelon to release lycopene from its cell structure is 70°C.

This research aimed to determine the activity of steamed watermelon juice in improving the spatial memory in a rat model of Electroconvulsive Shock (ECS)-induced dementia and to determine the effective dose to increase spatial memory. Steamed watermelon juice was expected to be able to improve spatial memory and function as a natural treatment, which would be potentially developed into phytopharmaceutical preparations, as well as a reference for further research to support the development of science.

## 2. MATERIALS AND METHODS

### 2.1 Materials

The research apparatus included an electroconvulsive meter Model EC 02 from Orchid Scientific and Innovative India Pvt. Ltd. (B 59, MIDC, Ambad,

Nashik-422010 India) and a radial arm maze designed with eight equidistantly spaced arm (53.5 cm x 12.5 cm) extending from a 24 cm diameter central platform elevated 11.5 of the floor.

The fresh watermelons (*Citrullus lanatus* Thunb.) were obtained from Cipanas, West Java and determined at LIPI Biological Research Center, Bogor from the Herbarium Bogoriense collection, Vitamin E (*tocopherol*) was acquired from PT. Cortico Mulia Sejahtera. The test animals were white male rats (*Rattus norvegicus*) strain *Sprague Dawley* aged 2-3 months and weighed  $\pm$  200 gram.

## 2.2 Methods

### 2.2.1 Identification of Watermelon

The watermelons were taxonomically identified from the Herbarium Bogoriense collection at LIPI Biological Research Center, Bogor.

### 2.2.2 Preparation of Steamed Watermelon Juice

Around 100 grams of watermelons were steamed for 5 minutes at a temperature of 70°C and the temperature was measured with a thermometer (Maulida & Zulkarnaen, 2010). The steamed watermelon was drained and the juice was extracted using a juicer machine.

### 2.2.3 Phytochemical Screening

The steamed watermelon juice was tested for the presence of alkaloids, flavonoids, steroids, tannins, triterpenoids, and saponins.

### 2.2.4 Treatment of Test Animal

This research is approved by the Ethics Committee of Health Research, Faculty of Medicine, University of Indonesia No. 278/UN2. F1/ETHICS/2017.

A total of 30 white male *Sprague Dawley* rats (*Rattus norvegicus*) aged 2-3 months old and weighed with  $\pm$  200 g was divided into six (6) groups, each consisting of five (5) rats. These six groups were: Group 1 as normal control, Group 2 as negative control, Group 3 as positive control, Group 4 as a treatment group receiving 0.92 g/kg BW of steamed watermelon juice, Group 5 as a treatment group receiving 1.85 g/kg BW of steamed watermelon juice, and Group 6 as a treatment group receiving 3.7 g/kg BW of steamed watermelon juice.

Table 1: The phytochemical screening results of steamed watermelon juice.

Compounds	Results
Alkaloids	+
Flavonoids	-
Tannins	-
Steroids and Triterpenoids	+
Saponins	+

The animals were acclimatized for at least five (5) days to the laboratory conditions before the behavioral experiments. They were placed in a natural light/dark cycle (12 h each) and given free access to standard food and water. From the second to the fourth day of the acclimatization, they were trained in the maze environment. After all group of rats fasted for 12 hours, they were subjected to preliminary radial arm maze test for 12 days. This test was called the Radial Arm Maze Test I (RAM I). The rats were placed on the central platform and the food (pellets) were hidden in a container at the end of the arm. The rat performance represented the ratio of the number of reward eaten pellets to the total number of arms entered in 10 minutes. There were a total of eight (8) arms in the maze.

After the RAM I was completed, Groups 2, 3, 4, 5, and 6 received ECS with a magnitude of 112 mA/150 V for 1 sec/day. Afterward, the experimental animals were rested for 10 minutes. Then, they were placed in the maze again and re-tested to prove that the ECS induced a dementia model. The ECS was performed for eight (8) days.

The positive control group (Group 3) received Vitamin E at a dose of 20.7 mg/kg BW, Groups 4, 5, and 6 were given steamed watermelon juice at different doses, namely 0.92 g/kg BW, 1.85 g/kg BW and 3.7 g/kg BW, respectively. The steamed watermelon juice and vitamin E were administered orally once a day every morning for 14 days.

The next stage was a post-test, i.e., a radial arm maze test for 12 consecutive days, referred to as RAM II. Before the post-test the rats fasted for 12 hours. The performance of the test rats in the RAM assay was calculated with the formula (1) (Sari & Ar Rochman, 2015):

$$\text{Performance (\%)} = \frac{(\text{Number of eaten reward (food)})}{(\text{Number of arms entered by the rats})} \times 100\% \quad (1)$$

Table 2: The observation and assessment results of the performance (%) of the experimental rat groups in RAM I.

Day	The Performance of the Groups (in %)					
	1	2	3	4	5	6
1	23.3	5	3.3	20	0	10.6
2	0	26.6	23.3	32.5	9.8	5.3
3	68	23.1	57.6	58.6	43.3	76
4	20	53.8	28.3	37.5	34.4	39.2
5	25	73.8	74.2	68	61.1	100
6	20	64.2	35	63.6	73	37.5
7	7.3	46.2	74.2	60	55	57.5
8	20	56.6	40	46	65.8	57.1
9	20	56.6	40	46	65.8	53.1
10	79.6	33.3	89.2	36.6	100	68
11	100	97.1	100	42	87.1	76.6
12	100	80	96	40	90	80

### 2.2.5 Data Analysis

The mean percentage of the performance of the experimental groups in RAM II was tested for normality using the Shapiro-Wilk test and homogeneity using the Levene's test. Afterward, the data were analyzed with one way ANOVA at 95 % significance level ( $\alpha = 0.05$ ). If the results showed a statistically significant difference, then the data analysis was continued with the Tukey test (Priyatno, 2012).

## 3 RESULTS AND DISCUSSION

The phytochemical screening showed that steamed watermelon juice was positive for alkaloids, terpenoids, steroids and saponins (Table 1).

Before the induction of dementia with ECS, all groups of experimental animals were subjected to the radial arm maze test I; henceforth referred to as RAM I, this test identified the basic memory of each rat and the whole six groups, and affirmed that all of them had the same condition before the treatment. The results of RAM I are described in Table 2. The entire groups showed remarkable spatial memory every day during the 12 days observation. Spatial learning and memory help animals find locations that provide, among other things, food and safety, and are thereby crucial for survival (Dogru *et al.*, 2003).

The performance of each test group in selecting the arms and feeding in RAM I showed an increase every day. Even though Group 1 had lowest average score



Table 3: The performance (%) of the experimental rat groups in the RAM test after the induction of ECS.

Day	The Performance of the Groups (in%)					
	1	2	3	4	5	6
1	50.3	24	0	0	59.3	20
2	54	40	20	0	30	55
3	60.3	37.3	0	0	46.3	53.3
4	81.5	52	0	0	20	18.3
5	94	14	16.6	10	6.6	40
6	85.6	20	6.6	0	26.6	0
7	87	80	12	0	50	10
8	86.6	55	35	0	40.6	20

of performance among all the test groups (38.3%), it scored 100% on Day 11 and 12. Meanwhile, Group 5 had the highest average (57.1%), followed by Group 3 and 6 (55%), Group 2 (51.4%), and Group 4 (45.8%).

After the 12-day observation in RAM I, the animal model of dementia was prepared. Except for the normal control group, all of the experimental groups were given ECS with an intensity of 150 V/112mA for 1 sec/day for eight (8) days. Each rat was electrocuted at the top of the ear and jammed until the electricity caused muscle spasms and stiffness. After rested for 10 minutes, the rats were tested in the maze to prove that the group had already developed dementia. The observation results are summarized in Table 3.

The results of the RAM assays after the ECS showed a decrease in the percentage of feeding and arm entries. When compared with normal controls (no ECS), all of the treatment groups showed low performances. Groups 4 even failed seven times (0%) and it was the lowest performance of all test groups. These results are the effect of ECS-induced memory degradation. They indicate poor working and reference memory capabilities, as well as low learning abilities in rats. During the maze test, the rats behaved as if in confusion and remained silent in the maze. They moved slowly and did not recognize the food placed at the end of the arm. The ECS decreased their memory because it involved the formation of free radicals that attacked the walls and nerve cells of their brains. These cells are composed of unsaturated fatty acids. Therefore, ECS would lead to peroxidation of fatty acids and neurotoxicity, causing neurotransmitter death and, subsequently a decline in memory (Luo *et al.*, 2011).

Groups 4, 5, and 6 received steamed watermelon juice at different doses namely 0.92 g/kg BW, 1.85 g/kg BW, and 3.7 g/kg BW respectively, for 14 days.

Table 4: The Performance (%) of the Experimental Rat Groups in RAM II.

Day	The Performance of the Groups (%)					
	1	2	3	4	5	6
1	39.3	13.3	22.1	18.7	0	0
2	40	0	29.1	50	37.5	0
3	51.6	13.3	50	62.5	33.5	27
4	94.4	21	87.5	100	83.3	46.6
5	66.6	10	45.8	55.3	100	66.6
6	50	20	26.1	80	72.2	56.2
7	66.6	20	66.6	87.5	91.6	75
8	66.6	13.2	61.1	83.3	44.4	66.6
9	50	20	91.6	100	66.6	66.6
10	66.6	20	100	100	88.8	33.3
11	66.6	20	100	83.3	100	33.3
12	66.6	0	100	75	100	33.3

The steamed watermelon juice was freshly made each day. The watermelon was prepared in small pieces, weighed using an analytical scale according to the needs of the three (3) treatment groups steamed at 70°C for 5 minutes and rested until the temperature decreased. Afterward, the steamed watermelon was drained and ready for juice extraction. The positive control received Vitamin E at a dose of 20.7 mg/kg BW dissolved in Na CMC. After these treatments, the rats were subjected to the second performance test, i.e., RAM II to identify any increases in spatial memory. The observation results are presented in Table 4.

Table 4 shows an increase in spatial memory during the 12 day observation in RAM II. The spatial memory of the treatment groups began to increase on Day 4 and reached its peak on the last day of the observation. Group 4 showed the highest performance with an average of 74.6%. The spatial memories of Groups 3, 5, and 6 increased by 65%, 68.1% and 42% respectively. The normal control (Group 1) had a different performance with negative control (Group 2). Furthermore, the performance of the negative control in RAM II was significantly different from its performance in RAM I. This group showed the lowest performance demonstrating the poor working and reference memory capabilities and the low learning ability of rats.

Group 4 and 5 each receiving steamed watermelon juice at the doses of 0.92 g/kg BW and 1.85 g/kg BW, respectively, scored 100% three times and ranked as the highest performance groups. This result suggest that these doses are effective to increase the spatial

memory of rats with ECS induced dementia. The memory improvement may be caused by an increase in the synaptic strength between the neurons in rat's hippocampus or referred to as neuronal plasticity, especially in the *gyrus dentatus* and CA1 hippocampus that are responsible for spatial information (Poirer *et al.*, 2008).

Interestingly, unlike the rest of the treatment groups, Group 6 that received steamed watermelon juice at a dose of 3.7 g/kg BW showed a decreased performance, even lower than normal control by 42%. This group failed two times (i.e., 0% performance) on the first and second day of observation. Its second lowest constant rate (33.3%) occurred on Day 10, or one day after the increased performance of the negative control group, until the end of the observation. The poor working memory may be behind this result. This memory is responsible only for the information received from one session of RAM test. It is erased shortly before the next session of RAM test, and therefore the rats try to habituate themselves to the maze environment again and form a new working memory, while making mistake (i.e., incorrect arm entries). According to Crusio & Schwegler (2005), the working memory is closely related to spatial memory. The RAM test also observed reference memory that, for example, encoded and retained information about the food at the end of some arms. This memory is always useful in any maze test because it has already faded before 24 hours. In this case, the rats must form new memory again from scratch allowing for a 24-hour maze test (Rao *et al.*, 2003).

The average percentage of the group performance in RAM II was subjected to homogeneity and normality tests. The normality test yielded  $p$ -value 0.142 ( $p > 0.05$ ), while the homogeneity test resulted in  $p$ -value 0.1 ( $p > 0.05$ ). The data analysis was continued with a one-way ANOVA to determine any statistically significant difference between the treatment groups in completing the maze test after the induction of ECS. The ANOVA test produced  $p$ -value 0.000 ( $p < 0.05$ ), representing a significant difference between the groups in RAM II.

The subsequent Tukey test proved that in the positive control group (Group 3), as well as the treatment groups receiving 0.92 g/kg BW (Group 4) and 1.85 g/kg BW (Group 5) of steamed watermelon juice, the spatial memory in the rat model of dementia improved. These three groups had a similar spatial memory improvement to the normal control group. The two treatment groups i.e., Group 4 and 5 showed comparable results as well, meaning, that the groups receiving 0.92 g/kg BW (Group 4) and 1.85 g/kg BW

(Group 5) experience an equivalent spatial memory enhancement activity. Meanwhile, receiving steamed watermelon juice at a dose of 3.7 g/kg BW, Group 6 exhibited significantly different activities with the normal and positive control groups, but it was comparable to the negative control. In other words, the treatment in Group 6 was not effective in increasing the spatial memory of the experimental animals. The effective doses of steamed watermelon juice in this study were 0.92 g/kg BW and 1.85 g/kg BW. The average percentage of spatial memory improvement in these doses exceeded the positive control group (Vitamin E, 20.7 mg/kg BW).

Theoretical oxidative damage is preventable if free radicals and antioxidants in the body are in balance. In this condition antioxidants will not exhibit any effects (Asri, 2014). The antioxidative properties of lycopene contained in steamed watermelon juice are allegedly one of the factors that prevent ECS-induced oxidative damage. Lycopene is a highly unsaturated straight-chain hydrocarbon with a total of 13 double bonds, eleven (11) of which are conjugated. This unique nature makes lycopene molecule a very potent antioxidant. An in vitro study affirms that lycopene is twice as potent as  $\beta$ -carotene and its singlet oxygen quenching ability is ten times greater than a tocopherol (Rao *et al.*, 2003).

## 4 CONCLUSIONS

This study concludes that the administration of steamed watermelon juice at the doses of 0.92 g/kg BW and 1.85 g/kg BW can improve the spatial memory in a rat model of dementia, which is comparable to the positive control. Treatment groups that received these doses had corresponding spatial memory enhancement activities.

## REFERENCES

- Artés-Hernández F., Robles P. A., Gómez P. A., Tomás-Callejas A., & Artés F., 2010. Low UV-C illumination for keeping overall quality of fresh-cut watermelon. *Postharvest Biology and Technology-Journal*. 55:114–120.
- Asri, W., 2014, Peran Antioksidan Bagi Kesehatan, *Jurnal Biotek Medisiana Indonesia*, Vol. 3. 2. 2014: 59-68
- Crusio, E., W., & Schwegler, H., 2005. Learning Spatial Orientation Tasks in The Radial Maze and Structural Variation in The Hippocampus in Inbred Mice. *Behavior and Brain Function* 1:3; DOI: 10.1186/1744-9081,1-3

- Dogru, E., J., Gumusbas, U., & Kara, F., 2003, Individual Variation in the Spatial Reference and Working Memory Assessed under Allothetic and Idiothetic Orientation Cues in Rat, *Acta Neurobiologiae Experimental*, 63: 17-23
- Fish W. W., & Davis A. R., 2003. The effects of frozen storage conditions on lycopene stability in watermelon tissue. *Journal of Agricultural and Food Chemistry*.51:3582–3585.
- Foy C. J., Passmore A. P., Vahidassr M. D., Young I. S., & Lawson J. T., 1999, Plasma chain-breaking antioxidants in Alzheimer's disease, vascular dementia and Parkinson's disease. *QJM: An International Journal of Medicine*, 92: 39-45
- Huh Y. C., Solmaz I., & Sari N., 2008. Morphological characterization of Korean and Turkish watermelon germplasm. In: Pitrat M, editor. *Cucurbitaceae. Proceedings of the IXth EUCARPIA meeting on genetics and breeding of Cucurbitaceae, Avignon (France)*, May 21-24th. 2008. pp. 327–333
- Luo, J. S., Min, K., Wei, P., Li, J., Dong, J., & Liu, Y.F. 2011, Propofol Protects Against Impairment of Learning-memory and Imbalance of Hippocampal Glu/GABA Induced by Electroconvulsive Shock in Depressed Rats, *Journal of Anesthesia*, Vol 25: 657– 665.
- Maulida, D., & Zulkarnaen, N., 2010, Ekstraksi Antioksidan (Likopen) dari Buah Tomat dengan Menggunakan Solven Campuran, n-Heksana, Aseton, dan Etanol, *Skripsi*, Semarang: Jurusan Teknik Kimia, Fakultas Teknik, Universitas Diponegoro.
- Mao, P., 2013. Oxidative Stress and Its Clinical Application in Dementia. *Journal of Neurogenerative Disease* Vol. 2013, Article ID 319898, 15 pages
- Perhimpunan Dokter Spesialis Saraf Indonesia, 2015, *Panduan Paktik Klinik Diagnosis dan Penatalaksanaan Demensia*, Perhimpunan Dokter Spesialis Saraf Indonesia: Jakarta.
- Poirier, G. L., Amin, E., & John P., 2008, Qualitatively Different Hippocampal Subfield Engagement Emerges with Mastery of a Spatial Memory Task by Rats, *The Journal of Neuroscience*. 28(5):1034 –1045
- Priyatno, D, Belajar Praktis Analisis Parametrik dan Nonparametrik Dengan SPSS & Prediksi Pertanyaan Pendadaran Skripsi dan Tesis. Gava Media: Yogyakarta
- Rao, L.G., Guns, E., & Rao, A.V., 2003. Lycopene: Its role in Human Health and Disease, *AGROFood industry hi-tech*, July-Agustus 2003
- Rao, A.V., & Rao, L.G., 2007. Carotenoids and Human Health. *Pharmacological Research* 55, 207-216.
- Sari, D. C. R., & Ar Rochmah, M, 2015. The Effects of Extract of *Centella asiatica* Leaf on the Retention of Spatial Memory in Rats (Sprague Dawley) After Chronic Electrical Stress, *Knowledge Life Science, The 3rd International Conference on Biological Science*, pp. 159-167.
- Schmidt, R., Fazekas, F., Hayn, M., Schmidt, H., Kapeller, P., Roob, G., Offenbacher, H., Schumacher, M., Eber, B., Weinrauch, V., Kostner, G. M., & Esterbauer, H., 1997. Risk factors for microangiopathy-related cerebral damage in the Austrian stroke prevention study. *Journal of The Neurological Science*, 6; 152 (1):15-21.
- Snowdon, D. A., Gross, M. D., & Butler, S. M, 1996. Antioxidants and reduced functional capacity in the elderly. *The Journal of Gerontology Series A - Biological Sciences and Medical Sciences*, 51: M10-M16.

With high appreciation presents

# Certificate

to

*Maifitrianti, M.Farm., Apt.*

For her/his attendance as **Poster Presenter**  
in the

**Muhammadiyah International Conference on Health and  
Pharmaceutical Development (MICH-PhD)**

***"Current Innovation in Chronic Diseases Treatment"***

With SKP **IAI** as **Poster Presenter: 3 SKP** number: 263/SK-SKP/PP.IAI/IV/2018



**Dr. Hadi Sunaryo, M.Si., Apt.**  
Dean of Faculty of Pharmacy and Science - UHAMKA



**Nurhannah, M.Farm., Apt.**  
Chairman of Organization Committee

