

PAPER • OPEN ACCESS

Preface

To cite this article: 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **755** 011001

View the [article online](#) for updates and enhancements.

You may also like

- [The teaching and learning of addition and subtraction of integers through manipulative in Brunei Darussalam](#)
N Sahat, K A Tengah and R C I Prahmana
- [Arithmetic learning with the use of graphic organiser](#)
F L Sai, M Shahrill, A Tan et al.
- [Identifying the Factors Affecting Students' Performances in Primary School Mathematics](#)
N A Abdullah, M Shahrill, J Yusof et al.

Annual Conference on Health and Food Science Technology (ACHOST) 2020



1ST Annual Conference on Health and Food Science Technology (ACHOST) 2020

We are glad to introduce you the proceedings of the first Annual Conference on Health and Food Science Technology (ACHOST) 2020. The 1st ACHOST 2020 addresses challenges and innovations the field of Food Science and Health, Environmental Science and Issue, Earth Science and Technology. It also provides a premier interdisciplinary platform for researchers, educators and practitioners to present and discuss the most recent innovations, trends, and concerns as well as practical challenges encountered and solutions adopted in the fields of applied science, technology, and engineering.

As we may aware, the World Health Organization officially declared the novel coronavirus COVID-19 a pandemic. Governments around the world are now issuing restrictions on travel, gatherings, and meetings in an effort to limit and slow the spread of the virus. The health and safety of the author and reseacher community is our first priority and we are supporting these efforts. Therefore, the ACHOST conference was held virtually on 25 November 2020.

The ACHOST conference is hosted by PT. Kresna Acitya Nusantara Mediatama and co-hosted by Universitas Muhammadiyah Sidoarjo, Relawan Jurnal Indonesia. This year, we held this flexible online conference to gather experts and scholars around the globe with the aim to continue disseminating the latest advanced research in the field of Food Science and Health, Environmental Science and Issue, Earth Science and Technology. The conference was held from Yogyakarta as the host of the event. The ACHOST 2020 event is virtually implemented with a model that all invited speakers are given time to present their material for about 30-45 minutes each. It then followed by a question and answer by the participants with a direct questioning system, through chat forums and Q&A forums provided by the zoom application. Overall, the conference took 6 hours.

The number of participants who joined the zoom room was recorded around 341 participants. The authors or participants are came from 5 countries, namely Indonesia, Malaysia, Brunei Darussalam, Phillipine and India. Indonesian Participants are come from 19 Provinces of 33 Provinces.

The committee of the conference are honored to have invited following renowned experts as our keynote speakers. Wahyu Caesarendra, PhD from Universiti Brunei Darussalam; Andri Pramesyanti, Ph.D from Universitas Pembangunan Nasional Veteran Jakarta.

We are glad to share with you that around 155 pre-registered authors are submitted their work in the conferences. However, its about 97 papers are selected and accepted for the conferences. All the papers have been through rigorous review by a panel of reviewers who provide critical comments and corrections, and have contributed substantially to the improvement of the quality of the papers to meet the requirements of International publication standard and IOP EES Scope.

We would like to express our sincere gratitude to the Chairman, the distinguished keynote speakers, as well as all the participants. We also want to thank the publisher for publishing the proceedings. May the readers could enjoy the gain some valuable knowledge from it. We are expecting more and more experts and scholars from all over the world to join this international event next year.

Chair of the Organizing Committee

Dwi Fajar Saputra

Editors

Dwi Fajar Saputra

Mochammad Tanzil Multazam

Robbi Rahim

Host Organizer

PT. Kresna Acitya Nusantara Mediatama

Relawan Jurnal Indonesia

Committee

1. Andri Pramesyanti, Ph.D (UPNV Jakarta – Indonesia)
2. Ifdil, Ph.D (Universitas Negeri Padang - Indonesia)
3. Dr. Aditya H.P.K Putra (Universitas Muslim Indonesia - Indonesia)
4. Dr. Prantasi Harmi Tjahjanti, M.T (Universitas Muhammadiyah Sidoarjo - Indonesia)
5. Fredy Kurniawan, ST, MT, M.Eng., Ph.D (Universitas Narotama - Indonesia)
6. Dr. rer. nat. Bintoro Anang Subagyo (Institut Teknologi Sepuluh Nopember - Indonesia)
7. Ahmad Fudholi, Ph.D (Universiti Kebangsaan Malaysia - Malaysia)

Table of contents

Volume 755

2021

◀ Previous issue Next issue ▶

Annual Conference on Health and Food Science Technology, 25 November 2020, Yogyakarta, Indonesia

Accepted papers received: 16 April 2021

Published online: 05 May 2021

Open all abstracts

Preface

OPEN ACCESS 011001

Preface

+ Open abstract  View article  PDF

OPEN ACCESS 011002

Peer review declaration

+ Open abstract  View article  PDF

Papers

OPEN ACCESS 012001

The Effect of Eggshell Organic Fertilizer on Vegetative Growth of Cayenne Pepper (*Capsicum frutescens* L)

Robert Devi Anugrah, Rafvenia, Meitiyani and Luthpi Safahi

+ Open abstract  View article  PDF

OPEN ACCESS 012002

Spray Drying of Rosella (*Hibiscus sabdariffa* L.) Powder: Effect of Shelf Life on Physicochemical Properties and *Cyanidin 3-O—glucoside*

UA Nur Fitriani, Muhammad Yusuf and Fika Soleha Ilyas

+ Open abstract  View article  PDF

OPEN ACCESS 012003

Acanthocephaliasis And Sparganosis Occurrence in An Asian Vine Snake (*Ahaetulla*

Phaiolepis): A Perspective of Neglected Zoonotic Disease
This site uses cookies. By continuing to use this site, you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Aditya Yudhana, Ratih Novita Praja and Arif Supriyanto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012004

Mechanical Properties of Whey Composite Edible Film with the Addition of Clove Essential Oil and Different Types of Plasticizer

Fahrullah, M Ervandi, M A Indrianti, D O Suparwata, I A Yasin, Y A Gobel, S Mokoolang and M M Mokoginta

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012005

The Effect of 96% Ethanol Extract of Binahong Leaf on Hyperglycemia White Male Rats Using Total Cholesterol and Triglyceride Parameters

Dwitiyanti and Tera Rorenza

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012006

Dragonfly Community in Flowing and Stagnating Water in the Cibodas Botanical Garden Area

Nur Afni Febrianti, Susanti Murwitaningsih, Paskal Sukandar and Suci Lestari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012007

The Effect of Feeding Maggot Fluor (*Hermetia illucens*) on Fish Feed for Growth of Sangkuriang Catfish (*Clarias* sp)

Ridho Budi Al Rizki, Maryanti Setyaningsih, Meitiyani and Irdalisa

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012008

The Effects of Probiotics Administration on Silkworms (*Tubifex* sp) Natural Feed on Growth of Kanca Fish (*Tor soro*)

S. Mulatsih, N.U. Hartanti and A. Najib

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012009

Sustainability Level of Aquaculture Utilization and Its Management Strategy In Conservation Areas Regional Water (KKPD) Biak Numfor District, Papua

Sutaman

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012010

Specific Immunomodulatory Effect of Water Extract of *Stachytarpheta jamaicensis* herbs

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012011

Antihypertensive Effects of Purslane (*Portulaca oleracea*) Extract in Animal Model of Hypertension

Afifah Bambang Sutjiatmo, Suci Nar Vikasari and Falna Bintussolihah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012012

Zebrafish Embryos Immune System induction with BSA as an initial Screening model for Covid-19 Treatment

Dita Arya Widatama and Bambang Retnoaji

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012013

Antibacterial Potential of *Nicotiana tabacum* L. var Virginia Pyrolysis Extract Against *Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, and *Pseudomonas aeruginosa*

S A Fernanda, B A Amru, H A Rahmani, M Gozan, N S Irsyad, M Bahar, O S Puspita, F Zulfa and A Pramono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012014

Meistera chinensis fruit properties: Chemical compound, antioxidant, antimicrobial, and antifungal activity

Musdalipah, Karmilah, Selfyana Austin Tee, Eny Nurhikma, Yulianti Fauziah, Adryan Fristiohady, I Sahidin and Agung Wibawa Mahatva Yodha

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012015

Effect of ammonium chloride and disodium hydrogen phosphate with molasses as a substrate for *Saccharomyces cerevisiae* in the production of β -glucan

Ririn Puspawati, Putranti Adirestuti and Jesiana Agustine Arifin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012016

Proteolytic and Clot Lysis Activity Screening of Crude Proteases Extracted from Tissues and Bacterial Isolates of *Holothuria Scabra*

Nur Hidayati, Hayatun Fuad, Hendra Munandar, Dewi Seswita Zilda, Nurrahman Nurrahman, Miswar Fattah, Oedjijono Oedjijono, Amin Samiasih and Stalis Norma Ethica

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012017

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Pramulani Mulya Lestari, Supandi and Mawar Nusafitri

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012018

The Characteristic of Gelatin Extracted from The Skin of Adult and Sub-Adult Striped Catfish (*Pangasius hypophthalmus*) Using Acid-Base Pretreatment: pH and FTIR

Dewi Hidayati, Gina R. Sabiyla, Endry N. Prasetyo, Noor Nailis Sa'adah and Fredy Kurniawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012019

Assessment of Stranded Marine Debris on Karimunjawa Island

M R Ansyari, I M Radjawane, T Suprijo, F Hanifah and A Tarya

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012020

Application of Geographical Information Systems (GIS) for Mapping Food Vulnerability in Denpasar, Bali

I Ketut Sardiana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012021

Mapping of Coastal Sedimentation using Landsat Satellite Image and Bathymetry Map (Case study: Batu Gong Coastal, Southeast Sulawesi, Indonesia)

Laode Muhammad Golok Jaya

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012022

Application of Supply Chain Requirements for Smallholders: Impact on Sustainable Palm Oil Management Policies in Indonesia

Herdis Herdiansyah, Ratih Dyah Kusumastuti, Palupi Lindiasari Samputra, Neny Indriyana and

Nanik Ambar Suharyanti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012023

Green open space implementation on the underground building

I N Indira and H Herdiansyah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012024

Epidemiological Characteristics of the First Cluster of COVID-19 in Magetan District – East Java, 2020

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012025

Social Consideration for Blue Carbon Management

C.C Pricillia, M.P Patria and H. Herdiansyah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012026

Sustainable Development of Papalele's Household from Environmental Issue

Simon Pieter Soegijono and Esther Kembauw

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012027

Ants Respon to Air Humadity in Small Islands of Haruku

Fransina Latumahina and Gun Mardiatmoko

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012028

Analysis of Changes in Health of Coastal Mangrove Forest on the East Coast of Lampung

Ferdy Ardiansyah and Rahmat Safe'i

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012029

Bontomarannu Education Park Reviewed from Conservation, Social, and Environmental Aspects

Rismawati, Syahribulan and Slamet Santosa

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012030

Collaborative Model to Increase Empowerment of Dryland Farmer (Case Study West Halmahera District, North Maluku Province)

Sitti Aminah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012031

Diversity of terrestrial ferns (*Pteridophytes*) in Ciliwung Telaga Warna Puncak Bogor tea estate in West Java

Basar Wiranto, Husnin and Susilo

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

[View our Privacy and Cookies policy.](#)

012032 

Amphibian Diversity (Order Anura) in Bogor, Indonesia Amphibian Diversity in Cimisblung Bogor West Java

Agus Pambudi Dharma, Apriyanto Saputra and Eka Kartika

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012033

Amphibian Diversity in the Kawasan Taman Wisata Alam, Bogor, Indonesia

Miftah Fauzi, Paskal Sukandar, Maryanti Setyaningsih and Rosi Ritonga

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012034

Integration of Village Development with the Company's CSR Program in the Environmental Priority Program Formulation

D Mardhia, N Kautsari, I W Ayu, Y Hartono and G Gunawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012035

Factors related to the incidence of stunting in Nupabomba and Guntarano Villages, Tanantovea District, Donggala Regency

Hasanudin, Tjitrowati Djaafar, Saharudin and Andi Bungawati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012036

Biodiversity of the "Karang Jeruk" Coral Reef Ecosystem in Tegal Regency, Central Java, Indonesia

Noor Zuhry, Djoko Suprpto, Supriharyono and Boedi Hendrarto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012037

Success Rate Of Mangrove Crab (*Scylla Serrata*) Molting With Different Salinity Treatments In Controlled Containers

Suyono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012038

Assessment of Spring Water Quality Affected by Agricultural and Human Activities in Bali Island

I Gusti Ngurah Agung Suryaputra, I Wayan Yudi Artawan and Made Vivi Oviantari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012039

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#) using Different Enzymet Methods



Kristhina Priskila Rahael, Santi Penina Tua Rahantoknam and Syahibul Kahfi Hamid

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012040

The effect of giving Kelakai (*Stenochlaena palustris*) juice on the volume of breastfeeding for postpartum mothers in the working area of the Pekauman Health Center, Banjarmasin

Dede Mahdiyah, Dwi Rahmawati and Desy Susila Waty

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012041

Isolation, Characterization, Molecular Identification of Probiotic Bacteria from Meconium

Dhanly Arie Irza, I Nyoman Ehrich Lister, Saipul Sihotang and Edy Fachrial

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012042

Extract of *Caesalpinia sappan* L. heartwood as food treatment anti-diabetic: a narrative review

A F Muti, D L C Pradana and E P Rahmi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012043

The effect of Indonesian honey *Tetragonula* sp. and Indonesian royal jelly *Apis mellifera* (*Ceiba pentandra*) to human preputium cell proliferation in serum-free DMEM

Q S Fachrani, M A Dhifanra, Y Nugraha, R Yulianti, M Sahlan, S Pambudi, A Nurhasanah and A Pramono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012044

Embryogenic Callus Induction of Todolo Toraja Coffee Leaf Cells (*Coffea arabica* Var. Typica) with the Addition of 2, 4-Dichlorophenoxyacetic Acid (2, 4-D) and Furfurylaminopurine (Kinetin) in Vitro

Putri Damayanti, Andi Ilham Latunra and Eva Johanes

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012045

Citrus maxima Pectin as Superdisintegrant: Preparation and Evaluation of Dextromethorphan Hydrobromide Orodispersible Film

Nining, Rahmah Elfiyani and Siti Nurhasanah

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

012046

Formulation and Development of Grape Seed Oil (*Vitis Vinifera* L) Emulgel Peel-Off Mask using Gelling Agent Hydroxy Propyl Methyl Cellulose (HPMC)

Fith Khaira Nursal, Nining and At Syamsiyah Rahmah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012047

Soy Milk Filter Design Using Dfma Method

Irfan Santosa, Galuh Renggani Wilis and Urip Mulyadi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012048

Implementation of Experimental Designs to Improve Bioethanol Quality From Banana Beans Through the Destilation Process

Angga Adhy Nugroho, Agus Wibowo and Saufik Luthfianto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012049

The Utilization of Tofu Waste Water as An Addition of Nutrition in Hydroponic Media to Lettuce Growth (*Lactuca sativa* L.)

Arnita Chaorlina, Maryanti Setyaningsih and Hilman Faruq

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012050

Effectiveness of Feeding Trash Fish and Spinach Extract on Mud Crab (*Scylla Serrata*) Feed for Molting Acceleration With the Popeye Method

Suyono, NU Hartanti and NHSA

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012051

Effect of Differences in Salt Concentration on the Quality of Rebon Shrimp Paste (*Acetes* Sp) in Tegal District

S Mulyani, P M Vestiyati, Kusnandar, H K Alamsyah and S W Simanjuntak

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012052

The Effectiveness Of Cod Reduction In Tofu Waste Using Active Mud And Oxygenation Methods

Wati Sukmawati and Wijastuti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012053

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

Application of Microcapsules Food with Different Dosage to Support The Growth of Fish Cork Seed (*Channa Striata*)



Hayati Soeprapto, Hadi Pranggono and Febri Mustafat Ridwan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012054

The Effectiveness of Sludge in the Leaf Fermentation Process

Yuni Astuti, Syamsul Arifin Syarifudin, Maesyaroh and Ranti An Nisa

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012055

The Characteristic of Yellow Pumpkin Flour That Has Been Processed Using Shard Gourd Method

Ida Farikha Azizah, PN Ika Mulawati, Ngatinem and Novi Kuswardani

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012056

Actualization of local community participation in critical land management in Gorontalo

D O Suparwata, D Rukmana, A N Tenriawaru and R Neswati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012057

The Effect Of Transfer Of Land Function On The Development Of The Area Through Community Empowerment

R.R.A Tarigan, H.B. Tarmizi, A Purwoko and R. Ginting

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012058

Environmental Factors for *Holothuria scabra* Sea Cucumber Cultivation

Santi Penina Tua Rahantoknam, Mariana Yermira Beruatjaan, Meyske Angel Rahantoknam and Maria Kristina Ohoiwutun

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012059

Natural Antibacterial and its Use

Sandriana J Nendissa, Meta Mahendradatta, Zainal and Februadi Bastian

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012060

The Effect of Ethanol Concentrations as The Extraction Solvent on Antioxidant Activity of Katuk (*Sauropus androgynus* (L.) Merr.) Leaves Extracts

Ni Putu Ermi Hikmawanti, Sofia Fatmawati and Anindita Wulan Asri

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Katuk is widely popular with its benefits for breastfeeding mothers. Katuk is also known as a plant with a high antioxidant content. This study aims to determine the effect of using variations in the ethanol concentration as an extracting solvent in producing Total Phenolics Content (TPC) and Total Flavonoids Content (TFC) and their activities in reducing DPPH free radicals. The dried katuk leaves were extracted by cold maceration method. The solvent used for extraction is ethanol with 3 variations in concentration: 50%, 70%, and 96% (absolute ethanol). TPC and TFC were determined by colorimetric method using a UV-Vis spectrophotometer. TPC was stated to be equivalent to gallic acid, while TFC was stated to be equivalent to quercetin. DPPH free radical scavenging activity was measured based on the IC₅₀ value. The results showed that Katuk leaf extract produced from 50% ethanol solvent was able to produce TPC (42.18 ± 0.30 mg GAE / g), TFC (11.18 ± 0.38 mg QE / g) and reduction activity against DPPH radicals (IC₅₀ = 88.33 ± 3.53 ppm). These were higher than ethanol with other concentrations. However, various things need to be considered when using this solvent given the high water content in the solvent.

<https://doi.org/10.1088/1755-1315/755/1/012060>

OPEN ACCESS

012061

The Use of Bioremediation Technology on Oil-Contaminated Soil

Putri Alvernia, Suyud Utomo, Tri Edhi Budhi Soesilo and Herdis Herdiansyah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012062

The Effect Of Microwave AID Extraction Temperature Of Kedawung Leaves (*Parkia biglobosa*) On Antioxidant Activity And Flavonoid Leavels

Ferika Indrasari and Buanasari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012063

Preparation and Evaluation of Plant Extract Microcapsules Using Chitosan

Buanasari, Warlan Sugiyo and Heri Rustaman

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012064

Twenty weeks of *Centella asiatica* improved cognitive function of women elderly with dementia

Lisna Anisa Fitriana, Kusnandar Anggadiredja, Setiawan and I Ketut Adnyana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#)

012065

Literature Review: Potential Pharmacological Activity of *Luffa cutangula* L. Roxb

Erna Harfiani and Dhigna Citra Pradana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012066

The Effect of Adding Kepok Banana Peels (*Musa paradisiaca*) to Powder Media on the Growth of White Oyster Mushrooms (*Pleurotus ostreatus*)

Riris Risnawati, Meitayani and Susilo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012067

Traditional Medicinal Plants in Bima Communities: A Bacterial Activities Test and Phytochemicals

N Azmin, M Nasir, H Hartati, A Ariyansyah and F Fahrudin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012068

Evaluation of Ipomea carnea Growth Response in Plant Media That was Exposed by Sidoarjo Mud

S.C Sutarman, S Pamungkas and M. Abror Arifin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012069

Material and Life Service of the Septic Tank Have an Influence on the Biological Pollution of Groundwater (Case Study Kelurahan Pademangan Barat, North Jakarta)

Novita Anggraini, Cindy Rianti Priadi and Herdis Herdiansyah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012070

Soil Water Content Estimation using Pedotransfer Functions in Drylands of Sumbawa Regency

I W Ayu, N D Lestari, G Gunawan and S Soemarno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012071

Sustainable Development Through the One Village One Product (OVOP) Approach for Local Commodities

Aveanty Miagina, Hernansi Bisio and Esther Kembauw

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012072

The Development Strategy of Conservation for *Cacatua sulphurea* in Bontomarannu

Education Park, Gowa Regency Animal Park
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Irny Novriyani, Syahribulan and Eddy Soekendarsi

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012073

Design and Development of a Geospatial-Based Information Systems for Disaster Management of Adolescent Reproductive Health in Nusa Tenggara Barat Province In 2020

M. Sapoan Hadi, Sutanto P. Hastono and Artha Prabawa

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012074

The Role of Diffusion of Innovation in Agricultural to Compete in Asean Community

Harisa Mardiana and Esther Kembauw

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012075

Village Development Program for Producing Iodized Salt for Sustainable Development Community

D Mardhia, D Syafikri, F Yahya, N Andriyani and G Gunawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012076

Reducing solid waste through waste banks: an empirical study in Kepulauan Riau, Indonesia

W E Yudiatmaja, Edison, T Samnuzulsari, Yudithia, S R I Rezeki, Suyito, D Akbar and A Alfiandri

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012077

Education of Waste Management Based on Zero Waste in Kendal District (Case Study: Waste Recycling Craft Community (Kerdus), Kendal District, Central Java)

Nanik Hidayati, Neli Hajar and Feri Setiyanto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012078

The Utilization of Vegetable Waste as a Nutrient Addition in Hydroponic Media for the Growth of Green Mustard (*Brassica juncea L.*)

Hilman Faruq, Elsa Novelia, Maryanti Setyaningsih and Ranti An Nisa

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012079

Monthly Rainfall Components in Ambon City: Evidence from the Serious Time Analysis

Lexy J. Sinay and Esther Kembauw

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

[see our Privacy and Cookies policy](#) [PDF](#)



-
- OPEN ACCESS** 012080
The Effect of Climate on the Outbreak of Covid-19: A Review
Candra Kirana, Samsul, Hesty Yariska Hapsir, Fitria Nurmalita and Ramadhan Tosepu
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012081
An Impact of Climatic Change on Water-borne Diseases: A Review
Muhammad Fadhil, Ruswan, Mutia Ismail, Helyani and Ramadhan Tosepu
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012082
Risk of Heat Stroke due to Extreme Heat Changes: A Review
Mirnawati, Nurnaningsih, Rina Asriani, I Amirullah and Ramadhan Tosepu
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012083
A Systematic literature (Impact of Climate Change on Filariasis)
Roswati, Santi, Ahmad Syafii, Eke Mahendra and Ramadhan Tosepu
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012084
Literature Review: Impact Of Temperature And Rainfall On Incident Malaria
Ladjumadil Ahmad Tiu, Waode Elfianti Wahid, Waode Yuli Andriani, Mirnawati and Ramadhan Tosepu
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012085
Increased Risk of Hepatitis A due to Weather Changes:A Review
Adibin, Aisnah, Indrawati, Sumriati and Ramadhan Tosepu
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012086
Fertilizer Application, Climate Change and Rice Production in Rural Java
Suko Irawan and Ernoiz Antriyandarti
[+ Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012087
Impacts and Adaptation Strategies on Climate Variability and Change of Coastal Communities along Banate Bay
B Johnny, Ed Dolor and D.
[+ Open abstract](#) [View article](#) [PDF](#)



OPEN ACCESS

012088

Impact of Weather and Climate on Diarrhea Incidence : A review

Idham Malik, Sri Anjayati, Pipit Musdhalifa, Dwiana Binti and Ramadhan Tosepu

[+ Open abstract](#) [View article](#) [PDF](#)**OPEN ACCESS**

012089

A Systematic Literatur The Impact Of The Climate To The Case Of Tuberculosis (TB): A Review

Al Edy Dawu, Resti Nur Pratiwi, Samsiah Winda, Agus Supriyanto Suparno and Ramadhan Tosepu

[+ Open abstract](#) [View article](#) [PDF](#)**OPEN ACCESS**

012090

Study of the Use of Block Compos on the Growth of Teak (*Tectona grandis*) in Used Lands of Kapur Stone Mine

Supiana Dian Nurtjahyani, Dwi oktafitria, Sriwulan, Ahmad Zaenal Arifin, Eko Purnomo, Aris Santoso and Ali Mustofa

[+ Open abstract](#) [View article](#) [PDF](#)**JOURNAL LINKS**[Journal home](#)[Journal scope](#)[Information for organizers](#)[Information for authors](#)[Contact us](#)[Reprint services from Curran Associates](#)

PAPER • OPEN ACCESS

The Effect of Ethanol Concentrations as The Extraction Solvent on Antioxidant Activity of Katuk (*Sauropus androgynus* (L.) Merr.) Leaves Extracts

To cite this article: Ni Putu Ermi Hikmawanti *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **755** 012060

View the [article online](#) for updates and enhancements.

The Effect of Ethanol Concentrations as The Extraction Solvent on Antioxidant Activity of Katuk (*Sauropus androgynus* (L.) Merr.) Leaves Extracts

Ni Putu Ermi Hikmawanti*, Sofia Fatmawati, Anindita Wulan Asri

Faculty of Pharmacy and Science, Universitas Muhammadiyah Prof. DR. HAMKA Klender, East Jakarta, DKI Jakarta, 13460, Indonesia

*ermy0907@uhamka.ac.id

Abstract. Katuk is widely popular with its benefits for breastfeeding mothers. Katuk is also known as a plant with a high antioxidant content. This study aims to determine the effect of using variations in the ethanol concentration as an extracting solvent in producing Total Phenolics Content (TPC) and Total Flavonoids Content (TFC) and their activities in reducing DPPH free radicals. The dried katuk leaves were extracted by cold maceration method. The solvent used for extraction is ethanol with 3 variations in concentration: 50%, 70%, and 96% (absolute ethanol). TPC and TFC were determined by colorimetric method using a UV-Vis spectrophotometer. TPC was stated to be equivalent to gallic acid, while TFC was stated to be equivalent to quercetin. DPPH free radical scavenging activity was measured based on the IC₅₀ value. The results showed that Katuk leaf extract produced from 50% ethanol solvent was able to produce TPC (42.18 ± 0.30 mgGAE / g), TFC (11.18 ± 0.38 mgQE / g) and reduction activity against DPPH radicals (IC₅₀ = 88.33 ± 3.53 ppm). These were higher than ethanol with other concentrations. However, various things need to be considered when using this solvent given the high water content in the solvent.

1. Introduction

Sauropus androgynus (L.) Merr in Indonesia is known as katuk (family Phyllanthaceae). Katuk contains high nutrients in the form of vitamins, minerals, fiber, lipids, carbohydrates, as well as bioactive compounds such as phenolics, tannins, flavonoids, anthocyanins, phytosterols, and so on. Katuk is included in the group of green vegetables with ethnomedicine potential, including as an antitussive, tonic, antipyretic, breast milk facilitator, and others (Petrus, 2013). Katuk was also reported to have aphrodisiac activity (Rusdi et al., 2018) and fertility enhancers in male rats given orally (Hikmawanti et al., 2020). The antioxidant capacity of katuk leaves associated with various pharmacological activities has also been discovered such as antimicrobial, cytotoxic against cancer cells, anti-inflammatory and wound healing (Petrus, 2013; Khoo et al., 2015).

Antioxidants are substances that are used in small concentrations to inhibit and/or to reduce oxidation caused by an oxidant. Plants are a source of natural ingredients with great natural antioxidant potentials, such as phenolic compounds (phenolic acids, flavonoids, tannins, anthocyanins, lignins), carotenoids, vitamins (vitamins A, C, E) and so on. To obtain these antioxidant compounds, an extraction process is necessary (Altemimi et al., 2017). Extraction is a procedure performed to obtain metabolites in plants such as alkaloids, phenolics, flavonoids, glycosides, and others using selective solvents. Solvent



selection is one of the stages of preparation for extraction (Azwanida, 2015). The selectivity of the solvent to extract the target compound from a plant material is related to the polarity compatibility of the two. Ethanol is a solvent that is safe for human consumption as a solvent for natural substances for both food and natural medicinal purposes. Absolute ethanol and aqueous ethanol have been used successfully to extract phenolic derivative antioxidant compounds from natural ingredients with good results (Sultana et al., 2009).

The purpose of this study is to determine the total phenolic content (TPC), total flavonoid levels (TFC), and the antioxidant activity against DPPH radicals from katuk leaf extract produced from three types of variations in the ethanol concentrations as an extracting solvent (50%, 70% and 96%). It is important to discover ethanol concentrations to produce katuk leaf extract with a high content of antioxidant compounds. In addition, with the right solvent, extracts with guaranteed safety and quality can be produced and applied widely.

2. Methodology

Materials

Fresh leaves of katuk were obtained from the Biopharmaca Cultivation Conservation Unit (UKBB), Center for Tropical Biopharmaca Study, LPPM IPB, Bogor, West Java, Indonesia. Plants were determined in the same place. Ethanol as a solvent extraction in analytical grade was obtained from Merck in Darmstadt, Germany. Gallic acid, quercetin, and DPPH (2,2-diphenyl-1-picryl-hydrazyl) were obtained from Sigma-Aldrich Co. in St. Louis, the US.

Extraction

The extraction of dried katuk leaf powder (150.0 g) was carried out by cold maceration method following the Indonesian Herb Pharmacopoeia. The solvent used was ethanol with various concentrations, namely 50%, 70% and 96% (absolute ethanol). Each extraction process was carried out for 24 hours. The filtrate was separated from the residue using Whatmann filter paper. The residue was remacerated 3 times. The filtrate from each solvent was collected and evaporated using a rotary evaporator N-1200 BS series from EYELA (Shanghai, China) at 50 ° C and using a water bath at 50 ° C until a thick extract was obtained (Ministry of Health Republic of Indonesia, 2008).

Determination of the Physico-Chemical Properties of the Extract

The physico-chemical parameters of katuk leaf extract including organoleptic, percentage of extract yield, drying loss, and total ash content were determined according to the procedure stated in the Indonesian Herb Pharmacopoeia (Ministry of Health Republic of Indonesia, 2008).

Phytochemical screening

Phytochemical screening was carried out qualitatively using detection reagents based on the procedures explained in Hanani (2015) and the Indonesian Herb Pharmacopoeia (Ministry of Health Republic of Indonesia, 2008). The classes of compounds identified in the extract included phenolics, flavonoids, tannins, saponins, alkaloids, steroids and triterpenoids.

Determination of Total Phenolics Content (TPC)

TPC determination followed Yang et al., (2007)'s procedure with modification using Folin-Ciocalteu Reagent (FCR) and gallic acid as standard. The calibration curve was made using gallic acid solution with a concentration range of 18-66 ppm. The relationship between the concentration of gallic acid (x) and its absorbance (y) was plotted to produce a linear line equation ($y = bx \pm a$). Each 1000 ppm katuk leaf ethanol extract in ethanol was pipetted as much as 0.3 mL. It was then added 1.5 mL FCR reagent (which has been diluted 1:10) and homogenized. The solution was incubated for 3 minutes, then 1.2 mL of 7.5% Na₂CO₃ solution and water were added to obtain a total volume of 10 mL. The mixture was incubated again for 60 minutes at a room temperature. The absorbance of the extract solution was measured using a UV-Vis Shimadzu UV-1601 Series (Kyoto, Japan) spectrophotometer with a

maximum wavelength of 756.5 nm at room temperature. The absorbance obtained was plotted into a linear line equation and then the TPC was calculated using the formula: phenolic content in solution ($\mu\text{g} / \text{mL}$) multiplied by the dilution factor and the solution volume (mL) and divided by the extract weight (g). TPC was expressed in mg which is equivalent to gallic acid per gram of extract. Each extract was tested with 5 repetitions and reported as mean \pm SD.

Determination of Total Flavonoids Content (TFC)

TFC determination followed the procedure in Chang et al., (2002) with modifications. The standard used was quercetin. Quercetin solutions with a concentration range of 33-129 ppm were used to create a calibration curve. The relationship between the concentration of quercetin (x) and its absorbance (y) was plotted to produce a linear line equation ($y = bx \pm a$). Each ethanol extract of katuk leaves with a concentration of 7500 ppm in methanol was pipetted 0.5 mL. It was then added 1.5 mL of methanol and 0.1 mL of 10% AlCl_3 . 0.1 mL of sodium acetate 1M was added and made sufficient with aquadest up to 5 mL. The solution was incubated for 60 minutes at room temperature. The absorbance of the extract solution was measured with a UV-Vis Shimadzu UV-1601 Series (Kyoto, Japan) spectrophotometer at a wavelength of 434 nm. The absorbance obtained was plotted into a linear line equation and then the TFC was calculated using the formula: the level of flavonoids in solution ($\mu\text{g} / \text{mL}$) multiplied by the dilution factor and the solution volume (mL) and divided by the extract weight (g). TFC was expressed in mg which is equivalent to quercetin per gram of extract. Each extract was tested with 5 repetitions and reported as mean \pm SD.

Antioxidant Activity Testing

The antioxidant activity test of katuk leaf extract was carried out using the DPPH method following the procedure of Wan et al., (2011) with modifications. Quercetin was used as a comparison. The extract was dissolved in methanol and diluted into 5 variations of concentration: 20, 40, 60, 80, and 100 ppm. Quercetin solution was dissolved in methanol and diluted to 2, 4, 6, 8, and 10 ppm. Each variation of the concentration extract solution or quercetin was pipetted 1 mL and was added 1 mL of fresh 0.5 mM DPPH in methanol. The solution was made sufficient with methanol to a total volume of 5 mL. The mixture in a tube lined with aluminum foil paper was incubated for 30 minutes at a room temperature in dark conditions. Blank solution was made from 1 mL DPPH mixed with 4 mL methanol. Furthermore, the absorption was measured at a wavelength of 515.5 nm. Each concentration series of each sample was tested triplo. Results were reported as mean \pm SD. The calculation of the percentage of DPPH radical inhibition used the formula:

$$\text{Inhibition of DPPH (\%)} = \frac{Ab - As}{Ab} \times 100$$

where Ab is the absorbance of the blank and As is the absorbance of the sample.

Data Analysis

Data were analyzed with a one-way analysis of variance (ANOVA) followed by Tukey's test ($\alpha \leq 0.05$).

3. Result and Discussion

The ethanol extract of katuk leaves in this study had organoleptic characteristics with a distinctive odor, thick shape, slightly sweet taste and dark brown color (Table 1). The three extracts had an ash content value in the range 5.74-8.23%, w / w. Ash content describes the amount of inorganic material contained in the katuk leaf extract, both obtained internally by plants and during the extraction process. The percentage of ash content of the katuk leaf ethanol extract obtained in this study (<10%) indicated that the extract had a fairly good quality. Ash content is a contaminant parameter that is difficult to remove in extracts but can still be controlled by proper post-harvest handling of plants and a maintained extraction process (Kunle et al., 2012). The drying shrinkage value describes the amount of compound lost during the 105°C heating process for several hours. The drying shrinkage value of the three katuk

leaf extracts was in the range of 1.96-4.96%, w / w (<5%) (Table 1.). This value range showed that the ethanol extract of katuk leaves had a good quality.

The characteristic parameter of the extract determined was the chemical content of the ethanol extract of katuk leaves. The results of phytochemical screening of the three ethanol extracts of katuk leaves showed the presence of alkaloids, phenolics, flavonoids, tannins, saponins and steroids (Table 2.). Although the ethanol solvent used for extraction was from different concentrations, it was still able to attract the same type of compound class to the katuk leaves. Ethanol has the ability to attract glycosides (Houghton & Raman, 1998), polyacetylenes, sterols (Tiwari et al., 2011), polyphenols, tannins, flavonols, terpenoids, and alkaloids (Azmir et al., 2013).

The choice of ethanol as the extraction solvent was considered to provide many advantages over other organic solvents, which is relatively safer (less toxic). Ethanol is a protic organic solvent with a polarity index value of 5.2 (Synder, 1974; Abarca-Vargas et al., 2016) and a dielectric constant of 24.55. Absolute ethanol contains 0.01% by volume of water (Covington & Dickinson, 1973). The 50% ethanol solvent in this study was able to extract more metabolites from the dry powder of katuk leaves (37.77%) than other ethanol solvents (70% ethanol > 96% ethanol). The ethanol solvent polarity of the three concentrations used in this study was influenced by the high concentration of water contained in ethanol. The more water contained in it, the higher its polarity compared to absolute ethanol (Tiwari et al., 2011). Solvents with high polarity had the ability to extract a class of compounds with a wider polarity. This allowed non-phenolic polar compounds such as carbohydrates and proteins to be dissolved during the extraction process which resulted in increased extraction yields (Do et al., 2014).

Phenolic is a plant-produced metabolite with a distinctive structure in the form of an aromatic that binds to one or more -OH rings. Meanwhile, flavonoids are phenolic derivative pigment metabolites which are widely found mainly in higher plants. One of the quantitative determinations of phenolics and flavonoids can be done colorimetrically using the UV-Vis spectrophotometer technique (Khoddami et al., 2013). This technique is considered relatively simple, fast, and is able to provide satisfactory quantitative results (Cong-cong et al., 2017). Quantitative determination of phenolics was carried out using the Folin-Ciocalteu Reagent (FCR) containing tungsten and molybdenum. The product of this reaction was a blue solution ($(\text{PM}_6\text{W}_{11}\text{O}_{40})^{4-}$) which was absorbed at a wavelength of about 760-765 nm (Prior et al., 2005; Khoddami et al., 2013). FCR has a weakness, namely that this reagent is also able to react with other compounds such as vitamin C, aromatic amines and saccharides (Cong-cong et al., 2017). Meanwhile, the quantitative determination of flavonoids was carried out using 10% AlCl_3 reagent which was carried out under alkaline conditions such as if 7.5-20% Na_2CO_3 was added. The uptake of this reaction product can be measured at a wavelength of about 410-423 nm (Khoddami et al., 2013). Based on the results in Table 3., the best TPC (42.18 mgGAE / g) and TFC (11.18 mgQE / g) were obtained in katuk leaf extract extracted with 50% ethanol (70% ethanol > 96% ethanol) solvent. These results indicated that 50% ethanol solvent is the best solvent for extracting antioxidant compounds from katuk leaves. Petrus (2013) explained that dried katuk leaves had TPC and TFC of 1150.95-2300.00 mgGAE / 100 g and 1040 mgRE / 100 g, respectively.

Antioxidant activity testing in this study was carried out using DPPH radicals. This method is considered relatively inexpensive, effective, efficient and has good sensitivity of a UV-Vis spectrophotometer. DPPH is a free radical with a dark blue color. With the presence of an antioxidant, the antioxidant would donate its Hydrogen atom to the DPPH radical (becoming a DPPH-H). This reaction is characterized by a decrease in absorbance due to DPPH loses its reactivity (Alam et al., 2013; Huyut et al., 2017). The range of quercetin concentrations as a comparison used in this test was 2-10 ppm. Meanwhile, the extract concentration used was 20-100 ppm. The concentration range used was adjusted to the reactivity of each material against the DPPH radical. In addition, other considerations were based on the limitations of the minimum and maximum sample absorption values that can still be read by the UV-Vis spectrophotometer instrument at a wavelength of 515-517 nm (Maesaroh et al., 2018). The antioxidant activity in this study was determined based on the IC_{50} value of the sample. The IC_{50} value showed the sample concentration needed to inhibit 50% of DPPH radicals. This value was obtained from the results of calculations using linear regression analysis. The smaller the IC_{50} value of

an antioxidant, the higher its power to inhibit free radicals (Do et al., 2014). Based on the results in Table 3., the ethanol extract of 50% katuk leaves had the highest antioxidant activity with an IC₅₀ value of 88.33 ppm (70% ethanol > 96% ethanol). This showed that the ethanol extract of 50% katuk leaves had a strong DPPH radical scavenging activity (<100 ppm).

The phenolic and flavonoid contents in katuk leaves play an important role in its activity as a source of antioxidants. The number of hydroxyl groups in a phenolic molecule affects its capacity as an antioxidant. Phenolic has a tendency to donate hydrogen atoms or electrons from its hydroxyl groups to free radicals (Dai & Mumper, 2010). As for flavonoids, the number and location of the aromatic hydroxyl groups in their structure affect their antioxidant capacity (Fernandez-Panchon et al., 2008).

4. Conclusion

Based on this study, the variation in the concentration of ethanol used as the extraction solvent was able to have an effect on the acquisition of phenolic and flavonoid compounds as well as the DPPH free radical scavenging activity from the katuk leaf extract. The 50% ethanol solvent was able to extract high amounts of phenolic and flavonoids from katuk leaf powder using the cold maceration method compared to ethanol solvent with a concentration of 70% and 96%. The best antioxidant activity was also obtained from katuk leaf extract which was extracted using 50% ethanol solvent. Thus, the selection of this solvent was proven to be easy and efficient to use to produce katuk leaf extract which is rich in antioxidants, both as a raw material for functional food and for natural medicine. However, it is necessary to consider again the amount of time and high energy required in the evaporation process, as well as the stability associated with the safety of the extract produced from the solvent so that the quality and quantity of the extract are guaranteed.

5. Acknowledgment

The authors would like to thank Dr. apt. Hadi Sunaryo, M.Si. as Dean of the Faculty of Pharmacy and Science, Universitas Muhammadiyah Prof. DR. HAMKA for the support of laboratory facilities which made this research completed properly. Our gratitude are also directed to the Scientific Publication Development and Development Unit, Universitas Muhammadiyah Prof. DR. HAMKA who facilitated the publication of this research.

References

- [1] Abarca-Vargas, R., Malacara, C. F. P., & Petricevich, V. L. (2016). Characterization of chemical compounds with antioxidant and cytotoxic activities in bougainvillea x buttiana holttum and standl, (Var. rose) extracts. *Antioxidants*, 5(45), 1–11.
- [2] Alam, M. N., Bristi, N. J., & Rafiquzzaman, M. (2013). Review on in vivo and in vitro methods evaluation of antioxidant activity. *Saudi Pharmaceutical Journal*, 21(2), 143–152.
- [3] Altemimi, A., Lakhssassi, N., Baharlouei, A., & Watson, D. G. (2017). Phytochemicals: Extraction, Isolation, and Identification of Bioactive Compounds from Plant Extracts. *Plants*, 6(42), 1–23.
- [4] Azmir, J., Zaidul, I. S. M., Rahman, M. M., Sharif, K. M., Mohamed, A., Sahena, F., ... Omar, A. K. M. (2013). Techniques for extraction of bioactive compounds from plant materials: A review. *Journal of Food Engineering*, 117(4), 426–436.
- [5] Azwanida, N. N. (2015). Medicinal & Aromatic Plants A Review on the Extraction Methods Use in Medicinal Plants , Principle , Strength and Limitation. *Medicinal and Aromatic Plants*, 4(3), 1–6.
- [6] Chang, C.-C. , Yang, M.-H. , Wen, H.-M. , & Chern, J.-C. (2002). Estimation of Total Flavonoid Content in Propolis by Two Complementary Colorimetric Methods. *Journal of Food and Drug Analysis*, 10(3), 178–182.
- [7] Cong-cong, X. U., Bing, W., Yi-qiong, P. , Jian-sheng, T. A. O., & Tong, Z. (2017). Advances in extraction and analysis of phenolic compounds from plant materials. *Chinese Journal of*

- Natural Medicines*, 15(10), 0721–0731.
- [8] Covington, A. K. ., & Dickinson, T. (1973). Introduction and Solvent Properties. In *Physical Chemistry of Organic Solvent Systems* (p. 5). New York: Plenum Press.
- [9] Dai, J., & Mumper, R. J. (2010). Plant phenolics: Extraction, analysis and their antioxidant and anticancer properties. *Molecules*, 15(10), 7313–7352.
- [10] Do, Q. D., Angkawijaya, A. E., Tran-Nguyen, P. L., Huynh, L. H., Soetaredjo, F. E., Ismadji, S., & Ju, Y.-H. (2014). Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. *Journal of Food and Drug Analysis*, 22, 296–302.
- [11] Fernandez-Panchon, M. S., Villano, D., Troncoso, A. M., & Garcia-Parrilla, M. C. (2008). Antioxidant Activity of Phenolic Compounds: From In Vitro Results to In Vivo Evidence. *Critical Review in Food Science and Nutrition*, 48, 649–671.
- [12] Hanani, E. (2015). *Analisis Fitokimia (In Bahasa)*. Jakarta: Buku Kedokteran EGC.
- [13] Hikmawanti, N. P. E., Rusdi, N. K., & Yulida, S. (2020). Evaluation of sperm quality in male rats treated with *Sauropus androgynus* (L.) Merr. leaf fractions. *Pharmaciana*, 10(2), 193–200.
- [14] Houghton, P. J. ., & Raman, A. (1998). Laboratory Handbook for the Fractionation of Natural Extracts. In *Laboratory Handbook for the Fractionation of Natural Extracts*. London: Springer Science Business Media.
- [15] Huyut, Z., Beydemir, F. ., & Gülçin, E. (2017). Antioxidant and Antiradical Properties of Selected Flavonoids and Phenolic Compounds. *Biochemistry Research International*, 2017, 1–10.
- [16] Khoddami, A., Wilkes, M. A., & Roberts, T. H. (2013). Techniques for analysis of plant phenolic compounds. *Molecules*, 18(2), 2328–2375.
- [17] Khoo, H. E., Azlan, A., & Ismail, A. (2015). *Sauropus androgynus* Leaves for Health Benefits: Hype and the Science. *The Natural Products Journal*, 5, 115–123.
- [18] Kunle, O. F. ., Egharevba, H. O. ., & Ahmadu, P. O. (2012). Standardization of Herbal Medicines - A Review. *International Journal of Biodiversity and Conservation*, 4(3), 101–112.
- [19] Maesaroh, K. ., Kurnia, D. ., & Anshori, J. A. (2018). Perbandingan Metode Uji Aktivitas Antioksidan DPPH, FRAP, dan FIC terhadap Asam Askorbat, Asam Galat dan Kuersetin. *Chimica et Natura Acta*, 6(2), 93–100.
- [20] Ministry of Health Republic of Indonesia. (2008). *Farmakope Herbal Indonesia (Indonesian Herb Pharmacopoeia)* (1st Ed). Jakarta: Ministry of Health Republic of Indonesia.
- [21] Petrus, A. J. A. (2013). *Sauropus androgynus* (L.) Merrill - a potentially nutritive functional leafy-vegetable. *Asian Journal of Chemistry*, 25(17), 9425–9433.
- [22] Prior, R. L., Wu, X., & Schaich, K. (2005). Standardized Methods for the Determination of Antioxidant Capacity and Phenolics in Foods and Dietary Supplements. *Journal of Agricultural and Food Chemistry*, 53, 4290–4302.
- [23] Rusdi, N. K., Hikmawanti, N. P. E., Maifitrianti., Ulfah, Y. S., & Annisa, A. T. (2018). Aktivitas Afrodisiaka Fraksi dari Ekstrak Etanol 70% Daun Katuk (*Sauropus androgynus* (L.) Merr) Pada Tikus Putih Jantan. *Pharmaceutical Sciences and Research (PSR)*, 5(3), 123–132.
- [24] Sultana, B., Anwar, F., & Ashraf, M. (2009). Effect of Extraction Solvent/Technique on the Antioxidant Activity of Selected Medicinal Plant Extracts. *Molecules*, 14, 2167–2180.
- [25] Synder, L. R. (1974). in The Netherlands. *Journal of Chromatography*, 92, 223–230.
- [26] Tiwari, P., Kumar, B., Kaur, M., Kaur, G., & Kaur, H. (2011). Phytochemical screening and Extraction: A Review. *Internationale Pharmaceutica Scientia*, 1(1), 98–106.
- [27] Wan, C., Yu, Y., Zhou, S., Liu, W., Tian, S., & Cao, S. (2011). Antioxidant activity and free radical-scavenging capacity of *Gynura divaricata* leaf extracts at different temperatures. *Pharmacognosy Magazine*, 7(25), 40–45.
- [28] Yang, J., Paulino, R., Janke-Stedronsky, S., & Abawi, F. (2007). Free-radical-scavenging activity and total phenols of noni (*Morinda citrifolia* L.) juice and powder in processing and storage. *Food Chemistry*, 102(1), 302–308.

Table 1. Characteristics of Katuk Leaf Ethanol Extract

No	Parameter	Katuk Leaf Extract		
		Ethanol 50%	Ethanol 70%	Ethanol 96%
1	Organoleptis Organoleptic			
	Smell	Typical	Typical	Typical
	Shape	Thick extract	Thick extract	Thick extract
	Taste	Little sweet	Little sweet	Little sweet
	Color	Dark brown	Dark brown	Dark brown
2	Extraction Yield (% w/w)	37.77 ± 0.93	36.26 ± 3.38	33.55 ± 2.77
3	Ash content (% w/w)	5.74 ± 0.48	7.13 ± 1.08	8.23 ± 0.61
4	Shrink drying (% w/w)	4.96 ± 1.19	1.96 ± 0.75	4.06 ± 0.82

Note: Data presented are mean (n = 5) ± SD.

Table 2. Results of Phytochemical Screening of Ethanol Extract of Katuk Leaves

No	Compound Group	Katuk Leaf Extract		
		Ethanol 50%	Ethanol 70%	Ethanol 96%
1	Alkaloids	+	+	+
2	Phenolic	+	+	+
3	Flavonoids	+	+	+
4	Tannin	+	+	+
5	Saponins	+	+	+
6	Steroids	+	+	+
7	Terpenoids	-	-	-

Description: (+) = detected compound identified; (-) = no identified compound was detected

Table 3. Total Phenolic and Flavonoid Levels of Katuk Leaf Ethanol Extract

Parameter	Katuk Leaf Extract		
	Ethanol 50%	Ethanol 70%	Ethanol 96%
TPC (mgGAE/g)	42.18 ± 0.30*	25.33 ± 0.13	16.25 ± 0.16
TFC (mgQE/g)	11.18 ± 0.38*	8.87 ± 0.14	5.68 ± 0.34
IC₅₀ (ppm)	88.33 ± 3.53	90.04 ± 1.00	95.73 ± 2.95*

Note: Data presented are mean (n = 5) ± SD. * It differs significantly from other extracts ($\alpha = 0.05$).