

Putri Avidianto Excelinda - Utilization of Millet Seed Flour (*Panicum miliaceum* L), Chia Seed Flour (*Salvia hispanica*) and Sesame Seeds (*Sesamum indicum*) on Making High-Fiber Flakes

by Putri Avidianto Excelinda Uploaded By Arik Suprapti

Submission date: 18-Aug-2021 04:48PM (UTC+0700)

Submission ID: 1632783832

File name: A-057_PUTRI_AVIDAINTO_EXCELINDA_-_Putri_A_E.pdf (201.44K)

Word count: 5275

Character count: 25990

Utilization of Millet Seed Flour (*Panicum miliaceum* L), Chia Seed Flour (*Salvia hispanica*) and Sesame Seeds (*Sesamum indicum*) on Making High-Fiber Flakes

Keywords: chia seeds, dietary fiber, millet seeds, sesame seeds

Abstract: Millet seeds contain protein and fiber, so that it is potential to be used as raw materials for food. One of food that is very popular is flakes which can be made with blending millet seeds, chia seeds, and sesame seeds. These seeds will add nutritional value of the flakes. The purpose of this study was to produce high-fiber flakes that have a good sensory property and to identify the physical and chemical properties. The design of this study used a completely randomized design of one factor and two replications. The process of making flakes includes ingredients mixing, steaming, flattening, molding, and baking. The treatment factors in this study were flakes formulation with ratio of millet and chia seed flour (4 levels), namely F1 (90%:5%), F2 (85%:10%), F3 (80%:15%), F4 (75%:20%). Analysis of millet seed flour includes analysis of yield, proximate and dietary fiber. Organoleptic test includes hedonic test and hedonic quality test using the line scale method. Data analysis used the Kruskal wallis test, if the p-value ≤ 0.05 followed by the Man Whitney test. The results of this study showed that millet seed flour obtained water content of 16.3%, ash content of 2.17%, protein content of 9.74%, fat content of 0.11%, carbohydrate 71.7%, dietary fiber 12.5%, and 72.2% yield. The results of the organoleptic test showed that flakes were not significantly different from the hedonic test ($p > 0.05$), while they were significantly different to the hedonic quality of color and aroma ($p \leq 0.05$). The best formula is F4 with ash content of 3.07%, water content of 1.96%, protein content of 6.74%, fat content of 14.1%, carbohydrate content of 74.2%, dietary fiber content of 26.1%, and calories produced 450.7 kcal. The resulting flakes can be claimed as flakes with high dietary fiber.

1 INTRODUCTION

Proso millet (*Panicum miliaceum* L) or white millet is one of the small seed cereals that is usually for bird feed. The low use of millet seeds in food products is due to the lack of references and knowledge about the characteristics and content of millet seeds (Marta, 2016). The result of chemical analysis in research Prabowo (2010) nutrient content of white millet seed flour is 9.19% of water, 1.80% of ash, 2.58% of fat, 11.29% of protein, 56.53% of starch, 74.52% of carbohydrate, and 2.01% of crude fiber. According Kumar et al., (2018) that proso millet has a higher content of essential amino acids (leucine, isoleucine, and methionine) than wheat, and is rich in vitamins and minerals such as copper and magnesium, vitamin B6 and folic acid. India, Russia, Nigeria, and China used Millet seeds as a food source of carbohydrates (Amadou et al., 2014). The products made include bread, pasta, porridge, biscuits and drink fermentation, both of whole millet seeds or combined with other grains (Das et al., 2019). Several recent studies develop millet seeds in the manufacture of product such as instant baby porridge, dry noodles,

snack bar, and millet sprout flour (Husna et al., 2012; Atmaja and Sari, 2017; Adi Sarno et al., 2018; Dewi et al., 2018).

Chia seeds have been approved as a novel food by the European Parliament and of the Council (E.U Commission, 2015). Chia seeds are report be a safe as food because it has no side effects or allergen (EFSA, 2009). Chia seeds are a good source of omega-3 and omega-6 fatty acids. In addition, chia seeds have a high protein content (16-26%) and are rich in essential amino acids, especially leucine, lysine, isoleucine, and valine, as well as a source of dietary fiber (23-35%) and antioxidants (Vázquez-Ovando et al., 2010; Marcinek et al., 2017). Chia seeds have to used as a staple food by the Aztecs and Mayans (Muñoz et al., 2013). Recent research has shown that chia seeds can using to produce foods such as biscuits, pasta, and bread (Borneo, 2012; Oliveira et al., 2015; Romankiewicz et al., 2017).

Sesame seeds are known as the "Queen of oilseed" due to their high oil content and resistance to oxidation and rancidity (Myi, 2020). Traditionally, sesame seeds have to be used as a health food in Asian countries and is use to enhance taste and aroma of food, including as a topping for soups, salad, cereals

and yoghurt (Asghar et al., 2014). Sesame seeds contain large amounts of oil, protein, carbohydrates, essential mineral, methionine, and tryptophan, as well as secondary metabolites such as lignans, saponins, flavonoids, and phenolic compounds. In addition, as a source of calcium, phosphorus, and iron, and rich in vitamins B and vitamin E (Hegde, 2012).

Food sources of fiber are commonly found in plant foods such as cereals, fruits, vegetables, grains, and nuts. Based on Riskesdas (2018), it is stated that 95.5% of the population aged ≥ 5 years lack fiber consumption (vegetables and fruits). WHO recommends dietary fiber intake in the range of 25-30 g/person/day. Meanwhile, based on the RDA, in 2019 the average dietary fiber recommendation for Indonesians was 30 g/person/day. Dietary fiber is able to provide a feeling of fullness for longer, because foods with high fiber content take longer to digest so it is good as a food source at breakfast. One of the popular alternative breakfast products is flakes, especially for children (Aulia et al., 2017). The limited time in preparing breakfast made the interest in flakes increased. This makes flakes as one of the right choices for the ready to eat product category that is easy, fast, and practical to serve. Flakes are generally made from corn (corn flakes) and wheat (wheat flakes) (Situmorang et al., 2017). Commercial flakes have nutritional value per 100 g ranging 310-367 kcal of energy, 1.7 g of fat, 80-83 g carbohydrates, 3.3-17 g of fiber, and 6.7 -10.3 g of protein.

In this study, the millet seeds are selected as raw material for flakes to replace corn flour because the protein content of millet seed flour is higher (11.29 g/100g) than corn flour (8.78 g/100g), and product based on millet seed flour have a glycemic index content (50.2-64.7) which is lower than product from corn flour (78.5-86.3) (McSweeney, 2014). In addition, millet seed flour is gluten-free so it is safe for people has disability from celiac disease (gluten-intolerance) from which they cannot consume gluten-containing foods such as wheat (Sarita and Singh, 2016). Dietary fiber in millet seeds also has positive effects on health, such as anti-inflammatory, antidiabetic, anti-hypercholesterolemia, and antiradiation (Sulistyaningrum and Aqil, 2017).

In improving the nutritional value of flakes, required other ingredients besides flour millet seeds like chia seeds and sesame seeds which is a functional food ingredient with bioactive components. Chia seeds flour has a high protein content (26.2%), high dietary fiber (30.24%), and high levels of alpha-linolenic acid (68.0%) which are good for health (Ayza and Coates, 2011). Sesame seeds are high in

protein (19.3 g/100 g), rich in calcium (1125 mg/100 g) and iron (9.5 mg/100 g) (TKPI, 2017). The sesame seeds used in this study were roasted sesame seeds as a sprinkling on the flakes to produce a nutty aroma which is expected to increase acceptance of the product. Product flakes-based millet, chia, and sesame seeds are expected to be an alternative for nutritional needs at breakfast. In addition, efforts to create products with a low glycemic index, rich in fiber, and gluten-free.

2 MATERIAL AND METHODS

2.1 Material

Millet seeds, chia seeds, sesame seeds, tapioca flour, milk powder, sugar and salt were purchased commercially from local market.

2.2 Methods

2.2.1 Millet Seed Flour

Millet seeds were cleaned for extraneous materials, and then soaked in water for 24 hours. The water soaking needs to be replaced with clean water every 12 hours. After that, blanching for 2 minutes and continued drying in an oven at 100°C for 30 minutes. After obtaining the dry millet seeds, they were milled and sieved of 60 mesh with two repetitions in order to obtain more optimal millet seed flour.

2.2.2 Chia Seed Flour

The chia seeds are roasting for 6 minutes, stirring frequently. Then roasted chia seeds are milled using blander and sieved using a sieve of 60 mesh.

2.2.3 Sesame Seed Roasted

The sesame seeds in this study were not made into flour, but only through a roasting process. The sesame seeds are roasting for 5 minutes while stirring then removed.

2.2.4 Blend Formulation and Preparation of Flakes

The flakes formula used consisted of 4 levels of treatment using a ratio of millet seed flour to chia seed flour are F1 (90%:5%), F2 (85%:10%), F3 (80%:15%), dan F4 (75%:20%). Roasted sesame

seeds added a fixed amount of 10 grams of each formulation. The formulation for making these flakes refers to Khairunissa et al. (2018). The main ingredients were mixed according to the treatment and added 5 grams of tapioca flour as a binder for the dough. Then add 10 grams of powdered milk, 15 grams of sugar, and 1 gram of salt to give the flakes a taste. The entire dough is mixing with water as much as 45% of the total dough until completely homogeneous. Furthermore, the dough is steamed to help pre-gelatinize the starch for 10 minutes, in this process there is a change in the starch granules and the breakdown of hydrogen bonds that function to maintain the structure so that it cannot return to its original shape (Muflihani, 2016). Then the dough is flattened with a noodle maker on a scale of 3 and printed manually using a knife, then baked using an oven at 120°C for 10 minutes.

2.2.5 Physical Properties of Flour

Physical properties that can be seen is the yield of flour. Yield is the percentage of the main raw material that can be used as a final product. The purpose of the yield is to find out how much flour is produced from the raw materials used. The measurement of flour yield is calculated based on the ratio of the weight of flour obtained with the weight of the raw material expressed in percent (%).

2.2.6 Proximate Analysis of Ingredients and Flakes Samples

The protein, fat, ash, moisture content of the flour and flakes with the best formula determined by the SNI 01-2891-1992 proximate analysis. Carbohydrate content was estimate by difference and caloric value was calculated by using Atwater factor ($4 \times \text{protein}$, $9 \times \text{fat}$, and $4 \times \text{carbohydrate}$). Dietary fiber with AOAC 1990.9855.29 proximate analysis.

2.2.7 Sensory Evaluation

Sensory evaluation was carried out by 50 consumer panelists. Sensory analysis of flakes includes hedonic test and hedonic quality test. The hedonic test was carried out to determine the most preferred flakes formula by the panelists. The assessed attributes include color, texture, taste, aroma, and overall. The score scale used is a score of 1 (very dislike) to a score of 7 (very much like). The hedonic quality test is more specific, not just likes or dislikes. The score scale used to determine the color quality is a score of 1 (very not strong dark cream

color) to a score of 7 (very strong dark cream color), for the aroma used a score of 1 (very not strong nutty aroma) to 7 (very strong nutty aroma), for the taste used a score of 1 (not very strong sweet/savory taste) to 7 (very strong sweet/ savory taste), for the texture used a score of 1 (very not crispy) to 7 (very crispy).

2.2.8 Statistical Analysis

Data were analysed statistically by Analysis of Variance (ANOVA) when the data were normally distributed, and if there were significant differences then conducted a further test using Duncan's Multiple Range Test (DMRT). Meanwhile, data that were not normally distributed were analysed using the Kruskal Wallis test, and if there were significant differences, then carried out further using the Mann Whitney test.

3 RESULT AND DISCUSSION

3.1 Physical Properties of Flour

The yield of millet seed flour of 72.2% is not much different from Mahendra (2019) studies ranged between 65.27% - 84.60%. This is related to the processing method carried out such as the soaking process in this study which reduces water-soluble components such as potassium, sodium, and tannin in millet seeds, so the yield of millet seed flour produced is not much different from previous studies (Mahendra, 2019). The yield of chia seed flour produced was 85.0% is no different from Riemersman (2016) study of 94.0%. The high and low yields produced are due to the evaporation of substances contained in chia seeds during the roasting process, such as evaporation of water and the process of browning the seed (Purnamayanti et al., 2017).

3.2 Chemical Composition of Millet Flour

The chemical composition of millet flour is shown in table 1. Millet flour produced in this study has a moisture content of 16.3% was higher than millet flour produced at the research Prabowo (2010) and Dewi et al (2018) respectively by 9.19% and 8.35%. The protein content of millet seed flour produced was 9.74% lower than the research conducted by Prabowo (2010) at 11.29% but not much different from the research conducted by Dewi et al (2018) which was 9.77%. This decrease is due to protein denaturation

by high heating. The heating process can damage amino acids as protein-forming so that the heating temperature can reduce protein levels (Yuniarti et al., 2013). The fat content of millet seed flour produced was 0.11% lower than that of millet seed flour in Prabowo (2010) study of 2.58% and Dewi et al (2018) of 7.57%. The decrease in fat content was caused by the active lipase enzyme during the soaking process which was secreted by lactic acid bacteria. Lipase enzymes will hydrolyse fats into simpler compounds such as fatty acids and glycerol (Mahendra, 2019). According to Rani et al (2013) fat content can also decrease due to the soaking process followed by blanching. The dietary fiber content of millet seed flour produced was 12.5% higher than the research conducted by Dewi et al (2018) which was 5.93%. The dominant dietary fiber content in millet seeds is insoluble dietary fiber (Manish, 2018).

Table 1: Chemical composition of millet flour

Components	Millet Flour
Moisture (g/ 100 g)	16.3
Ash (g/ 100 g)	2.17
Protein (g/ 100 g)	9.74
Fat (g/ 100 g)	0.11
Carbohydrate (g/ 100 g)	71.7
Dietary fiber (g / 100 g)	12.5

3.3 Sensory Evaluation of Flakes

The hedonic characteristics of flakes made from millet seed flour, chia seed flour, and sesame seeds such as color, aroma, taste, texture and overall acceptability are shown in table 2. There were no significant differences in all sensory properties of flakes. The hedonic quality there are differences in color and aroma attributes, shown in table 8. The color in each formulation is influenced by the percentage of chia seed flour added to the flakes product. The higher the percentage of chia seed flour then the resulting color will be darker. This is also supported by the statement of Pizarro et al (2013) that the use of 0% to 30% chia seed flour will reduce the brightness value of pound cake. The aroma of flakes is obtained from millet seed flour which has a nutty aroma and a distinctive aroma of chia seed flour (Casper et al., 2014). Supported by research by

Shivakumar et al (2014) the addition of millet seed flour to cheese making gives a nutty aroma. Coorey et al (2012) the addition of 5% chia seed flour can affect the aroma of the chips. According to Hatamian et al (2020) the aroma of chia seed flour is obtained from the process of roasting the seeds before they become flour.

Table 2: Hedonic score of flakes

	Color	Aroma	Taste	Texture	Overall
F1	4.40± 1.29 ^a	4.62± 1.22 ^a	4.73± 1.31 ^a	5.17± 1.31 ^a	4.99± 1.22 ^a
F2	4.52± 1.11 ^a	4.28± 1.21 ^a	4.67± 1.21 ^a	4.90± 1.09 ^a	4.86± 1.18 ^a
F3	4.41± 1.27 ^a	4.45± 1.26 ^a	4.41± 1.33 ^a	4.57± 1.25 ^a	4.74± 1.36 ^a
F4	4.48± 1.46 ^a	4.40± 1.29 ^a	4.97± 1.14 ^a	4.97± 1.17 ^a	5.08± 1.11 ^a
LSD 5%	0.98	0.61	0.17	0.09	0.54

Table 3: Hedonic quality score of flakes

	Color	Aroma	Sweetness	Savoriness	Texture
F1	3.82 ± 1.21 ^a	3.56± 1.19 ^a	4.07± 0.98 ^a	3.93± 1.27 ^a	5.27± 1.27 ^a
F2	4.56 ± 0.97 ^b	4.22± 1.33 ^{bc}	3.94± 1.06 ^a	4.20± 1.12 ^a	5.11± 1.14 ^a
F3	4.73 ± 1.12 ^b	4.15± 1.29 ^b	3.78± 0.95 ^a	4.14± 1.01 ^a	5.13± 1.19 ^a
F4	5.06 ± 1.21 ^b	4.71± 1.14 ^{cd}	4.31± 1.03 ^a	4.36± 1.23 ^a	5.28± 1.20 ^a
LS D 5%	0.00	0.00	0.65	0.40	0.83

3.4 Chemical Composition of Flakes

The result of the hedonic and hedonic quality test obtained the selected formulations is F4 be obtained with the use of 75% millet seed flour and 20% chia seed flour. The nutritional content of selected flakes is shown in the table 4. The selected flakes will be compared with commercial flakes and the cereal quality requirements according to SNI No. 01-4270-1996 to determine the suitability of the nutritional quality of the selected formula.

Table 4: Analysis of selected flakes composition

Components	Selected Flakes	Commercial	SNI
Moisture (g/ 100 g)	1.96	-	Max.3
Ash (g/ 100 g)	3.07	-	Max.4
Protein (g/ 100 g)	6.74	6.7	Max.5
Fat (g/ 100 g)	14.1	1.7	Max.7
Carbohydrate (g/ 100 g)	74.2	80	Min.60
Dietary fiber (g / 100 g)	26.1	3.3	-
Energy (kcal)	450.7	366.7	-

Based on the results of the proximate analysis and dietary fiber, it showed that the fat content and dietary fiber content of the selected flakes were significantly higher than those of commercial flakes made from corn flour. This is due to the use of chia seed flour as much as 20% in selected flakes which is a source of fiber. Based on all the nutritional parameters of the selected flakes product, it has complied with SNI 01-4270-1996 for cereal quality. The fat content of the selected flakes is much higher. This can happen because of the oil content in chia seeds and sesame seeds. The fat content of chia seed flour is 32.59% (Arumsari dan Sofyaningsih, 2020). Research conducted by Rendón-Villalobos *et al.* (2012) stated that the manufacture of corn tortillas with the addition of 20% chia seed flour resulted in a much higher fat content (10.95%) than the control (4.08%) which could be attributed to concentration of chia seed oil. According to Mohd Ali *et al.*, 2012 chia seeds contain 25-40% fat, which is in the form of polyunsaturated fatty acids such as omega-3 (alpha linolenic acid / ALA) and omega-6 (linoleic acid). The dominant fatty acid content in sesame seeds in the form of oleic and linoleic acids ranges from 80-85% of the total amount (Mahmood Biglar, 2012). These fatty acids are included in the essential fatty acids needed by health.

These selected flakes have fulfilled nutritional claims as high-fiber foods because they contain more than 6 g per 100 g solid weight. The high content of dietary fiber in the selected flakes was caused by several ingredients such as chia seed flour and millet seed flour which contain high dietary fiber. In this study, the dietary fiber content of millet seed flour reached 12.5 g per 100 g. While the content of dietary fiber in chia seed flour is 30.24 g per 100 g (Arumsari dan Sofyaningsih, 2020). Increased dietary fiber can also result from the development of resistant starch

during the heating and cooling process (Takhellambam *et al.*, 2016). The dietary fiber content of millet seeds and chia seeds is included in the insoluble dietary fiber (Rana, 2019). Insoluble fiber includes lignin, cellulose, and hemicellulose. The main component found in insoluble fiber is lignin, which plays an important role in the protection of unsaturated fats and is responsible for the hypercholesterolemic activity associated with fiber intake (Tolba *et al.*, 2011). Chia seed flour has a much higher fiber content, so it will determine the percentage of fiber content of the flakes.

3.6 Nutrition Facts of Flakes

Determination of the serving size of flakes is adjusted to the suggestion of serving commercial products from corn flakes is 30 g and served with milk. The energy and nutritional contributions of flakes based on millet seed flour, chia seed flour and sesame seeds are shown in table 5.

Table 5: Nutrition facts of flakes

6	Nutrition Facts		
serving size 1 cup (30g)			
Servings Per Container About 1			
Amount Per Serving		Flakes	Flakes with 200ml Milk
Calories		135	257
Calorie from Fat		36	99
			% Daily Value
Total Fat	4 g	6%	17%
Total Carbohydrate	22 g	7%	10%
Dietary Fiber	8 g	27%	27%
Protein	2 g	3%	14%

4 CONCLUSIONS

The results showed that the best formulation of flakes from millet seed flour, chia seed flour, and sesame seeds based on the organoleptic test is F4 (75% millet seed flour: 20% chia seed flour). The results of the sensory test analysis in the study of the comparison of millet seed flour and chia seed flour with hedonic parameters of color, aroma, taste, texture and overall showed that there was no significant difference. Meanwhile, the hedonic quality was significantly different to the color and aroma quality of the flakes. Analysis of the nutritional content of the best flakes formula that is ash content of 3.07%, water content of 1.96%, protein content of

6.74%, fat content of 14.1%, carbohydrate content of 74.2%, dietary fiber content of 26.1%, and calories produced 450.7 kcal per 100 grams. The resulting flakes can be claimed as flakes with high dietary fiber.

REFERENCES

- Adi Sarno, I. P., Wulandari, Y. W., & Suhartatik, N. (2018). Karakteristik Snack Bars dengan Variasi Suhu Pemanggangan dan Perbandingan Tepung Millet Kuning (*Panicum sp*) dengan Tepung Pisang Raja Bandung (*Musa paradisiaca* L.). *Jurnal Teknologi Pangan*, 12(2), 47–53. <https://doi.org/10.33005/Jtp.V12i2.1288>
- Amadou, I., Gounga, M. E., Shi, Y. H., & Le, G. W. (2014). Fermentation and Heat-Moisture Treatment Induced Changes on The Physicochemical Properties of Foxtail Millet (*Setaria italica*) Flour. *Food and Bioproducts Processing*, 92(1), 38–45. <https://doi.org/10.1016/j.fbp.2013.07.009>
- Arumsari, I., & Sofyaningsih, M. (2020). Evaluasi Zat Gizi Tepung Chia (*Salvia hispanica* L.) dan Tepung Wijen (*Sesamum indicum* L.) sebagai Alternatif Tepung Tinggi Serat dan Protein. 5(1), 27–33. <https://doi.org/10.22236/Argipa.V5i1.4950>
- Asghar, A., Majeed, M. N., & Akhtar, M. N. (2014). A Review on The Utilization of Sesame as Functional Food. *Am. J. Food. Nutr.*, 4(1), 21–34. <https://doi.org/10.5251/Ajfn.2014.4.1.21.34>
- Atmaja, R. P., & Sari, R. Y. (2017). Pembuatan Tepung Millet Terfermentasi dan Pemanfaatannya dalam Produk Mie Kering Fermentation. *Journal of Chemical Information And Modeling*, 53(9), 1689–1699. <https://doi.org/10.1017/Cbo9781107415324.004>
- Aulia, T., Suhaedi, I., & Rusmarilin, H. (2017). Tepung Pisang dan Persentase Kuning Telur terhadap Mutu Flakes Talas. *Ilmu dan Teknologi Pangan*, 5(2), 333–342.
- Borneo, R. (2012). Chia (*Salvia hispanica*) Can be Used to Manufacture Sugar-Snap Cookies with an Improved Nutritional Value. *International Journal of Food Studies*, 1(2), 135–143. <https://doi.org/10.7455/Ijfs/1.2.2012.A4>
- Casper, J. L., Cargill, I., Plymouth, M., Atwell, W. A., & Champlin, M. (2014). Gluten-Free Ingredients. In *Gluten-Free Baked Products* (Pp. 23–47). <https://doi.org/10.1016/B978-1-891127-80-9.50003-0>
- Coorey, R., Grant, A., & Jayasena, V. (2012). Effects of Chia Flour Incorporation on The Nutritive Quality and Consumer Acceptance of Chips. *Journal of Food Research*, 1(4), 85. <https://doi.org/10.5539/Jfr.V1n4p85>
- Das, S., Khound, R., Santra, M., & Santra, D. K. (2019). Beyond Bird Feed: Proso Millet For Human Health and Environment. *Agriculture (Switzerland)*, 9(3). <https://doi.org/10.3390/Agriculture9030064>
- Dewi, I. S., Ekawati, I., & Pratiwi, I. (2018). Pengaruh Lama Perkecambahan Millet (*Panicum milliaceum*) Terhadap Karakteristik Flakes. *Jurnal Ilmu Dan Teknologi Pangan*, 7(4), 175–183.
- Hatamian, M., Noshad, M., Abdanan-Mehdizadeh, S., & Barzegar, H. (2020). Effect of Roasting Treatment on Functional and Antioxidant Properties of Chia Seed Flours. *Nfs Journal*, 21(March), 1–8. <https://doi.org/10.1016/J.Nfs.2020.07.004>
- Hegde, D. M. (2012). *Sesame. Handbook Of Herbs and Spices: Second Edition*, 2, 449–486. <https://doi.org/10.1533/9780857095688.449>
- Husna, E. A., Affandi, D. R., & Anandito, R. B. K. (2012). Karakterisasi Bubur Bayi Instan Berbahan Dasar Tepung Millet (*Panicum sp*) dan Tepung Kacang Hijau (*Phaseolus radiatus*) Dengan Flavor Alami Pisang Ambon (*Musa paradisiaca* Var. *Sapientum* L.). 1(1).
- Khairunissa, Harun, N., & Rahmayuni. (2018). Pemanfaatan Tepung Talas dan Tepung Kacang Hijau dalam Pembuatan Flakes [Utilization Of Taro Flour and Mung Bean Flour in Making Flakes]. *Jurnal Sagu Universitas Riau*, 17(1), 2018.
- Kumar, A., Tomer, V., Kaur, A., Kumar, V., & Gupta, K. (2018). Millets: A Solution to Agrarian and Nutritional Challenges. *Agriculture and Food Security*, 7(1), 1–15. <https://doi.org/10.1186/S40066-018-0183-3>
- Pizarro, P. L., Almeida, E. L., Sammán, N. C., & Chang, Y. K. (2013). Evaluation of Whole Chia (*Salvia hispanica* L.) Flour and Hydrogenated Vegetable Fat In Pound Cake. *Lwt - Food Science and Technology*, 54(1), 73–79. <https://doi.org/10.1016/J.Lwt.2013.04.017>
- Mahendra, P. E. D., Yusasrini, N. L. A., & Pratiwi, I. D. P. K. (2019). Pengaruh Metode Pengolahan Terhadap Kandungan Tanin dan Sifat Fungsional Tepung Proso Millet (*Panicum miliaceum*). *Jurnal Ilmu dan Teknologi Pangan (Itepa)*, 8(4), 354. <https://doi.org/10.24843/Itepa.2019.V08.I04.P02>
- Mahmood Biglar. (2012). Profiling of Major Fatty Acids In Different Raw and Roasted Sesame Seeds Cultivars. *African Journal Of Biotechnology*, 11(24), 6619–6623. <https://doi.org/10.5897/Ajb11.1966>
- Manish, C. El. (2018). Nutritional and Nutraceutical Properties of Millets: Clinical Journal of Nutrition And Dietetics, 1(1), 1–10.
- Marcinek, K., & Krejpcio, Z. (2017). Chia Seeds (*Salvia hispanica*): Health Promoting Properties and Therapeutic Applications – A Review. *Roczniki Panstwowego Zakladu Higieny*, 68(2), 123–129.
- Marta, H. (2016). Sifat Fungsional dan Amilografi Pati Millet Putih (*Pennisetum glaucum*) Termodifikasi Secara Heat Moisture Treatment dan Annealing. *Jurnal Aplikasi Teknologi Pangan*, 5(3), 76–84. <https://doi.org/10.17728/Jatp.175>
- Mcsweeney, M. (2014). Proso Millet As An Ingredient In Foods Common To North Americans. (October).
- Mohd Ali, N., Yeap, S. K., Ho, W. Y., Beh, B. K., Tan, S.

- W., & Tan, S. G. (2012). The Promising Future of Chia, *Salvia Hispanica* L. *Journal of Biomedicine And Biotechnology*, 2012. <https://doi.org/10.1155/2012/171956>
- Muflihani, Y. (2016). Uji Organoleptik Formula Flakes Dari Pasta Ubi Jalar Dengan Penambahan Tepung Jalejo. *Prosiding Seminar Hasil Penelitian Tanaman Aneka Kacang dan Umbi.*, (Widowati 2011), 603–610.
- Muñoz, L. A., Cobos, A., Diaz, O., & Aguilera, J. M. (2013). Chia Seed (*Salvia Hispanica*): An Ancient Grain and A New Functional Food. *Food Reviews International*, 29(4), 394–408. <https://doi.org/10.1080/87559129.2013.818014>
- Myint, D., Gilani, S. A., Kawase, M., & Watanabe, K. N. (2020). Sustainable Sesame (*Sesamum indicum* L.) Production Through Improved Technology: An Overview Of Production, Challenges, and Opportunities In Myanmar. *Sustainability (Switzerland)*, 12(9), 1–21. <https://doi.org/10.3390/Su12093515>
- Oliveira, M. R., Novack, M. E., Santos, C. P., Kubota, E., & Da Rosa, C. S. (2015). Evaluation of Replacing Wheat Flour with Chia Flour (*Salvia hispanica* L.) In Pasta. *Semina: Ciencias Agrarias*, 36(4), 2545–2553. <https://doi.org/10.5433/1679-0359.2015v36n4p2545>
- Prabowo, B. (2010). Kajian Sifat Fisikokimia Tepung Millet Kuning Dan Tepung Millet Merah. *Skripsi*, 1–39.
- Purnamayanti, N. P. A., Gunadnya, I. B. P., & Arda, G. (2017). Pengaruh Suhu dan Lama Penyangraian Terhadap Karakteristik Fisik dan Mutu Sensori Kopi Arabika (*Coffea Arabica* L.). *Jurnal Beta (Biosistem Dan Teknik Pertanian)*, 5(2), 39–48.
- Rana. (2019). Characterization Of Chia Seed Flour and Wellbeing Endorsing Possessions. *International Journal Of Food Science, Nutrition And Dietetics*, 8(5), 419–426. <https://doi.org/10.19070/2326-3350-1900075>
- Rani, H., Zulfahmi, & Widodo, Y. R. (2013). Optimasi Proses Pembuatan Bubuk (Tepung) Kedelai Optimization Process Soybean Flouring. *Jurnal Penelitian Pertanian Terapan*, 13(3), 188–196.
- Rendón-Villalobos, R., Ortiz-Sánchez, A., Solorza-Feria, J., & Trujillo-Hernández, C. A. (2012). Formulation, Physicochemical, Nutritional and Sensorial Evaluation of Corn Tortillas Supplemented with Chia Seed (*Salvia hispanica* L.). *Czech Journal Of Food Sciences*, 30(2), 118–125. <https://doi.org/10.17221/393/2010-Cjfs>
- Riemersman, C. N., & Maria, R. A. (2016). Whole Chia Flour As Yield Enhancer, Potential Antioxidant and Input Of N-3 Fatty Acid In A Meat Product. *Food And Nutrition Sciences*, 07(10), 855–865. <https://doi.org/10.4236/Fns.2016.710085>
- Romankiewicz, D., Hassoon, W. H., Cacak-Pietrzak, G., Sobczyk, M. B., Wirkowska-Wojdyba, M., Ceglinska, A., & Dziki, D. (2017). The Effect of Chia Seeds (*Salvia hispanica* L.) Addition on Quality and Nutritional Value of Wheat Bread. *Journal Of Food Quality*, 2017. <https://doi.org/10.1155/2017/7352631>
- Sarita, & Singh, E. (2016). The Pharma Innovation Journal 2016; 5(8): 42–46 Millet's Anti-Nutrients and Their Therapeutic Effects. 5(8), 42–46. Retrieved From <http://www.slideshare.net/Poshadri/Cereals-Millet-Processing>.
- Shivakumar, Arunkumar.H., & Venkatesh.M.V. (2014). Process Optimization For The Production of Paneer (Soft Cheese) Kheer Blended with Foxtail Millet and Finger Millet Flour. *Quest Journals Journal Of Research In Agriculture And Animal Science*, 2(6), 2321–9459. Retrieved From www.questjournals.org
- Situmorang, C., Swamilaksita, D. P., Anugrah, N., Gizi, P. I., Kesehatan, F. I., & Unggul, U. E. (2017). Substitusi Tepung Kacang Hijau dan Tepung Kacang Kedelai pada Pembuatan Bean Flakes Tinggi Serat dan Tinggi Protein Sebagai Sarapan Sehat.
- Sulistyaningrum, A., & Aqil, M. (2017). Karakteristik Tepung Jewawut (Foxtail Millet) Varietas Lokal Majane dengan Perlakuan Perendaman. 11–21.
- Zulfianto. N. A. (2017). Tabel Komposisi Pangan Indonesia. Jakarta: Kementerian Kesehatan RI.
- Takhellambam, R. D., Chimmad, B. V., & Prkasam, J. N. (2016). Ready-To-Cook Millet Flakes Based on Minor Millets For Modern Consumer. *Journal of Food Science and Technology*, 53(2), 1312–1318. <https://doi.org/10.1007/S13197-015-2072-0>
- Tolba, R., Wu, G., & Chen, A. (2011). Adsorption Of Dietary Oils Onto Lignin for Promising Pharmaceutical and Nutritional Applications. *Bioresources*, 6(2), 1322–1335. <https://doi.org/10.15376/Biores.6.2.1322-1335>
- Vázquez-Ovando, J. A., Rosado-Rubio, J. G., Chel-Guerrero, L. A., & Betancur-Ancona, D. A. (2010). Dry Processing of Chia (*Salvia hispanica* L.) Flour: Chemical Characterization of Fiber and Protein Article. *Cyta - Journal Of Food*, 8(2), 117–127. <https://doi.org/10.1080/19476330903223580>
- Yuniarti, D. W., Sulistiyati, T. D., & Suprayitno, E. (2013). Pengaruh Suhu Pengeringan Vakum terhadap Kualitas Serbuk Albumin Ikan Gabus (*Ophiocephalus striatus*). *Thpi Student Journal*, 1.

Putri Avidianto Excelinda - Utilization of Millet Seed Flour (*Panicum miliaceum* L), Chia Seed Flour (*Salvia hispanica*) and Sesame Seeds (*Sesamum indicum*) on Making High-Fiber Flakes

ORIGINALITY REPORT

9%

SIMILARITY INDEX

8%

INTERNET SOURCES

5%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1	www.hindawi.com Internet Source	2%
2	www.mdpi.com Internet Source	1%
3	link.springer.com Internet Source	1%
4	mafiadoc.com Internet Source	1%
5	Nutrition & Food Science, Volume 43, Issue 5 (2013-09-28) Publication	1%
6	www.arrowheadmills.com Internet Source	1%
7	"Oilseeds: Health Attributes and Food Applications", Springer Science and Business Media LLC, 2021 Publication	1%

8	Viviane da Costa Borges, Sibeles Santos Fernandes, Elessandra da Rosa Zavareze, Claudia Monika Haros et al. "Production of gluten free bread with flour and chia seeds (Salvia hispânica L)", Food Bioscience, 2021 Publication	1 %
9	repository.uin-suska.ac.id Internet Source	1 %
10	jurnal.uns.ac.id Internet Source	<1 %
11	chimie-biologie.ubm.ro Internet Source	<1 %
12	www.ojafr.ir Internet Source	<1 %
13	www.scidoc.org Internet Source	<1 %

Exclude quotes

Off

Exclude matches

< 17 words

Exclude bibliography

On