

Jurnal JYoungPharm_10_2_s93

by Yesi .

Submission date: 21-Apr-2020 08:46AM (UTC+0700)

Submission ID: 1303232394

File name: Jurnal_JYoungPharm_10_2_s93.pdf (211.27K)

Word count: 3063

Character count: 16865

Antioxidant Activity of *Rubus fraxinifolius* Poir. and *Rubus rosifolius* J. Sm. Leaves

Yesi Desmiaty^{1,2}, Berna Elya^{1*}, Fadlina Chany Saputri¹, Muhammad Hanafi^{2,3}, Rini Prastiwi⁴

¹Faculty of Pharmacy, Universitas Indonesia, Depok, INDONESIA.

²Faculty of Pharmacy, Pancasila University, Jakarta, INDONESIA.

³Research Centre for Chemistry Indonesian Institute of Sciences, Jakarta, INDONESIA.

⁴Faculty of Pharmacy, Muhammadiyah Prof. Dr. Hamka University, Jakarta, INDONESIA.

ABSTRACT

Objective: *Rubus fraxinifolius* and *R. rosifolius* are found growing in mountainous areas of West Java and potentially act as an antioxidant. The objective of this research was to examine the antioxidant activity of *Rubus rosifolius* J. Sm. and *R. fraxinifolius* Poir. leaves. **Methods:** Dried leaves powder were extracted using Soxhlet apparatus with n-hexane, ethyl acetate, and methanol. The extracts were evaporated, and antioxidant activity was determined using DPPH free radical scavenger methods as well as FRAP reduction capability. Total phenolic contents from the extract gave the best antioxidant activity was measured by the Folin-Ciocalteu method. **Results:** IC₅₀ of DPPH scavenging activity of n-hexane, ethyl acetate and methanolic extracts of *R. fraxinifolius* were >200; 140.97; and 4.48 ppm, while for *R. rosifolius* were >200; 29.67; and 5.17 ppm, respectively. Percent FRAP capacity of *R. fraxinifolius* in 25 ppm were 55.65; 89.47; and 97.65 %, while for *R. rosifolius* were 73.51; 82.83; and 96.5, respectively. Total phenolic content of the methanolic extract of *R. fraxinifolius* and *R. rosifolius* were 39.0 + 26.5 and 80.62+21.6 mg GAE/g extract.

Conclusion: Both methanolic extracts showed a significant optimum trapping capability of free radicals and the hexane extracts did not show antioxidant activity. Thus, it may be concluded that the methanolic extracts of both leaves possess potent antioxidant properties and had high total phenolic content. This investigation provides promising results to emphasize the importance of antioxidant capabilities of both plants.

Key words: Antioxidant, *Rubus fraxinifolius*, *Rubus rosifolius*, Total phenolic content.

Correspondence

Berna Elya, Faculty of Pharmacy, Universitas Indonesia, Depok, INDONESIA.

Phone: +62-21-7270031

Email: berna.elya@ui.ac.id, icapps2017.farmasiui@gmail.com

DOI: 10.5530/jyp.2018.2s.18

INTRODUCTION

Indonesia is a tropical country and has numerous mountains with thousands of plants unexplored including chemical and biological activities. One of the genus that lives in the highlands of Indonesia is *Rubus* (Rosaceae). From the literature review, the collection of herbarium and exploration life collection of Indonesian Institute of Sciences (LIPI), it is known that there are 25 species of *Rubus* genus scattered across Indonesian mountain forests, including *R. alpestris*, *R. chrysophyllus*, *R. ellipticus*, *R. fraxinifolius*, *R. lineatus*, *R. moluccanus*, *R. pyrifolius*, dan *R. rosifolius*.¹ Fruit of *R. fraxinifolius* Poir. and *R. rosifolius* J. Sm. in the West Java area (Sundanese) are known as "beberetean" or "arben".^{2,3} Both fruits are edible, has similarity in shape which is small, have a red color, and have a sweet rather than sour taste. *Rubus* genus is a part of berries plants, which are small, soft, commonly edible, colorful and have important nutritional content.⁴

Some *Rubus* were reported to have a potential activity of antioxidant, anti-bacteria, anti-elastase, anti-collagenase, anti-thrombotic, and potential for the treatment of radical generated disorders, mainly cancer, and other inflammatory diseases as well as a large content of polyphenols and flavonoids.⁵⁻¹³ *R. fraxinifolius* Poir. had economic value due to its ability to produce fruit throughout the year.¹⁴ Leaves of *R. rosifolius* J. Sm. were reported to have 5,7-dihydroxy-6,8,4'-trimethoxyflavonol, 5-hydroxy-3,6,7,8,4'-pentamethoxyflavone, and tormentic acid which had anti-proliferative activity against ovary cancer cell.¹⁵ The main isolated compound from hexane extract of *R. rosifolius* herb was 28-methoxy-tormentic acid and was reported to have a potential analgesic activity.¹⁶

Fruit of *R. fraxinifolius* Poir. was reported have good antioxidant activity (FRAP method)¹⁷ but only few studies had established this result. It has been reported in *Rubus* species that there are phenolic compounds such as ellagic acid (usually found as glycosylated glycosylation polymers), gallic acid, chlorogenic acid, and caffeic acid.¹⁷

This study was aimed to examine the antioxidant capacity of both leaves extract using DPPH scavenging and FRAP methods. DPPH can only be dissolved in organic media; this condition becomes an important limitation when it comes to interpreting the role of hydrophilic antioxidants. The ferric ion reducing antioxidant power (FRAP) method is based on the reduction of a ferroin analog, the Fe³⁺ complex of tripyridyltriazine Fe(TPTZ)³⁺ to the intensely blue colored Fe²⁺ complex Fe(TPTZ)²⁺ by antioxidants in low pH medium. Both methods can be measured the product using spectrophotometry instrument.

MATERIALS AND METHODS

Chemical and reagents

Diphenyl-1-picrylhydrazyl (DPPH), Folin-Ciocalteu's phenol reagent, 2,4,6-tri(2-pyridyl)-s-triazine (TPTZ), gallic acid were purchased from Sigma-Aldrich®, Methanol, HCl, FeCl₃·6H₂O, aqua demineralisata.

Plant Material

Rubus rosifolius leave were collected from Jayagiri, Mount Tangkuban Perahu and *Rubus fraxinifolius* leaves were collected from Cibodas,

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Mount Gede-Panglango National Park in November 2016. Both plants was identified by Research Center for Biology, Indonesian Institute of Sciences (LPI), Cibinong, Indonesia. Fresh samples were cleaned, air dried and grounded into a fine powder by laboratory mill.

Extraction

Each dried powdered of leaves (20g) were extracted using Soxhlet apparatus by n-hexane, ethyl acetate, and methanol. The organic solvents were evaporated using rotary vacuum evaporator, and then were dried using vacuum oven.

Determination of Total Antioxidant Capacity

The DPPH radical-scavenging activity was determined using the method proposed by Blois.¹⁸ DPPH was dissolved in pure methanol (0.1 mM). The DPPH solution (180 µL ml) was added to 20 µL of extracts in methanol. The mixture was shaken vigorously and allowed to stand at room temperature in the dark for 30 min. The decrease in absorbance of the reaction mixture was monitored at 517 nm at 96-well microplate reader (Versamac).¹⁹ Determinations were performed in triplicate. The ability to scavenge DPPH radical was calculated by the following equation: DPPH radical scavenging activity (%) = $\frac{[(\text{Abs control} - \text{Abs sample}) / (\text{Abs control})] \times 100}{1}$ where Abs control is the absorbance of DPPH radical and methanol; Abs sample is the absorbance of DPPH radical + extract/control.⁸

Ferric Reducing Antioxidant Potential (FRAP)

FRAP reagent was prepared by mixing acetate buffer (300 mM, pH 3.6), a solution of 10 mM TPTZ in 40 mM HCl, and 20 mM FeCl₃ (10:1:1). The reagent (270 µl) and sample solutions (30 µl) were added to each well and mixed thoroughly. The absorbance was taken at 593 nm after 30 min 96-well microplate reader (Versamac). All solutions were used on the same day of preparation. All determinations were performed in triplicates.¹⁹ The antioxidant capacity was calculated using the formula FRAP capacity (%) = $\frac{[(\text{Abs sample} - \text{Abs control}) / \text{Abs sample}] \times 100}{1}$, where FRAP reagent was used as a control.

Total Phenolic Content Assay

Total phenolic content of both methanol extract was assessed by using a modified Folin Ciocalteu method (Singleton and Rossi, 1965). Folin Ciocalteu reagent (FCR) is a mixture of phosphotungstate and phosphomolybdate, which oxidizes the phenolates to a blue complex. Diluted sample (20 µL) in 95% ethanol and 100µl FCR (1:4 in aquadest) were mixed in a 96-well plate, shake in 5 min, then add 75 µl of sodium carbonate (100 g/L). After a 120min incubation period at room temperature, the absorbance of the reaction mixture was measured at 765 nm with a microplate spectrophotometer (Versamac). The gallic acid was used as a reference (10-200 mg/L). All samples were performed in triplicate. The results were expressed as milligrams of gallic acid equivalent (GAE) per gram of the extract.

RESULTS

Extraction

The dried extracts from both *Rubus* leaves were weighed and the yield of each extract was calculated. The extraction rendement of n-hexane, ethyl acetate and methanolic extracts of *R. fraxinifolius* were 1.76%; 5.85%; and 6.22%, while for *R. rosifolius* were 2.11%; 6.93%; and 6.32%, respectively.

Determination of Total Antioxidant Capacity

The DPPH scavenging activity of selected *Rubus* species and standard (Vitamin C) are depicted in Table 1. The result showed IC₅₀ of DPPH scavenging activity of n-hexane, ethyl acetate and methanolic extracts of

Table 1: Results of antioxidant activity testing of DPPH free radical scavenger

DPPH test	Fraction	IC ₅₀ (ppm)
<i>Rubus fraxinifolius</i> leaves	n-hexane	>200 ppm
	Ethyl acetate	140.97
	methanol	4.48
<i>Rubus rosifolius</i> leaves	n-hexane	>200 ppm
	Ethyl acetate	29.67
	methanol	5.17
Vitamin C		2.24




Figure 1: FRAP Capacity of Leaves Extract (25 ppm).




Figure 2: Total Phenolic Content of Methanolic Leaves Extract.

R. fraxinifolius were >200; 140.97; and 4.48 ppm, while for *R. rosifolius* were >200; 29.67; and 5.17 ppm, respectively. Both n-hexane fraction gave no activity with IC₅₀ more than 200ppm, and methanol fraction of both leaves had strong activity to scavenging free radical DPPH. Vitamin C which serves as positive control has IC₅₀ 2.24 ppm.

Ferric Reducing Antioxidant Potential (FRAP)

The FRAP capacity of both *Rubus* species are given in Figure 1. Percent FRAP capacity of *R. fraxinifolius* in 25 ppm were 55.65%; 89.47%; and 97.65%, while for *R. rosifolius* were 73.51%; 82.83%; and 96.50%, respectively. All fraction showed potential activity to reduction of the complex of ferric ion to the ferrous ion and correlated with antioxidant activity which involves Single Electron Transfer (SET) mechanism. All experi-

ments were triplicated and analyzed using ANOVA two-factor with replication using Microsoft Excel 2016 showed significant differences ($p < 0.05$) in the FRAP capacity between extracts and both species.

17 Total Phenolic Content Assay

The total phenolic content of both methanolic leaves extracts are shown in Figure 2. Total phenolic content of the methanolic extract of *R. fraxinifolius* and *R. rosifolius* were 3.14 ± 26.5 mg GAE/g extract and 80.62 ± 21.6 mg GAE/g extract. All results are presented as mean \pm standard deviation (SD) of three replicates and were calculated using Microsoft Excel 2016.

DISCUSSION

The antioxidant activities of both *Rubus* species were investigated using two different *in vitro* antioxidant tests. DPPH (α -diphenyl- β -picrylhydrazyl) free radical scavenging method developed by Blois offers the first approach for evaluating the antioxidant potential of a compound, an extract or other biological sources. This method is the simplest, in which the compound or extract is mixed with DPPH solution and the absorbance is recorded after a certain period of time. (DPPH=C₁₈H₁₂N₅O₆, M=394.33).²⁰ The antioxidant capacity of extracts on DPPH free radical scavenging was due to its hydrogen-donating process. In the DPPH test, the purple color of DPPH solution turns yellow in the presence of antioxidant compounds.

FRAP method developed by Benzie and Strain, measured the antioxidants capacity in reducing ferric (Fe³⁺). FRAP method was used to determine the antioxidant activity which involves Single Electron Transfer (SET) mechanism. It is based on the reduction of the complex of ferric ion to the ferrous ion (Fe²⁺) at acidic condition. The latter form a blue complex (Fe²⁺/TPTZ (2,3,5-triphenyl-1,3,4-triazza-2-azoniacyclopenta-1,4-dienechloride)), which monitored by measuring the change in absorption at 593 nm, using a microplate reader.²¹ The FRAP assay is cheap, reagents are easy to prepare, reproducible, and the procedure is simple and quick so it can be done in every laboratory and researcher who interested in antioxidant assay. Based on the result showed that all fraction of both leaves gave potential FRAP capacity. Higher FRAP values give higher antioxidant capacity because FRAP value is based on reducing ferric ion, where antioxidants are the reducing agent. An antioxidant is a compound that can contribute an electron or a hydrogen atom to reduce the reacting radical.

Both methanol fractions of *R. rosifolius* and *R. fraxinifolius* showed a significant optimum trapping capability of free radicals. However, the hexane fractions did not show any antioxidant activity. In the methanol extract contains polar compounds such as phenolic compounds. These phenolic compounds are known to have antioxidant activity, especially phenolic acids and flavonoids. Phenolic acid is the main class of phenolic compounds, widely found in plants especially in fruits and vegetables.¹⁹ It has been reported in *Rubus* species that there are phenolic compounds such as ellagic acid (usually found as glycosylated glycosylation polymers), gallic acid, chlorogenic acid, and caffeic acid.¹⁷

Polyphenolic compounds are known to react strongly in scavenge free radicals by donating hydrogen atoms or electron chelating metal ions. The use of FCR is the most commonly used method for assessing TPC. In fact, this method may give a false positive reaction and do not provide the actual measurement of phenolic compounds due to the presence of reducing compounds such as ascorbic acid and other reducing sugars, which can also react with the FCR. Various modification attempts have been made to improve the specificity of the FCR based test originally developed by Singelton and Rossi (1965). The total phenolic content of methanolic extracts varied and methanol leaf extract of *R. rosifolius* showed higher phenolic content than *R. fraxinifolius*. The Pearson

correlation coefficient (r) was calculated to show correlation between TPC and FRAP activity, it was considered statistically extremely significant with $r=0.9939$.

CONCLUSION

Thus, it may be concluded that both methanol extracts showed better radical scavenging activity than other selected solvents and had high total phenolic content. This investigation gave the promising result to emphasize the importance of antioxidant activities of both leaves.

11 ACKNOWLEDGEMENT

This work was funded by Universitas Indonesia for PITTA grant in 2017.

10 CONFLICT OF INTEREST

The authors declare no conflict of interest.

ABBREVIATIONS

DPPH: 2,2- Diphenyl-1-picrylhydrazyl; **TPTZ:** 2,3,5-triphenyl-1,3,4-triazza-2-azoniacyclopenta- 1,4-dienechloride; **GAE:** Gallic Acid Equivalent; **FRAP:** Ferric Ion Reducing Antioxidant Power.

REFERENCES

- Surya M, Ismaini L, Destri. Keragaman Buah Raspberries (*Rubus* spp.) Asal Indonesia. In: Seminar nasional Biologi. Semarang: Jurusan Biologi FMIPA Univ. Negeri Semarang. 2015;296-305.
- Zandee M, Kalkman Lab Exp Plant Taxonomy C, Rijksherbarium Leiden L. The genus *Rubus* (Rosaceae) in Malaysia. 1. Subgenera *Chamaebatus* and *Idaebatus*. BLUMEA27 (I9B1). 1981;27(1):75-113.
- Kalkman C, Verheij EW, Coronel R. *Rubus*. PROSEA (Plant Resources of South-East Asia) Foundation. 1991. [cited 2016 Dec 8]. Available from: http://proseanet.org/prosea/e-prosea_detail.php?frt=e&id=8
- Padmanabhan P, Palyath G. Berries and Related Fruits. 1st ed. Vol. 1, Encyclopedia of Food and Health. Canada: Elsevier Ltd. 2016;1:364-71.
- Park CM, Joung MS, Yang DO, Choi JW. The Study on the Effect of *Rubus coreanus* Miquel Extract as a Cosmetic Ingredient. J Soc Sci Korea. 2007;33(1):41-5.
- Bravo K, Alzate F, Osorio E. Fruits of selected wild and cultivated Andean plants as sources of potential compounds with antioxidant and anti-aging activity. Ind Crops Prod. 2016;85:341-52.
- Lino M. Elastase inhibitor raspberry. South Korea; Patent US8491946 B2. 2010. [cited 2016 Dec 1]. Available from: <https://www.google.com/patents/US8491946>
- Barcelo R. Phytochemical Screening and Antioxidant Activity of Edible Wild Fruits in Benguet, Cordillera Administrative Region, Philippines. Electron J Biol. 2015;11(3):80-9.
- Martini S, D'Addario C, Colacevich A, Focardi S, Borghini F, Santucci A, et al. Antimicrobial activity against *Helicobacter pylori* strains and antioxidant properties of blackberry leaves (*Rubus ulmifolius*) and isolated compounds. Int J Antimicrob Agents. 2009;34(1):50-9.
- Xie P, Zhang Y, Wang X, Wei J, Kang W, Wang L, et al. Antithrombotic effect and mechanism of *Rubus* spp. Blackberry. Food Funct. 2017;8(5):2000-12.
- Han N, Gu Y, Ye C, Cao Y, Liu Z, Yin J. Antithrombotic activity of fractions and components obtained from raspberry leaves (*Rubus chingii*). Food Chem. 2012;132(1):181-5.
- Plackal George B, Thangaraj P, Sulaiman C, Piramanayagam S, Ramaswamy SK. Bioassay directed isolation and biological evaluation of compounds isolated from *Rubus fairholmanus* Gard. Biomed Res Int. 2014.
- Iriwohan TS, Wrolstad RE. Polyphenolic composition of Marion and Evergreen Blackberries. J Food Sci. 2004;69(4).
- Surya M, Rahman W. Flowering and fruiting phenology of *Rubus* spp. In Cibodas Botanical Garden, Indonesia. Agrivita. 2012;34(2):193.
- Petreanu M, Ferreira EK, Sagaz AP, Vendramini-Costa DB, Ruiz AL, De Carvalho JE, et al. Uncommon trimethoxylated flavonol obtained from *Rubus rosaeifolius* leaves and its antiproliferative activity. Evidence-based Complement Altern Med. 2015;2015:6-11.
- Kanegusuku M, Sbars D, Bastos E. Phytochemical and analgesic activity of extract, fractions and a 19-hydroxyursane-type triterpenoid obtained from *Rubus rosaeifolius* (Rosaceae). Biol Pharm Bull. 2007;30(5):999-1002.
- Abu Bakar MF, Ismail NA, Isha A, Mei Ling AL. Phytochemical composition and biological activities of selected Wild Berries (*Rubus moluccanus* L., *R. fraxinifolius* Poir, and *R. alpestris* Blume). Evidence-based Complement Altern Med. 2016.

18. Blois MS. Antioxidant determinations by the use of a stable free radical. *Nature*. 1958;181(4617):1199-200.
19. Wojdylo A, Oszmianski J, Czemerys R. Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chem*. 2007;105(3):940-9.
20. Kedare SB, Singh RP. Genesis and development of DPPH method of antioxidant assay. *J Food Sci Technol*. 2011;48(4):412-22.
21. Alam N, Bristi NJ. Review on *in vivo* and *in vitro* methods evaluation of antioxidant activity. *Saudi Pharm J*. 2013;21(2):143-52.

Article History: Submission Date : 19-11-2017 ; Revised Date : 28-01-2018; Acceptance Date : 12-04-2018.

Cite this article: Desmiaty Y, Elya B, Saputri FC, Hanafi M, Prastiwi R. Antioxidant Activity of *Rubus fraxinifolius* Poir. and *Rubus rosifolius* J. Sm. Leaves. *J Young Pharm*. 2018;10(2)Suppl:s93-s96.

ORIGINALITY REPORT

9%

SIMILARITY INDEX

5%

INTERNET SOURCES

6%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1

res.mdpi.com

Internet Source

1%

2

www.coursehero.com

Internet Source

1%

3

Maria Kapiszewska, Agnieszka Cierniak, Monika A. Papiez, Agata Pietrzycka, Marek Stepniewski, Adam Lomnicki. "Prolonged Quercetin Administration Diminishes the Etoposide-Induced DNA Damage in Bone Marrow Cells of Rats", Drug and Chemical Toxicology, 2008

Publication

1%

4

Christian Mertz, Veronique Cheynier, Ziya Günata, Pierre Brat. "Analysis of Phenolic Compounds in Two Blackberry Species (*Rubus glaucus* and *Rubus adenotrichus*) by High-Performance Liquid Chromatography with Diode Array Detection and Electrospray Ion Trap Mass Spectrometry", Journal of Agricultural and Food Chemistry, 2007

Publication

<1%

5	www.wjgnet.com Internet Source	<1%
6	www.jove.com Internet Source	<1%
7	Submitted to Student Paper	<1%
8	www.pnas.org Internet Source	<1%
9	Submitted to Universidad Estadual Paulista Student Paper	<1%
10	www.florajournal.com Internet Source	<1%
11	T. Yuliani, E. Saepudin, T. A. Ivandini. "Anodic stripping voltammetry of Ni(OH) ₂ nanoparticles in acid solution using boron-doped diamond electrodes", AIP Publishing, 2018 Publication	<1%
12	Canini, A.. "Gas chromatography-mass spectrometry analysis of phenolic compounds from <i>Carica papaya</i> L. leaf", Journal of Food Composition and Analysis, 200711 Publication	<1%
13	M Amzad Hossain, Muhammad Dawood Shah, Mahyar Sakari. "Gas chromatography–mass spectrometry analysis of various organic	<1%

extracts of *Merremia borneensis* from Sabah",
Asian Pacific Journal of Tropical Medicine, 2011

Publication

14

www.apjtb.com

Internet Source

<1%

15

www.google.ml

Internet Source

<1%

16

www.jisikworld.com

Internet Source

<1%

17

Submitted to University of Birmingham

Student Paper

<1%

18

Medine Güllüce. "Biological activities of the essential oil and methanolic extract of *Micromeria fruticosa*(L) Druce ssp *serpyllifolia*(Bieb) PH Davis plants from the eastern Anatolia region of Turkey", Journal of the Science of Food and Agriculture, 05/2004

Publication

<1%

19

Submitted to 7996

Student Paper

<1%

20

ir.library.oregonstate.edu

Internet Source

<1%

21

Anna Tomlik-Wyremblewska, Raymond W.J.M. Van der Ham, Piotr Kosiński. "Pollen morphology of genus *Rubus* L. Part III. Studies

<1%

on the Malesian species of subgenera
Chamaebatus L. and Idaebatus L.", Acta
Societatis Botanicorum Poloniae, 2011

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On