

HARIYANTI-Synthesis and Preliminary In Vitro Anti-inflammatory Evaluation of Mannich Bases Derivatives of 4'-Methoxy-substituted of Asymmetrical Cyclovalone Analogs

by Hariyanti Uploaded By Mas'ud Syaiful

Submission date: 29-Mar-2023 01:36PM (UTC+0700)

Submission ID: 2049783162

File name: 1767-3686-1-PB_1_-_Dr_apr_Hariyanti_M.Si..pdf (1.34M)

Word count: 2703

Character count: 14612

10

Synthesis and Preliminary *In Vitro* Anti-inflammatory Evaluation of Mannich Bases Derivatives of 4'-Methoxy-substituted of Asymmetrical Cyclovalone Analogs

Nur Rahmawati¹, Hariyanti Hariyanti², Fadlina Chany Saputri¹, Hayun Hayun^{1,*}

1. Faculty of Pharmacy, Universitas Indonesia, Depok 16424, West Java, Indonesia
2. Faculty of Pharmacy, Universitas Muhammadiyah Prof. DR. HAMKA, Jakarta. 12130 Jakarta

Info Article

Submitted: 11-12-2019

Revised: 10-01-2020

Accepted: 20-03-2020

*Corresponding author

Hayun Hayun

Email:

hayun@farmasi.ui.ac.id

ABSTRACT

Two of Mannich bases derivatives of 4'-methoxy-substituted of asymmetrical cyclovalone analog (ACA) (**2a** and **2b**) were synthesized. The synthesized compounds and the other two Mannich bases derivatives of 4'-methoxy-substituted ACA (**2c** and **2d**) were evaluated for their *in-vitro* anti-inflammatory activity preliminary by protein denaturation inhibition method using a final concentration of 1.57 μ M. The study found that all the Mannich bases exhibited anti-inflammatory potential with inhibition ranging from 33.17- 42.47%. The activity of **2b** (42,47%) and **2d** (41.90%) was higher than that of diclofenac sodium (35.27%) and the parent compound **1** (38.16%). As a conclusion, **2b** and **2d** have a prospect as a potential candidate for an anti-inflammatory agent. Further study should be done using more specific methods.

Keywords: Mannich bases derivatives, asymmetrical, ayclovalone, synthesis, *in-vitro* anti-inflammatory, protein denaturation.

INTRODUCTION

Curcumin is well-documented to have anti-inflammatory activity with low toxicity. However, clinical usage of the compound is limited due to its stability and bioavailability (Prasad *et al.*, 2008; Anand *et al.*, 2008; Wang *et al.*, 1997). The mono-carbonyl analogs of curcumin (MACs), such as cyclovalone, showed a more stable chemical structure and better pharmacokinetic profile. Several of them were more active as anti-inflammatory agents than curcumin. (Liang *et al.*, 2008; Zhao *et al.*, 2013; Lamperti *et al.*, 2014; Zhao *et al.*, 2015).

In the past few years, some asymmetrical MACs were prepared, and among them indicated potent biological activity as anti-inflammation (Zhang *et al.*, 2014a; Zhang *et al.*, 2014b; Aluwi *et al.*, 2016). Introduction of morpholine Mannich base moiety into asymmetrical MACs bearing a cyclohexanone linker (= asymmetrical cyclovalone analogs, ACA) (Figure 1), increased the anti-inflammatory activity of the parent compound. 5-Morpholinomethyl-4'-methoxy ACA showed the highest activity, which almost comparable to diclofenac sodium (Putri *et al.*, 2018). Further investigation found that 5-dimethylaminomethyl

ACA (Figure 1, 4'-R = H) exhibited anti-inflammatory about four-fold than curcumin and equal to diclofenac sodium (Hayun *et al.*, 2019). However, the anti-inflammatory activity of Mannich bases derivatives of 4'-methoxy-substituted of ACA has not elucidated yet.




Figure 1. Asymmetrical Cyclovalone Analogs (ACA) (Putri *et al.*, 2018; Hayun *et al.*, 2019)

Hence, as a continuation of the study in exploring the anti-inflammatory activity of Mannich bases derivatives of ACA, in the present study, we synthesized 5-dimethylaminomethyl-4'-methoxy ACA (**2a**) and 5-(N-methylpiperazino)methyl-4'-methoxy ACA (**2b**) (Figure 2). The synthesized compounds and the other two Mannich bases derivatives of 4'-methoxy-substituted ACA (**2c** and **2d**) then were evaluated for their preliminary *in-vitro* anti-inflammatory activity.

14

MATERIAL AND METHODS

2,6-(4'-Methoxybenzylidene-4-hydroxy-3-methoxybenzylidene)cyclohexanone (= 4'-methoxy ACA) (**1**), 5-diethylaminomethyl-4'-methoxy ACA (**2c**), 5-morpholinomethyl-4'-methoxy ACA (**2d**), and cyclovalone, were documented by earlier researchers (Hayun *et al.*, 2017; Prasetyaningrum *et al.*, 2018; and Putri *et al.*, 2018). Diclofenac sodium was obtained from PT Kimia Farma, Indonesia, while other chemicals and solvent were supplied from chemical suppliers of Sigma-Aldrich or Merck. The compounds' melting point was measured by the melting point apparatus (Stuart Scientific, UK), infrared spectra and NMR spectra were recorded by FT-IR Spectrometer (Agilent Technologies, USA) and NMR spectrometer (Agilent Technologies, USA) respectively. While the mass spectra were recorded by LC-MS with ESI (+) mode (UNIFI, Waters, USA).

Synthesis of Mannich Bases derivatives of 4'-methoxy ACA (**2a** and **2b**)

The compounds **2a** and **2b** were synthesized using the method for preparation of **2c** and **2d** reported previously (Prasetyaningrum *et al.*, 2018; Putri *et al.*, 2018). A cooled solution in ethanol of **1** (2 mmol) was stirred and added a solution in ethanol of dimethylamine (for **2a**) or 1-methylpiperazine (for **2b**) (6 mmol) and formaldehyde solution (6 mmol). The mixture was further stirred for 30 minutes at r.t., then refluxed for 3 h (for **2a**) and 6 h (for **2b**) until the reaction was completed (TLC monitoring). After that, the products were isolated and then purified using a column chromatography technique to get **2a** and **2b**.

Preliminary *in-vitro* anti-inflammatory Evaluation.

The evaluation was done using inhibition of albumin denaturation technique, as reported previously with minor modification (Putri *et al.*, 2018). Mixture of 0.5mL of standard or test compounds solution in methanol (15.72 μ M) or solvent (for control) and 4.5mL Bovine Saline albumin (BSA) 0.5% in tris-buffer saline pH 6.3 was incubated at 37°C for 15min, then heated at 70°C in a water bath for 5min, and cooled to reach room temperature. The turbidity was measured at 660nm. The inhibition (%) of the denaturation was calculated with the formula:

$$\% \text{ inhibition} = \left[\frac{Ac - As}{Ac} \right] \times 100\%$$

where Ac was absorbance of control; As was absorbance with sample addition.

RESULT AND DISCUSSION

Chemistry

Scheme of the synthesis of compound **2a-b** (Figure 2).




Figure 2. Scheme of the synthesis of Mannich bases derivatives of 4'-methoxy-substituted ACA

Infrared, ^1H and ^{13}C NMR and mass spectral data obtained were presented in Table II. The infrared spectra **2a** and **2b** (Figures 3) showed the bands of C-O-C and C-N at 1,265-1,023 cm^{-1} ; C-H aliphatic at 2.935-2.829, while the α,β -unsaturated carbonyl groups, the C=C aromatic or ethylenic and alkyl were observed as strong band peaks at 1.658-1.507 and 1.459 cm^{-1} , respectively. The ^1H NMR (Figures 4), the protons of methylene connecting N of dimethylamine or N-methylpiperazine to aromatic ring appeared as a singlet at 3.71 ppm. Six protons from dimethylamine group in **2a** appeared as singlet peak at 2.34 ppm, eight protons from piperazine group in **2b** appeared as multiplet peak at 2.99ppm, and three protons of N-methylpiperazine group in **2b** appeared as a singlet at 2.31ppm. While, the two protons from the ethenyl chain appeared as two singlets at 7.61-7.67ppm (1H, respectively) confirming the structures were asymmetric (Silverstein *et al.*, 2005). The structures elucidation was supported further by ^{13}C NMR (Figures 5 and mass spectra (Figures 6). All data confirmed full agreement with the structures expected.

Table I. Physical and spectral data of compound **2a** and **2b**

Compd	Physical	Infra red (cm ⁻¹)	Spectra		Massa (M+H ⁺) (m/z)
			¹ H NMR, δ (ppm)	¹³ Carbon NMR, δ (ppm)	
2a	Red crystal, mp. 100-102°C	2,931-2,829, 1,658, 1,554, 1,507, 1,459, 1,246, 1,138, and 1,023.	7.67 (1H,s), 7.62 (1H,s), 7.50 (2H,d), 7.07 (1H,s), 23 3 (2 H,d), 6.91(1H,s), 24 3 (3H,s), 3.83 (3H,s), 3.72 (2H,s), 2.94 (4H,m), 2.34 (6H,s), 24 8 (2H,m)	189 (1C), 162 (1C), 150 & 149 (2C), 137-115 (13C), 63 (1C), 55 & 56 (2C), 45 (2C), 28-23 (3C)	Found: 408.21680 Calc for neutral mass of C ₂₅ H ₂₉ N ₀₄ = 407.20966; Mass Error = -0.3 mDa
2b	Yellow crystal, mp. 123-126°C	2,935-2,864, 1,655, 1,589, 1,508, 1,459, 1,246, 1,142, and 1,023.	7.67 (1H,s), 7.62 (1H,s), 36 6 (2H,d), 7.01 (1H,s), 23 9 (1H,s), 6.97 (2H,d), 3.84 (3H,s), 3.83 (3H,s), 3.75 (2H,s), 2.92(4H,m), 2.59 (8H,m), 2.30 (3H,s), 1.80 (2H, m)	192 (1C), 161 (1C), 149 & 148 (2C), 138-114 (13C), 59 (1C), 55 & 56 (2C), 53 & 56 (4C), 46 (1C), 30-24 (3C)	Found: 463.25921 Calc for neutral mass of C ₂₈ H ₃₄ N ₂ O ₄ = 462.25186; Mass Error = -0.3 mDa







Figure 3. Infra red spectrum of **2a** and **2b**



17
Figure 4. ¹H NMR spectrum of 2a and 2b



17
Figure 5. ¹³C NMR spectrum of 2a and 2b



17
Figure 6. Mass spectrum of 2a and 2b

²⁷ **In-vitro Anti-inflammatory activity**

Protein denaturation *in vivo* is one of the causes of inflammation. One of the mechanisms of antirheumatic activity of NSAIDs is by inhibiting this denaturation (Umapathy *et al.* 2010; Gunathilake *et al.*, 2018). Denaturation of protein can be induced by UV or heat. The denaturation is making protein aggregation, which can lead to the production of superoxide and nitric oxide by macrophages, which stimulates inflammation (Guzik *et al.*, 2003). Compounds inhibiting the heat-induced protein denaturation are considered to have potential anti-inflammatory activity (Chandra *et al.* 2012; Jagtap *et al.*, 2011).

²⁵ Table II. Inhibition (%) of heat-induced albumin denaturation by 1.57 μ M of the test compounds.

Compound	Inhibition (%) \pm SD	
1	38.16	0.14
2a	33.17	0.06
2b	42.47	0.02
2c	38.86	0.04
2d	41.90	1.63
Cyclovalone	19.64	0.09
Diclofenac Sodium	35.27	0.02




Figure 7. Inhibition \pm SD (%) (n=3) of heat-induced albumin denaturation the Mannich base derivatives of 4'-methoxy-substituted of asymmetrical cyclovalone analogs (ACA) (2a-d) compared to the parent compound (1), cyclovalone and sodium diclofenac at a final concentration of 1.57 μ M.

The preliminary activity of the synthesized compounds (Table II and Figure 7). At a final concentration of 1.57 μ M indicated that all the Mannich bases inhibited the heat-induced protein denaturation ranging from 33.17–42.47%. This study found that the activity of **2b** [5-(N-methylpiperazino)methyl-4'-methoxy ACA] (42.47%) and **2d** (5-morpholinomethyl-4'-methoxy ACA) (41.90%) was higher than that of diclofenac sodium (35.27%), the parent compound

1 (38.16%), and cyclovalone (19.64%). It was in line with the result of the introduction of the Mannich bases to dehydrozingerone reported earlier (Hayun *et al.*, 2018). Therefore, the compounds have a prospect as a potential candidate for an anti-inflammatory agent. However, to ensure their biological activities, further study should be done using more specific methods.

CONCLUSION

Two Mannich base derivatives of asymmetrical cyclovalone analogs (ACA) were synthesized successfully. Their preliminary *in-vitro* anti-inflammatory evaluation indicated that the compound **2b** [5-(N-methylpiperazino)methyl-4'-methoxy ACA] and **2d** (5-morpholinomethyl-4'-methoxy ACA) exhibited the highest activity. Their biological activities were higher than diclofenac sodium. However, further study should be done using more specific methods.

ACKNOWLEDGMENT

This work was supported by Universitas Indonesia (PITTA B, Research Grant 2019). Thank Laboratory of Organic Chemistry, Institut Teknologi Bandung, Indonesia, for recording NMR spectra; and to the Research Center for Chemistry, the Indonesian Institute of Sciences, Serpong, Indonesia, for recording mass spectra.

REFERENCES

- Aluwi MFMF., Rullah K., Yan BM., Leong SW., Bahari MNA., *et al.*, 2016. Synthesis of unsymmetrical monocarbonyl curcumin analogues with potent inhibition on prostaglandin E2 production in LPS induced murine and human macrophages cell lines. *Bioorg Med Chem Let.* 26(10):2531-2538. <http://doi.org/10.1016/j.bmcl.2016.03.092>
- Anand P., Thomas SG., Kunnumakkara AB., Sundaram C., Harikumar KB. *et al.*, 2008. Biological activities of curcumin and its analogues (congeners) made by man and mother nature. *Biochem Pharmacol.* 76:1596-1611. <http://doi.org/10.1016/j.bcp.2008.08.008>
- Chandra S., Chatterjee P, Dey P, Bhattacharya S. 2012. Evaluation of *in vitro* anti-inflammatory activity of coffee against the denaturation of protein. *Asian Pac J Trop Biomed.* 2:S178-S180. [https://doi.org/10.1016/S2221-1691\(12\)60154-3](https://doi.org/10.1016/S2221-1691(12)60154-3)

- 20 Gunathilake KDPP., Ranaweera KKDS., Rupasinghe HPV. 2018. *In Vitro* Anti-Inflammatory Properties of Selected Green Leafy Vegetables. *Biomedicines* 6, 107. <http://doi.org/10.3390/biomedicines6040107>
- 18 Guzik TJ., Korbut R., Adamek-Guzik T. 2003. Nitric oxide and superoxide in inflammation and immune regulation. *J Physiol Pharmacol.* 35(4):469-487. <https://www.ncbi.nlm.nih.gov/pubmed/14726604>
- 2 Hayun H., Maggad¹⁰ BP., Kurnia A., Hanifah A., Yuliandi M., et al., 2019. Anti-Inflammatory And Antioxidant Activity Of Synthesized Mannich Base Derivatives of (2E,6E)-2-[(4-Hydroxy-3-methoxyphenyl)methyl-idene]-6-(phenylmethylidene)cyclohexan-1-one. *Int J App Pharm.* 11 (Special Issue1):246-250. <http://dx.doi.org/10.22159/ijap.2019.v11s1.19448>
- 4 Hayun H., Jatmika C., Purwati EM., Salim S., Kurniawan R. et al. 2017. Synthesis and Free Radical-scavenging Activities of Di-Mannich Bases of Cyclovalone Derivatives. *Orient J Chem.* 33(6):2742-57. <http://dx.doi.org/10.13005/ojc/330607>
- 1 Hayun H., Arrahman A., Purwati EM., Yanuar A., Fortunata F., Suhargo F. et al. 2018. Synthesis, anti-inflammatory, and antioxidant activity of Mannich bases of dehydrozingerone derivatives. *J Young Pharm.* 10:s6-10:56-60. <http://doi.org/10.5530/jyp.2018.2s.2>
- 16 Jagtap VA., Agasimundim YS., Jayachandran E., Sathe BS. 2011. *In vitro* anti-inflammatory activity of 2-amino-3-(substituted benzylidene-carbohydrazide)-4,5,6,7-tetrahydrobenzothiofenes. *J Pharm Res.* 4(2):378-9. <http://jprsolutions.info/newfiles/journal-file-56cc7064d2aa84.22735300.pdf>
- 5 Lamperti M., Maspero A., Tonnesen HH., Bondani M., Nardo L. 2014. Elucidation of the relationships between H-bonding patterns and excited state dynamics in cyclovalone. *Molecules.* 19(9):13282-13304. <https://doi.org/10.3390/molecules190913282>
- 5 Liang G., Li X., Chen L., Yang S., Wu X., et al., 2008. Synthesis and anti-inflammatory activities of mono-carbonyl analogues of curcumin. *Bioorg Med Chem Lett.* 18(4): 1525-1529. <http://doi.org/10.1016/j.bmcl.2007.12.068>
- 7 Prasad S., Tyagi AK., Aggarwal BB. 2014. Recent developments in delivery, bioavailability, absorption and metabolism of curcumin: the golden pigment from golden spice. *Cancer Res Treat.* 46(1):2-18. <https://doi.org/10.4143/cr.11.14.46.1.2>
- Prasetyaningrum PW., Bahtiar A., Hayun H. 2018. Synthesis and cytotoxicity evaluation of novel asymmetrical mono-carbonyl analogs of curcumin (AMACs) against Vero, HeLa, and MCF7 Cell Lines. *Sci Pharm.* 86(2):25. <http://doi.org/10.3390/scipharm86020025>
- 2 Putri TN., Bahtiar A., Hayun H. 2018. Synthesis, antioxidant, and antiinflammatory activity of morpholine Mannich base of AMACs ((2E,6E)-2-((4-hydroxy-3-[morpholin-4-yl)methyl]phenyl)methylidene)(phenylmethylidene)cyclohexan-1-one) and its analogs. *21 App Pharm Sci.* 8:19-25. <http://doi.org/10.7324/JAPS.2018.8503>
- 2 Silverstein RM., Webster FX., Kiemle DJ. 2014. *Spectrometric Identification of Organic Compounds.* 7th ed. New York, USA: John Wiley and Sons, Inc.
- Umopathy E., Ndebia EJ., Meeme A., Adam B., Menziwa P., et al. 2010. An experimental evaluation of *Albica setosa* aqueous extract on membrane stabilization, protein denaturation and white blood cell migration during acute inflammation. *J Med Plants Res.* 4:789-795. <http://doi.org/10.5897/JMPR10.056>
- 12 Wang YJ., Pan MH., Cheng AL., Lin LI., Ho YS., et al., 1997. Stability of curcumin in buffer solutions and characterization of its degradation products. *J Pharm Biomed Anal.* 1997; 32 15:1867-1876. [http://doi.org/10.1016/s0731-7085\(96\)02024-9](http://doi.org/10.1016/s0731-7085(96)02024-9)
- 19 Zhao C., Liu Z., Liang G. 2013. Promising curcumin-based drug design: Mono-carbonyl analogues of curcumin (MACs). *Curr Pharm Des.* 19:2114-2135. <http://doi.org/10.2174/1381612811319110012>
- 9 Zhao C., Zhang Y., Zou P., Wang JJ., He W., et al., 2015. Synthesis and biological evaluation of a novel class of curcumin analogs as anti-inflammatory agents for prevention and treatment of sepsis in mouse model. *Drug*

¹ *Des Dev Ther.* 9:1663-1678.
<https://doi.org/10.2147/DDDT.S75862>
Zhang Y., Zhao L., Wu J., Jiang X., Dong L., *et al.*,
2014a. Synthesis and evaluation of a series
of novel asymmetrical curcumin analogs for
¹¹ treatment of inflammation. *Molecules*.
19:7287-7307.

<https://doi.org/10.3390/molecules19067287>
¹⁵ Zhang Y., Jiang X., Peng K., Chen C., Fu L., *et al.*,
2014b. Discovery and evaluation of novel
antiinflammatory derivatives of natural
bioactive curcumin. *Drug Des Devel Ther.*
8:2161-2171.
<https://doi.org/10.2147/DDDT.S69914>

HARIYANTI-Synthesis and Preliminary In Vitro Anti-inflammatory Evaluation of Mannich Bases Derivatives of 4'-Methoxy-substituted of Asymmetrical Cyclovalone Analogs

ORIGINALITY REPORT

40%
SIMILARITY INDEX

36%
INTERNET SOURCES

32%
PUBLICATIONS

22%
STUDENT PAPERS

PRIMARY SOURCES

1	innovareacademics.in Internet Source	4%
2	www.innovareacademics.in Internet Source	4%
3	www.preprints.org Internet Source	3%
4	www.jyoungpharm.org Internet Source	2%
5	journals.plos.org Internet Source	2%
6	m.japsonline.com Internet Source	2%
7	"Sustainable Agriculture Reviews 43", Springer Science and Business Media LLC, 2020 Publication	2%
8	Nasreddine El Omari, Fatima Ezzahrae Guaouguaou, Naoual El Meniy, Taoufiq	2%

Benali et al. "Phytochemical and biological activities of Pinus halepensis Mill., and their ethnomedicinal use", Journal of Ethnopharmacology, 2020

Publication

9	www.frontiersin.org Internet Source	2%
10	Akash Jain, Jasmine Chaudhary, Harpreet Khaira, Bhawna Chopra, Ashwani Dhingra. "Piperazine: A Promising Scaffold with Analgesic and Anti-inflammatory Potential", Drug Research, 2020 Publication	1%
11	www.mdpi.com Internet Source	1%
12	Submitted to Benedictine University Student Paper	1%
13	pdfs.semanticscholar.org Internet Source	1%
14	repository.uin-malang.ac.id Internet Source	1%
15	Morena Zusso, Giulia Mercanti, Federica Belluti, Rita Maria Concetta Di Martino, Andrea Pagetta, Carla Marinelli, Paola Brun, Eugenio Ragazzi, Rita Lo, Stefano Stifani, Pietro Giusti, Stefano Moro. "Phenolic 1,3-	1%

diketones attenuate lipopolysaccharide-induced inflammatory response by an alternative magnesium-mediated mechanism", British Journal of Pharmacology, 2017

Publication

16

www.ftstjournal.com

Internet Source

1 %

17

Jeong-Myeong Ha, Andrew Solovyov, Alexander Katz. "Postsynthetic Modification of Gold Nanoparticles with Calix[4]arene Enantiomers: Origin of Chiral Surface Plasmon Resonance", Langmuir, 2008

Publication

1 %

18

qmro.qmul.ac.uk

Internet Source

1 %

19

researchspace.auckland.ac.nz

Internet Source

1 %

20

ouci.dntb.gov.ua

Internet Source

1 %

21

link.springer.com

Internet Source

1 %

22

Submitted to Universiti Sains Malaysia

Student Paper

1 %

23

docksci.com

Internet Source

1 %

24

patents.google.com

Internet Source

1 %

25

Gwatidzo Luke, Chowe Leo, Musekiwa and Netai Mukaratirwa-Muchanyereyi Cexton. "In vitro anti-inflammatory activity of Vangueria infausta: An edible wild fruit from Zimbabwe", African Journal of Pharmacy and Pharmacology, 2018

Publication

1 %

26

Submitted to Universitas Indonesia

Student Paper

<1 %

27

Submitted to CVC Nigeria Consortium

Student Paper

<1 %

28

Farmiza Begum, Suman Manandhar, Gautam Kumar, Raghuvir Keni et al.

"Dehydrozingerone promotes healing of diabetic foot ulcers: a molecular insight", Journal of Cell Communication and Signaling, 2022

Publication

<1 %

29

op.niscair.res.in

Internet Source

<1 %

30

Gokul Sudhakaran, Pandurangan Prathap, Ajay Guru, Ravi Rajesh et al. "Anti - inflammatory role demonstrated both in vitro and in vivo models using nonsteroidal tetranortriterpenoid, Nimbin (N1) and its

<1 %

analogs (N2 and N3) that alleviate the domestication of alternative medicine", Cell Biology International, 2022

Publication

31

Chenguang Zhao, Yali Zhang, Peng Zou, Jian Wang, Wenfei He, Dengjian Shi, Huameng Li, Guang Liang, Shulin Yang. "Synthesis and biological evaluation of a novel class of curcumin analogs as anti-inflammatory agents for prevention and treatment of sepsis in mouse model", Drug Design, Development and Therapy, 2015

Publication

<1 %

32

Submitted to Emmanuel College

Student Paper

<1 %

33

Pradeep Pakalapati, Afrin Shanwaz. "Anaemic retinopathy in megaloblastic anaemia", Nepalese Journal of Ophthalmology, 2020

Publication

<1 %

34

www.annualreviews.org

Internet Source

<1 %

35

www.diva-portal.se

Internet Source

<1 %

36

www.freepatentsonline.com

Internet Source

<1 %

37

Atulkumar Kamble, Ravindra Kamble, Suneel Dodamani, Sunil Jalalpure et al. "Design,

<1 %

synthesis and pharmacological analysis of 5-[4'-(substituted-methyl)[1,1'-biphenyl]-2-yl]-1H-tetrazoles", Archives of Pharmacal Research, 2017

Publication

38

Nitesh Tamang, Gayathri Ramamoorthy, Mayank Joshi, Angshuman Roy Choudury, Siva Kumar B., Nageswara Rao Golakoti, Mukesh Doble. "Diarylidene cyclopentanone derivatives as potent anti-inflammatory and anticancer agents", Medicinal Chemistry Research, 2020

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off