

Investigation of Antioxidant and Antimicrobial Properties of Sunda Porcupine's

by Muhamad Arif Budiman Uploaded By Fauzan

Submission date: 03-May-2023 02:13PM (UTC+0700)

Submission ID: 2082875458

File name: ident_AO_and_AM_original_version_turnitin_1.pdf (346.16K)

Word count: 3929

Character count: 22056

**Investigation of Antioxidant and Antimicrobial Properties of Sunda Porcupine's
(*Hystrix javanica*, F.Cuvier 1823) Quills Ethanolic Crude Extract**

¹Muhamad Arif Budiman*, ²Pamungkas Rizki Ferdian**, ²Tri Hadi Handayani, ³Rizki Rabeca Elfirta, ⁴Masrukhin, ²Herjuno Ari Nugroho, ²Ni Luh Putu Rischa Phadmachanty, ²Wartika Rosa Farida, ⁵Ardya Widyastuti, ⁶Danita Dwi Sugiartanti,

¹Faculty of Medicine, Universitas Muhammadiyah Prof. DR. HAMKA. Tangerang City, Banten 15153, Indonesia

²Research Center for Applied Zoology, Cibinong, West Java 16911, Indonesia

³Research Center for Applied Microbiology, Cibinong, West Java 16911., Indonesia

⁴Research Center for Biosystematic and Evolution, Cibinong, West Java 16911, Indonesia

⁵DialoVet Animal Care, Bogor, West Java 16913, Indonesia

⁶Indonesian Research Center for Veterinary Science, Bogor 16114, Indonesia

*email: arifbudiman@uhamka.ac.id; **email: pamu003@brin.go.id.

ABSTRACT

The sunda porcupine (*Hystrix javanica*, F. Cuvier 1823) is a rodent-mammal species native to Indonesia and is utilized in traditional medicine for the treatment of various ailments. Some ethnic communities in Indonesia have traditional beliefs regarding sunda porcupine's quills, which are thought to relieve back pain and toothache. Despite this traditional knowledge, there is limited scientific research on the topic. The aim of this study was to identify active compound in an ethanolic crude extract of sunda porcupine's quills, and to evaluate its antioxidant and antimicrobial properties. The antioxidant activity was evaluated using DPPH-free radical scavenging assay while the antimicrobial activity was evaluated through microdilution resazurin assay. The total phenolic and flavonoid contents were also determined to support the antioxidant properties. The active compounds were identified using GC-MS with the NIST-11 library. The result showed that the extract possesses antioxidant properties (IC₅₀ 138,93 µg/mL) and antimicrobial properties against *E. coli*, *P. aeruginosa*, *S. aureus*, *B. subtilis* and *C. albicans* (IC₅₀ range 0.40-33.05 mg/mL). Total phenolic and total flavonoid content were 27.29 ± 2.20 mgGAE/g and 27.09 ± 1.66 mgQE/g, respectively. A total of 24 active compounds from the crude extract were identified. As much as 5 compounds serve as antioxidant agents, including: butylated hydroxytoluene; eicosane; 1-iodo-hexadecane; methyl ester hexadecanoic acid; and 2,6-dihexadecanoate-L-(+)-ascorbic acid. Furthermore, as much as 11 compounds serve as antimicrobial agents, including: tetradecane; pentadecane; 2-isopropyl-5-methyl-1-heptanol; hexadecane; butylated hydroxytoluene; eicosane; 1-iodo-hexadecane; methyl ester hexadecanoic acid; 3,5-bis(1,1-dimethylethyl)-4-hydroxy-, methyl ester benzenepropanoic acid; 2,6-dihexadecanoate-L-(+)-ascorbic acid; and octadecanoic acid. This study provides scientific validation for the use of the sunda

porcupine's quills in traditional medicine and highlights the potential for further research in animal bioprospecting.

Keywords: antioxidant, antimicrobial, GC-MS, sunda porcupine, *Hystrix javanica*

INTRODUCTION

Animal-based medicine ¹⁴ has been used for centuries in traditional medicine systems around the world. However, its use is not as widespread as that of plant-based medicine due to various factors such as cultural, religious, conservation, and abundance considerations. The medicine can be derived from diverse range of animal materials, including animal metabolites, body parts, or non-animal components such as bird nests, bee hives and cocoons. Inspired by ethnobiology and ethnomedicine, research into animal-based medicine has been conducted globally, including in Indonesia. Mardiasuti *et al.* (2021) documented the use of a variety of animal species in Indonesian traditional medicine, including 59 mammal species, 12 bird species, 37 reptile species, and 6 amphibian species, with porcupines being one of the species used.

The porcupine is widely used in Indonesian traditional medicine. In Kalimantan (Borneo), porcupine's quills are ground into flour and used to treat acne. Meanwhile the Dayak natives use the burning porcupine's quills to relieve back pain. In Java, some locals use the porcupine's quills as a medicine for toothaches and ulcers (Inayah, 2016; Krisyanto *et al.*, 2019). One species of the porcupine that is thought to be used in traditional medicine ¹⁰ is the sunda porcupine (*Hystrix javanica*, F. Cuvier 1823), which is reported to be found in certain regions of Indonesia including ¹⁰ Java, Bali, Sumbawa, Flores, Lombok, Madura, and Tonahdjampea (van Weers, ¹⁰ 1979; Van Weers, 1983; Woods & Kilpatrick, 2005; Aplin, 2016).

Recent research has investigated the potential of the sunda porcupine in animal-based medicine. Prawira *et al.* (2020) reported the rapid wound healing in this species, while Gifardi *et al.* (2022) demonstrated that sunda porcupine's quills hexane extract could inhibit the growth of *Staphylococcus aureus*, bacteria that infect the skin at certain concentration levels. Furthermore, Anita *et al.* (2018) reported that the tail meat of sunda porcupine possesses aphrodisiac potency.

The exploration of active compounds from sunda porcupine's quills is of interest since its utilization in traditional medicine and the limited current study in this area. Our study aims to identify active compounds in sunda porcupine's quills ethanolic crude extract as well as to evaluate its antioxidant and antimicrobial properties. This research is expected to provide new knowledge and contribute to the discovery of traditional medicine as a potential source of drugs.

MATERIALS AND METHODS

Sample Extraction

The simplisia of sunda porcupine's quills were obtained by drying the quills in an oven at 50 °C for 5 days. The simplisia were grounded to a size of 60 mesh and had a water content of 9.1%. A maceration method was used to extract the active compound inside simplisia with a simplisia to solvent ratio of 1:30 using 70% ethanol. The crude extract was obtained by evaporating all solvent using a rotary evaporator at 50 °C. The crude extract was then stored in 4 °C until further use.

Determining Antioxidant Activity using DPPH-Free Radicals Scavenging Assay

The DPPH (1,1-diphenyl-2-picrylhydrazyl) free radical was used to measure the antioxidant activity of the sunda porcupine's quills crude extract. The procedure

used was adapted from Handayani *et al.* (2022) and Aryal *et al.* (2019) with slight modifications. The sample was dissolved in methanol at concentrations ranging from 0 to 250 µg/mL. A mixture of 2 mL of the sample and 2 mL of 0.1 mM DPPH was incubated in the dark for 30 minutes. The absorbance of the sample was measured using Spectrophotometer UV-Vis at a wavelength of 517 nm. The antioxidant activity was calculated using the following formula:

$$\text{Antioxidant Activity (\%)} = \frac{\text{Absorbance blank} - \text{Absorbance sample}}{\text{Absorbance blank}} \times 100\%$$

In order to determine the IC₅₀ value, the antioxidant activity score obtained from the DPPH assay was plotted against the concentration of the sample. The concentration of the sample that caused a 50% reduction of DPPH was determined from the graph as the IC₅₀ value. Sample with lower IC₅₀ value were considered more effective in neutralizing free radicals.

Determining Total Phenolic Content (TPC)

A sample was first dissolved in distilled water to obtain a 1000 µg/mL solution. A standard curve was created using gallic acid with a range of serial concentrations from 0 to 200 µg/mL. A 0.2 mL sample or standard was added into 1.8 mL of distilled water and 0.2 mL of Folin-Ciocalteu reagent. The solution was homogenized and incubated for 6 minutes. After that, 2 mL of Na₂CO₃ 7% (w/v) was added to the solution, homogenized, and incubated for 90 minutes. The absorbance was then measured at 750 nm using a UV-Vis Spectrophotometer. TPC was calculated in mgGAE/g sample (Maeng *et al.* 2017).

Determining Total Flavonoid Content (TFC)

The determination of TFC followed the Dowd method as described by Aryal *et al.* (2019). A sample solution of 1000 µg/mL was prepared using distilled water. Quercetin was used as a standard with serial concentrations ranging 0 to 100 µg/mL. A 1 mL of prepared sample or standard was added to a mixture of 0.2 mL AlCl₃ 10% (w/v) in methanol, 0.2 mL CH₃COOK 1 M, and 5.6 mL distilled water. The solution was homogenized and incubated for 30 minutes. The absorbance was measured with a UV-Vis Spectrophotometer at 415 nm. TFC was calculated in mgQE/g sample.

Determining Antimicrobial Activity using Resazurin Assay

The antimicrobial activity was assessed using microdilution method incorporated with resazurin as the indicator of cell viability (EUCAST 2022; Sarker *et al.* 2007). The assay was conducted in 96-well plate against *Escherichia coli*, *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, and *Candida albicans*. Prior to antimicrobial assay, the target microbes were grown in Luria Bertani (LB) broth and incubated overnight in an incubator shaker. The target microbes were then adjusted using McFarland turbidity standard 0.5 and diluted 1000x, so that the final concentration of the cells was $\pm 1.5 \times 10^5$ CFU/mL. Antimicrobial assay of the extract was then conducted with concentration 100 mg/mL as the starting point and serially diluted to several concentrations followed by overnight incubation at 37°C. After the addition of 30 µL 0.1% resazurin, the cell suspension then incubated overnight and read under fluorescence with multimode reader Varioskan Lux (Thermo Scientific) with 530 excitation and 590 emission. The fluorescence data was used to determine the inhibition activity (%) using the following equation:

$$\% \text{ Inhibition} = \left(1 - \frac{\text{Fluorescence sample} - \text{Fluorescence of control (-)}}{\text{Fluorescence of control (+)} - \text{Fluorescence of control (-)}} \right) \times 100\%$$

The inhibitory concentration of 50% (IC₅₀) was determined through a dose-response relationship using linear regression analysis, with the transformation of the concentration to a logarithmic scale.

Identification of Active Compound Using GC-MS

The sunda porcupine's quills ethanolic crude extract was dissolved with dichloromethane to make a 1000 µg/mL solution. It was then filtered using a minisart syringe membrane 0.22 µm. The filtrate was injected into a GC-MS instrument equipped with an Rtx-5MS column (5% diphenyl: 95% dimethyl-polysiloxane) with length of 30 m and diameter of 0.25 mm. The mobile phase consisted of ultra high purity helium on 30 kPa. The injector temperature was set at 200 °C, the ion source at 230 °C, and the interphase at 280 °C, with a splitless injection mode. The oven temperature program was initiated at 60 °C and increased to 150 °C at a rate of 10 °C/minute and held for 3 minutes. The resulting chromatogram and m/z were compared with the NIST-11 database to identify the active compound.

RESULTS

Antioxidant Activity, TPC, and TFC

The antioxidant activity of sunda porcupine's ethanolic crude extract was assessed at varying concentrations, ranging from 0 to 250 µg/mL. The curve of antioxidant activity against the sample concentration was determined with a regression equation of $y = 0.3603x - 0.0551$ and $R^2 = 99.48\%$ as illustrated in Figure 1. The IC₅₀

of the extract's antioxidant against ²⁶ DPPH free radical scavenging activity was determined to be 138.93 µg/mL.

The standard curve of gallic acid and quercetin were used to ³³ determine the TPC and TFC as illustrated in Figure 2. The linear regression equation of gallic acid and quercetin were derived as $y = 0.005x + 0.0039$ ($R^2 = 99.98\%$) and $y = 0.0061x - 0.0066$ ($R^2 = 99.93\%$), respectively. The TPC and TFC of the extract were calculated to be 27.29 ± 2.20 mgGAE/g sample and 27.09 ± 1.66 mgQE/g sample, respectively.

Antimicrobial Activity

In present study, the antimicrobial activity of sunnda porcupine's quills ethanolic crude extract was evaluated for its antimicrobial activity against various microorganism, including ¹ *E. coli*, *P. aeruginosa*, *S. aureus*, *B. subtilis* and *C albicans*. The result is presented in Figure 3, which shows the linear regression of antimicrobial activity against the log concentration of the extract for each microorganism.

The data revealed that the extract possessed significant ²⁹ antimicrobial activity against all the tested microorganism. The highest antimicrobial activity was observed ⁸ against *S. aureus* with an IC_{50} of 0.40 mg/mL (Table 1), indicating that the extract could be a potential source of antibacterial agents, mainly to Gram positive bacteria.

The linear regression equation for ¹ *E. coli*, *P. aeruginosa*, *S. aureus*, *B. subtilis* and *C. albicans* were $y = 45.195x - 13.461$ ($R^2 = 90.42\%$), $y = 30.555x + 18.566$ ($R^2 = 94.07\%$), $y = 19.01x + 57.636$ ($R^2 = 91.23\%$), $y = 29.858x + 40.024$ ($R^2 = 92.87\%$), and $y = 77.819x + 68.222$ ($R^2 = 94.10\%$), respectively with *y-axis* representing the percentage of antimicrobial activity and *x-axis* representing the log concentration of the extract (Figure 3). These findings suggest that sunnda porcupine's ethanolic crude

extract could be a promising candidate for the development of novel antimicrobial agents.

Identification of Active Compound Using GC-MS

The CG-MS analysis of the extract resulted in the chromatogram as shown in Figure 4. After comparing the m/z data with the NIST-11 database, a total of 24 compounds were identified and are listed in Table 2. Among these compounds, 6 exhibit a relatively high intensity proportional to the percentage of area (more than 5%), including butylated hydroxytoluene (20.94%), 2,6-dihexadecanoate-L-(+)-ascorbic acid (14.60%), eicosane (8.86%), 5-methyl-1-phenylbicyclo [3.2.0] heptane (8.21%), pentadecane (6.88%), and hexadecane (6.30%). The highest intensity to the percentage of area in chromatogram was found at retention time of 13.952 min and presumably represent butylated hydroxytoluene. It has been reported by Ayaz *et al.* 1980 and Lim *et al.* 1987 that it possesses antioxidant and antimicrobial properties. Furthermore, the second highest intensity found at retention time of 23.860 min which presumably represent 2,6-dihexadecanoate-L-(+)-ascorbic acid also was known as antioxidant and antimicrobial agent (Hadi *et al.* 2016; Igwe & Okwunodulu 2014).

DISCUSSION

The sunda porcupine, an endemic mammal species of Indonesia, has a relatively wide distribution across several regions of the country, including Java, Bali, Sumbawa, Flores, Lombok, Madura, and Tonahdjampea (van Weers, 1979; Van Weers, 1983; Woods & Kilpatrick, 2005; Aplin, 2016). This broad distribution has fostered a connection between the sunda porcupine and the local people resulting traditional knowledge, including ethnobiology and ethnomedicine. This traditional knowledge is frequently not adequately documented and is instead passed down orally

from generation to generation, leading to difficulties in accessing this information. Some indigenous communities in Indonesia are reported to use the sunda porcupine's quills for medicinal purposes such as treating acne, relieving backpain, curing ulcer, and relieving toothache (Inayah, 2016; Krisyanto *et al.*, 2019). Furthermore, the sunda porcupine's quills is a unique skin derivate that provides an additional protective layer against harsh environment condition and acts as a defence tool against predators (Prawira *et al.* 2018). Therefore, it is plausible that the quills contain a certain compound that may be effective in combating environmental stress.

The present study investigates the active compounds found inside the sunda porcupine's quills, specifically focusing on their antioxidant and antimicrobial properties. The quills were extracted using a 70% ethanol solvent via the maceration method. Ethanol was chosen as the solvent due to its safety profile in comparison to other organic solvents. The maceration process was selected to minimize the risk of damaging the active compounds through the application of heat. The extract in this study was identified using GC-MS which successfully identified 24 active compounds as shown in Tabel 2. Most of these compounds were identified as biologically active, which aligns with previous studies.

The antioxidant activity of sunda porcupine's quills ethanolic crude extract was determined using DPPH free radicals scavenging assay in various concentrations. Furthermore, the antioxidant activities with their respective concentrations were plotted in a linear regression to determine the antioxidant IC₅₀. The antioxidant IC₅₀ of the extract in present study was 138.93 µg/mL, indicating ²⁸ the concentration of the extract required to neutralize 50% of free radicals. A lower IC₅₀ value indicates a smaller concentration of the sample needed to neutralize free radicals. The antioxidant properties mostly caused by the content of phenolic and flavonoid compound (Aryal *et*

al 2019; Maeng *et al.* 2017). The TPC and TFC of the extract was 27.29 ± 2.20 mgGAE/g and 27.09 ± 1.66 mgQE/g respectively. The flavonoids are a part of phenolics. The score of TPC and TFC are close, indicating the phenolics content are mostly in the form of flavonoids.

Moreover, the antioxidant properties of the extract were in line with the identified compound obtained from GC-MS analysis. There are 5 compounds with total of 47.33%, proportional to the percentage of area in chromatogram, responsible with antioxidant properties. These are butylated hydroxytoluene (20.94%), 2,6-dihexadecanoate-L-(+)-ascorbic acid (14.60%), eicosane (8.86%), methyl ester hexadecanoic acid (1.70%), 1-iodo-hexadecane (1.23%). Butylated hydroxytoluene was reported as an antioxidant to inhibit free radicals production for medicine and cosmetics (Ershov & Volod'kin 1962). The 2,6-dihexadecanoate-L-(+)-ascorbic acid is vitamin C derivative and it is important as a lipophilic antioxidant, antitumor, wound healing, and antimicrobial properties (Hadi *et al.* 2016; Igwe & Okwunodulu 2014). Eicosane is monoterpene hydrocarbon and is reported to have antioxidant and antiinflammatory properties by inhibiting the release of cytokines such as histamine, bradykinin, PGs, TXs, and LTs in rats (Kazemi 2015; Okechukwu 2020). Methyl ester hexadecanoic acid (methyl palmitate) is a fatty acid group with antioxidant, hypocholesterolaemia, and antiandrogenic properties (Astiti & Ramona, 2021; Hema *et al.* 2011). The 1-iodo-hexadecane has been reported in some plant extract and possesses antioxidant properties (Kim *et al.* 2022).

The present study also investigates the antimicrobial properties of sunda porcupine's quills ethanolic crude extract, in addition to its antioxidant properties. Resazurin assay was used in this study to determine the antimicrobial activity of the extract against *E. coli*, *P. aeruginosa*, *S. aureus*, *B. subtilis* and *C. albicans*. The use of

resazurin microdilution assay was selected since it can provide more accurate result through spectrophotometry, allowing for precise analysis signal readings. The antimicrobial activity was determined by measuring the resazurin readings after 24 hours of incubation for bacteria and 72 hours of incubation for yeast. Resazurin (7-Hydroxy-3H-phenoxazin-3-one 10-oxide) is a blue dye that can be irreversibly reduced by oxidoreductase in active bacteria to a pink and highly red fluorescent substance called resorufin (Chakansin *et al.* 2022). This method is based on detection of microbial viability by observing the colour change of resazurin from blue to purple or pink (Jia *et al.* 2020). The test was considered positive if the well contents were blue in colour, indicating the extract inhibits the growth of microbial, and negative if the well contents were pink, indicating the microbial is still growing in the medium wells. The test results were valid if negative control wells (without extract) remained pink (Germ *et al.* 2019).

The antimicrobial IC₅₀ values of sunda porcupine quills extract against *E. coli*, *P. aeruginosa*, *S. aureus*, *B. subtilis* and *C. albicans* was 23.65 mg/mL; 10.68 mg/mL; 0.40 mg/mL; 2.16 mg/mL; 33.05 mg/mL, respectively. These results indicate that the sunda porcupine quills extract has a broad antimicrobial range as it can inhibit the growth of Gram-positive bacteria, Gram-negative bacteria, spore-forming bacteria and yeast. In particular, the *S. aureus* bacteria species was highly sensitive for antimicrobial activity. However, the antimicrobial assay was limited to several species in present study. The further research related to the antimicrobial assay with other microbial species may complete the potential of sunda porcupine's quills as an antimicrobial agent. Sunda porcupine's quills ethanolic crude extract could inhibit the activity of *S. aureus* at the smallest concentration (0,4 mg/mL or 0,04% equivalent). The concentration of the sunda porcupines's quills ethanolic crude extract required to

inhibit *S. aureus* in the present study was found to be significantly lower (0.04%) than reported by Gifardi *et al.* (2022) for the sunda porcupine's quills extracted using hexane solvent, which required at least a concentration of 25% for inhibition of *S. aureus* growth. These findings suggest that the antimicrobial compounds in sunda porcupine's quills are more soluble in polar solvent such as ethanol 70%. Further research on the extraction and identification of the active compound in sunda porcupine's quills ⁸ could lead to the development of novel and effective antimicrobial agents.

Furthermore, the antimicrobial properties of the sunda porcupine ethanolic crude extract also appropriate with the GC-MS analysis. As much as 11 identified compounds have been reported possess antimicrobial properties. In total, it is about 70.12% of identified compound, proportional to the percentage of area in chromatogram, possess antimicrobial properties. These compounds include butylated hydroxytoluene (20.94%), 2,6-dihexadecanoate-L-(+)-ascorbic acid (14.60%), eicosane (8.86%), pentadecane (6.88%), hexadecane (6.30%), octadecanoic acid (3.48%), tetradecane (3.07%), ²⁰ 3,5-bis(1,1-dimethylethyl)-4-hydroxy-, methyl ester benzenepropanoic acid (1.76%), methyl ester hexadecanoic acid (1.70%), ⁹ 2-isopropyl-5-methyl-1-heptanol (1.30%), and 1-iodo-hexadecane (1.23%).

Butylated hydroxytoluene have been reported as antimicrobial to inhibit growth some microorganism (Ayaz *et al.* 1980; Lim *et al.* 1987). The 2,6-dihexadecanoate-L-(+)-ascorbic acid is ascorbic acid derivate which is potential to prevent and treat common cold, gum diseases, acne, skin infection, tuberculosis, dysentery, and dental caries (Igwe & Okwunodulu, 2014). Eicosane have been reported as antifungal, antibacterial, antitumor, inhibit foodborne pathogen, and has cytotoxic effect (Okechukwu 2020; Kazemi 2015; Akpuaka *et al.* 2013; Yogeswari *et al.* 2012; Hsouna ³²

et al. 2011). Pentadecane is reported as an antimicrobial compound by inhibited growth of *E. coli* and *S. typhi* (Firdaus *et al.* 2019; Martinac *et al.* 1987). Octadecanoic acid or stearic acid is fatty acid that displays antibacterial activity towards Gram-positive and Gram-negative bacteria (Abdalaziz *et al.* 2017; Casillas-Vargas *et al.* 2021). Hexadecane, tetradecane, 2-isopropyl-5-methyl-1-heptanol, and 1-iodohexadecane has reported to possess antifungal and antimicrobial properties (Selvin *et al.* 2009; Akpuaka *et al.* 2013; Yogeswari *et al.* 2012; Kim *et al.* 2022). Benzenepropanoic acid was also reported to be effective against microbes such as *E. coli*, *K. pneumoniae*, *P. aeruginosa*, and *C. albicans* (Amrati *et al.* 2021). Benzoic acid alone is known as a nonspecific antimicrobial agent with a wide spectrum of the activities against human pathogenic fungi and bacteria (Innocenti *et al.* 2009; Krátký *et al.* 2012). Methyl ester hexadecanoic acid was reported as antibacterial by disrupting bacterial cell wall and cell membrane (Astiti and Ramona 2021).

CONCLUSIONS

As much as 24 active compounds were identified from sunda porcupine's quills ethanolic crude extract using GC-MS. The extract was investigated and showed an antioxidant and antimicrobial properties. The IC₅₀ of antioxidant was 138.93 µg/mL, while the IC₅₀ of antimicrobial against *E. coli*, *P. aeruginosa*, *S. aureus*, *B. subtilis* and *C. albicans* were 23.65, 10.68, 0.40, 2.16, 33.05 mg/mL, respectively. The antioxidant properties also investigated through determination of TPC and TFC with value were 27.29 ± 2.20 mgGAE/g and 27.09 ± 1.66 mgQE/g, respectively. There were 5 identified compounds serve as antioxidant and 11 identified compounds serve as antimicrobial. The two highest intensity to the percentage of area in chromatogram were butylated hydroxytoluene (20.94% with RT=13.952 min) and 2,6-dihexadecanoate-L-(+)-

ascorbic acid (14.60% with RT= 23.860 min). These both compounds have been reported as antioxidant dan antimicrobial agent. This study provides scientific validation for the use of the sunda porcupine's quills in traditional medicine and highlights the potential for further research in animal bioprospecting.

41

ACKNOWLEDGMENTS

This research was supported by The Budget Implementation List of Indonesian Institute of Sciences Year 2021 (research grant number 46/A/DH/2021) and Faculty of Medicine, Universitas Muhammadiyah Prof. DR. HAMKA (Lemlit Grant no. 867/F.03.07/2022).

Figure Caption






Figure 1. Antioxidant Activity of Sunda Porcupine's Ethanollic Crude Extract using DPPH Free Radical Scavenging Assay




18
(a)




(b)


Figure 2. (a) Standard Curve of Gallic Acid and (b) Standard Curve of Quercetin.




36
(a)




(b)



(c)



(d)



(e)

Figure 3. Antimicrobial Activity of Sunda Porcupine's Quills Ethanolic Crude Extract Against (a) *E. coli*, (b) *P. aeruginosa*, (c) *S. aureus*, (d) *B. subtilis*, and (e) *C. albicans*.




Figure 4. Chromatogram of Sunda Porcupine's Quills Ethanolic Crude Extract

Table 1. The Antimicrobial IC₅₀ of Sunda Porcupine's Quills Ethanolic Crude Extract

Microorganism	Group	IC ₅₀ (mg/mL)
<i>Escherichia coli</i>	Gram negatif	23.65
<i>Pseudomonas aeruginosa</i>	Gram negatif	10.68
<i>Staphylococcus aureus</i>	Gram positif	0.40
<i>Bacillus subtilis</i>	Gram positif, sporadic	2.16
<i>Candida albicans</i>	Fungi	33.05

Table 2. The Active Compound Identified by GCMS Instrument with NIST 11 Database

No.	Retention Time (minute)	Area (%)	Name	Molecular Formula	Similarity (%)	Role(s)
1	8.921	3.07	Tetradecane	C ₁₄ H ₃₀	75	antifungal and antibacterial (Ozdemir et al. 2004)
2	9.167	6.88	Pentadecane	C ₁₅ H ₃₂	92	Antimicrobial (Firdaus et al. 2019; Martinac et al. 1987)
3	9.570	1.30	2-Isopropyl-5-methyl-1-heptanol	C ₁₁ H ₂₄ O	87	Antimicrobial (Selvin et al., 2009)
4	9.700	1.04	2-methyl-1-decanol	C ₁₁ H ₂₄ O	87	-
5	9.855	2.22	2,6,11-trimethyl-dodecane	C ₁₅ H ₃₂	91	-
6	11.024	1.08	E-14-Hexadecenal	C ₁₆ H ₃₀ O	84	-
7	13.199	1.33	2,6,11,15-tetramethyl-hexadecane	C ₂₀ H ₄₂	82	Flavoring agent (Pammi et al. 2021)
8	13.346	6.30	Hexadecane	C ₁₆ H ₃₄	90	Antifungal, antibacterial, (Akpuaka et al. 2013; Yogeswari et al. 2012)
9	13.952	20.94	Butylated Hydroxytoluene	C ₁₅ H ₂₄ O	95	Antioxidant and antibacterial (Ayaz et al. 1980; Lim et al. 1987)
10	18.020	1.15	Dodecyl ester chloroacetic acid	C ₁₄ H ₂₇ ClO ₂	88	-
11	18.361	8.86	Eicosane	C ₂₀ H ₄₂	87	Antifungal, antibacterial, inhibit foodborne pathogen, antitumor, cytotoxic effect (Kazemi 2015; Okechukwu 2020; Akpuaka et al. 2013; Yogeswari et al. 2012; Hsouna et al. 2011)
12	18.538	1.23	1-iodo-hexadecane	C ₁₆ H ₃₃ I	75	Inhibitory effect on AD-like lesions, antimicrobial, antioxidant, anticancer (Kim et al. 2022)
13	19.167	1.07	2-Methoxycarbonyl-2-methylbrendane	C ₁₂ H ₁₈ O ₂	65	-
14	20.459	1.73	2-fluoro-5,6-dimethoxybenzoic acid	C ₉ H ₉ FO ₄	59	-
15	20.539	2.70	Dimethyl (2E)-4-cyclopropyl-2-heptenedioate	C ₁₂ H ₁₈ O ₄	53	-
16	20.787	1.54	Hexadecanal	C ₁₆ H ₃₂ O	93	-
17	22.967	4.86	2-methyl-heptadecane	C ₁₈ H ₃₈	89	-
18	23.055	1.70	Methyl ester hexadecanoic acid	C ₁₇ H ₃₄ O ₂	92	antioxidant, antifungal, antimicrobial (Astiti & Ramona, 2021; Hema et al. 2011)
19	23.509	1.76	3,5-bis(1,1-dimethylethyl)-4-hydroxy-, methyl ester benzenepropanoic acid	C ₁₈ H ₂₈ O ₃	91	-
20	23.860	14.60	2,6-dihexadecanoate-L-(+)-ascorbic acid	C ₃₈ H ₆₈ O ₈	86	antioxidant, antitumor, wound healing, and antimicrobial properties (Hadi et al. 2016; Igwe & Okwunodulu 2014)
21	24.550	8.21	5-Methyl-1-phenylbicyclo[3.2.0]heptane	C ₁₄ H ₁₈	69	Antivirus (Poongulali & Sundararaman, 2016)
22	24.959	1.63	Heptadecyl- oxirane	C ₁₉ H ₃₈ O	93	-
23	27.667	3.48	Octadecanoic acid	C ₁₈ H ₃₆ O ₂	83	antibacterial and antifungal (Akpuaka et al. 2013)
24	27.845	1.32	Tetracosane	C ₂₄ H ₅₀	87	anticancer (Paudel et al. 2019)

Note: (-) Unknown roles

Investigation of Antioxidant and Antimicrobial Properties of Sunda Porcupine's

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

12%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	pdfs.semanticscholar.org Internet Source	2%
2	www.mdpi.com Internet Source	2%
3	www.ncbi.nlm.nih.gov Internet Source	1%
4	Chanthiwa Chakansin, Jakrawan Yostaworakul, Choochart Warin, Kornphimol Kulthong, Suwimon Boonrungsiman. "Resazurin rapid screening for antibacterial activities of organic and inorganic nanoparticles: Potential, limitations and precautions", Analytical Biochemistry, 2021 Publication	1%
5	www.researchgate.net Internet Source	1%
6	link.springer.com Internet Source	1%

Submitted to Universiti Teknologi Petronas

7	Student Paper	1 %
8	www.science.gov Internet Source	1 %
9	Sunil Kr Jha, Masahiro Imahashi, Kenshi Hayashi, Tadashi Takamizawa. "Data fusion approach for human body odor discrimination using GC-MS spectra", 2014 IEEE Ninth International Conference on Intelligent Sensors, Sensor Networks and Information Processing (ISSNIP), 2014 Publication	1 %
10	Submitted to Padjadjaran University Student Paper	1 %
11	Submitted to Universiti Putra Malaysia Student Paper	<1 %
12	Anni NURLIANI, Motoki SASAKI, Teguh BUDIPITOJO, Toshio TSUBOTA, Masatsugu SUZUKI, Nobuo KITAMURA. "An immunohistochemical study on testicular steroidogenesis in the Sunda porcupine (<i>Hystrix javanica</i>)", Journal of Veterinary Medical Science, 2019 Publication	<1 %
13	mdpi-res.com Internet Source	<1 %

14 Boreddy Srinivas Reddy. "Evaluation of antioxidant and antimicrobial properties of *Soymida febrifuga* leaf extracts", *Phytotherapy Research*, 07/2008
Publication <1 %

15 phcog.com
Internet Source <1 %

16 Greeshma Gopalan, Bernard Prabha, Alfred Joe, Thankappan Remadevi Reshmitha et al. " Screening of . seeds for antidiabetic properties and isolation of apiforol, a potential lead, with antidiabetic activity ", *Journal of the Science of Food and Agriculture*, 2018
Publication <1 %

17 juniperpublishers.com
Internet Source <1 %

18 www.hindawi.com
Internet Source <1 %

19 Lucía Valverde-Som, Ana Herrero, Celia Reguera, Luis Antonio Sarabia, María Cruz Ortiz. "A new multi-factor multi-objective strategy based on a factorial presence-absence design to determine polymer additive residues by means of head space-solid phase microextraction-gas <1 %

chromatography-mass spectrometry",
Talanta, 2023

Publication

20

www.rcdc.nd.edu

Internet Source

<1 %

21

escholarship.mcgill.ca

Internet Source

<1 %

22

Submitted to Higher Education Commission
Pakistan

Student Paper

<1 %

23

Valentina Nanni, Lorena Canuti, Angelo
Gismondi, Antonella Canini. "Hydroalcoholic
extract of Spartium junceum L. flowers
inhibits growth and melanogenesis in B16-F10
cells by inducing senescence", Phytomedicine,
2018

Publication

<1 %

24

patents.justia.com

Internet Source

<1 %

25

Jiangtao Hu, Yali Li, Ya Liu, Dong Il Kang, Hao
Wei, Byoung Ryong Jeong. "Hydrogen Sulfide
Affects the Root Development of Strawberry
During Plug Transplant Production",
Agriculture, 2020

Publication

<1 %

26

jddtonline.info

Internet Source

<1 %

27

www.ajol.info

Internet Source

<1 %

28

Jasril, L. Y. Mooi, A. M. Ali, M. A. Sukari, A. A. Rahman, A. G. Othman, H. Kikuzaki, N. Nakatani. " Antioxidant and Antitumor Promoting Activities of the Flavonoids from ", *Pharmaceutical Biology*, 2008

Publication

<1 %

29

core.ac.uk

Internet Source

<1 %

30

idr.nitk.ac.in

Internet Source

<1 %

31

repository-tnmgrmu.ac.in

Internet Source

<1 %

32

www.vbmv.org

Internet Source

<1 %

33

Girmantė Jurkšienė, Vaida Sirgedaitė-Šėžienė, Aušra Juškauskaitė, Virgilijus Baliuckas.

"Identification of *Alnus incana* (L.) Moenx. × *Alnus glutinosa* (L.) Gaertn. Hybrids Using Metabolic Compounds as Chemotaxonomic Markers", *Forests*, 2023

Publication

<1 %

34

Patrick M. Schlievert. " Chitosan Malate Inhibits Growth and Exotoxin Production of Toxic Shock Syndrome-Inducing Strains and

<1 %

Group A Streptococci ", Antimicrobial Agents and Chemotherapy, 2007

Publication

35

Sarah Heinrich, Adam Toomes, Lalita Gomez. "Valuable stones: The trade in porcupine bezoars", Global Ecology and Conservation, 2020

Publication

<1 %

36

Soumya Ranjan Nayak, Deepak Ranjan Nayak, Utkarsh Sinha, Vaibhav Arora, Ram Bilas Pachori. "An Efficient Deep Learning Method for Detection of COVID-19 Infection Using Chest X-ray Images", Diagnostics, 2022

Publication

<1 %

37

Viney Chawla, Shubhini A. Saraf. "Glyceryl Behenate and Its Suitability for Production of Aceclofenac Solid Lipid Nanoparticles", Journal of the American Oil Chemists' Society, 2010

Publication

<1 %

38

fppn.biomedcentral.com

Internet Source

<1 %

39

papiro.unizar.es

Internet Source

<1 %

40

topsecretapiaccess.dovepress.com

Internet Source

<1 %

41

www.dovepress.com

Internet Source

<1 %

42

Dong Zhang, Zhi Ma, Hanchi Chen, Weilin Ma, Jiawei Zhou, Qing Wang, Chan Min, Yuele Lu, Xiaolong Chen. "Efficient production of valinomycin by the soil bacterium, *Streptomyces* sp. ZJUT-IFE-354", 3 Biotech, 2021

Publication

<1 %

43

Sławomir Kocira, Agnieszka Szparaga, Anna Kocira, Ewa Czerwińska et al. "Modeling Biometric Traits, Yield and Nutritional and Antioxidant Properties of Seeds of Three Soybean Cultivars Through the Application of Biostimulant Containing Seaweed and Amino Acids", *Frontiers in Plant Science*, 2018

Publication

<1 %

44

Chien Huey Teh, Wasi Ahmad Nazni, Ab Hamid Nurulhusna, Ahmad Norazah, Han Lim Lee. "Determination of antibacterial activity and minimum inhibitory concentration of larval extract of fly via resazurin-based turbidometric assay", *BMC Microbiology*, 2017

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off