

Layanan Perpustakaan UHAMKA

Nurul Azmah N-Detection and quantification of pork and rat DNA in processed meats using multiplex quantitative Real-Tim...

 24/07/2024

 Fakultas Farmasi Dan Sains

 Universitas Muhammadiyah Prof. Dr. Hamka

Document Details

Submission ID

trn:oid::1:2995635009

Submission Date

Sep 2, 2024, 8:10 AM GMT+7

Download Date

Sep 2, 2024, 8:14 AM GMT+7

File Name

94212-344757-1-ED_clear_2_-_Nurul_Azmah_Nikmatullah.docx

File Size

329.6 KB

19 Pages

4,607 Words

24,264 Characters





19% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.




Filtered from the Report

- Bibliography

Match Groups

-  **11 Not Cited or Quoted 3%**
Matches with neither in-text citation nor quotation marks
-  **4 Missing Quotations 2%**
Matches that are still very similar to source material
-  **29 Missing Citation 11%**
Matches that have quotation marks, but no in-text citation
-  **11 Cited and Quoted 3%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 13%  Internet sources
- 15%  Publications
- 6%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 11 Not Cited or Quoted 3%**
Matches with neither in-text citation nor quotation marks
- 4 Missing Quotations 2%**
Matches that are still very similar to source material
- 29 Missing Citation 11%**
Matches that have quotation marks, but no in-text citation
- 11 Cited and Quoted 3%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 13% Internet sources
- 15% Publications
- 6% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	www.ncbi.nlm.nih.gov	1%
2	Internet	www.tandfonline.com	1%
3	Internet	jurnal.ugm.ac.id	1%
4	Internet	www.mdpi.com	1%
5	Student papers	Birkbeck College	1%
6	Internet	pdffox.com	1%
7	Publication	Fidel Toldrá, Leo M.L. Nollet. "Handbook of Seafood and Seafood Products Analyti...	1%
8	Internet	link.springer.com	1%
9	Internet	www.bu.edu.eg	1%
10	Publication	E P Sari, L R Kartikasari, M Cahyadi. "Detection of chicken contamination in beef ...	1%

11	Internet	sigaa.ufpa.br	1%
12	Publication	B Wards. "Detection of Mycobacterium bovis in tissues by polymerase chain react...	1%
13	Student papers	Universiti Malaysia Sabah	0%
14	Publication	Shafa'atu Murtala, Abel Simon Agbaji, Peter Maitalata Waziri, Samson Baranzan ...	0%
15	Publication	G. Schares, A. Maksimov, W. Basso, G. Moré et al. "Quantitative real time polymer...	0%
16	Student papers	Mahidol University	0%
17	Publication	Ariel Brunn, Zaharat Kadri-Alabi, Arshnee Moodley, Luca Guardabassi, Phil Taylor,...	0%
18	Publication	Eiseul Kim, Seung-Min Yang, Ik-Seon Kim, Hae-Yeong Kim. "Identification of novel...	0%
19	Publication	Hans-Ulrich Waiblinger, Carina Geppert, Daniela Bartsch, Katrin Neumann et al. "...	0%
20	Publication	Siska Siska, Hanifah Rahmi, Almawati Situmorang. "The Effectiveness of Technical...	0%
21	Internet	e-journal.uajy.ac.id	0%
22	Internet	repository.cshl.edu	0%
23	Internet	www.afia.co.id	0%
24	Publication	Artnice Mega Fathima, Laila Rahmawati, Anjar Windarsih, Suratno Suratno. "Adva...	0%

25	Publication	Jennie Pryce. "Advances in breeding of dairy cattle", Burleigh Dodds Science Publi...	0%
26	Publication	Leo M.L. Nollet, Fidel Toldra. "Handbook of Seafood and Seafood Products Analysi...	0%
27	Internet	doczz.net	0%
28	Internet	journal.uinsgd.ac.id	0%
29	Publication	Abdul Rohman, Yuny Erwanto, M.A. Motalib Hossain, Myrto Rizou, Turki M.S. Alda...	0%
30	Publication	Alexander Arie Sanata Dharma. "Exploring the Communication, Information, and ...	0%
31	Student papers	Manchester Metropolitan University	0%
32	Publication	TANABE, Soichi, Eiji MIYAUCHI, Akemi MUNESHIGE, Kazuhiro MIO, Chikara SATO, ...	0%
33	Internet	core.ac.uk	0%
34	Internet	docsdrive.com	0%
35	Internet	nexusacademicpublishers.com	0%
36	Internet	www.wjgnet.com	0%
37	Publication	Dongyou Liu. "Handbook of Nucleic Acid Purification", CRC Press, 2019	0%
38	Publication	Yicun Cai, Yuping He, Rong Lv, Hongchao Chen, Qiang Wang, Liangwen Pan. "Det...	0%

39	Publication	Al-Kahtani, Hassan A., Elsayed A. Ismail, and Mohammed Asif Ahmed. "Pork detec...	0%
40	Publication	D P Sinambela, D Bakti, M C Tobing, Y M Kusumah. "Molecular characterization of...	0%
41	Publication	Dongyou Liu. "Molecular Detection of Human Fungal Pathogens", CRC Press, 2019	0%
42	Publication	Jozef Golian, Zuzana Drdlová, Lucia Benešová. "Chapter 10 Objectification of Reli...	0%
43	Publication	Putri Widyanti Harlina, Vevi Maritha, Raheel Shahzad, Mohamad Rafi et al. "Comp...	0%
44	Publication	Sirazum Munira, Fatema Tuz Jahura, Md Munir Hossain, Mohammad Shamsul Ala...	0%

Detection and quantification of pork and rat DNA in processed meats using multiplex quantitative Real-Time PCR (m-qPCR)

Abstract

In addition to the issue of pork contamination, processed meats frequently contain traces of rat meat. The study aimed to detect and quantify the pork and rat DNA in cases of meat and processed meat adulteration. In the current study, two gene targets of the cytochrome *b* for pigs and the Mt-*atp6* of *Rattus norvegicus* for rats were used in the absolute multiplex quantitative real-time PCR (m-qPCR). The sample DNA was amplified with a standard as positive control in the various concentration of 1000 pg, 100 pg, 10 pg, 0.1 pg, 0.01 pg, and 0.001 pg. There were 25 processed meat samples and 5 fresh meat samples identified in this study. Among the total of 30 samples assessed, 6 samples were successfully detected and quantified their pork and rat DNA contamination. One sample was contaminated with pork DNA with a concentration of 2.451×10^{-4} pg ("Meatball 3). Five samples were contaminated with rat DNA with a concentration of 3.603×10^{-11} pg ("Sempol 3"), 2.196×10^{-10} pg ("Meatball 6"), 4.908×10^{-11} pg ("Siomay 3"), 1.489×10^{-10} pg ("Grinding 2"), and 3.564×10^{-10} pg ("Grinding 4"). The study discovered contamination of pork and rat in the samples.

Keywords: Multiplex PCR, Cytochrome b, Mt – *atp6*, Processed meat, qPCR.

1. Introduction

Halal food is essential for the Indonesian Muslim community, as mentioned in the Quran and Hadist. As the global halal food market is growing, however, there are increased risks of fraud and

adulteration. It has been reported that frequent incidents of meat fraud and adulteration occurred in ASEAN countries over 20 years (2000-2020) (Owolabi and Olayinka 2021). Indonesia itself also face high case of meat adulterations. They were mostly formalin meat, "glonggong" meat, rotten meat, and fake meat or species substitution (Anuar Ramli et al. 2018). In the case of species substitution, Indonesia has also faced numerous cases of meat and processed meat being mixed with pork (Sari 2017; Maulani et al. 2020; Nida et al. 2020; Mustaqimah et al. 2021; Siswara et al. 2022; Waluyo et al. 2023). Recently, many cases of adulterated meat with rats have also been discovered and the numbers are increased (Suryawan et al. 2020; Lestari et al. 2022).

A halal product regulation of Law Number 33 of 2014 concerning Halal Product Certification, amended by Law Number 11 of 2020 concerning Job Creation, is being implemented in Indonesia. The law demands that all products must undergo halal standard and certification. They have to be labelled as halal; thus, it has to be made using halal ingredients. These regulations have implications for legal consequences if they are not complied. Therefore, the regulation should strongly protect meat ingredients and their derivatives from adulteration. Despite the presence of this regulation, the practice of mixing a specific type of meat with meat from a different species often occurs to increase profits in certain industries. Meat adulteration is considered a crime as it introduces unsafe and low-quality products into the market. Such cases happen when meat is mixed with cheaper meats with similar characteristics, such as pork, rat, and a combination of pork and rat. Therefore, meat detection methods are vital to detect such contamination. The meat detection can provide halal authenticity as well as to prevent and decrease meat and processed meat adulteration cases in the market.

The most accurate method to detect meat adulteration is based on the DNA marker using polymerase chain reaction (PCR) (Tanabe, Miyauchi, et al. 2007; Sari 2017; Maulani et al. 2020;

Waluyo et al. 2023; Mustaqimah Septiani, and Roswiem 2021). PCR was selected because this technique has high sensitivity and can amplify the specific targets, even those of highly complex genomic sequences (Tanabe, Hase, et al. 2007). Many studies have used PCR to detect pork contamination in meat and processed meats. The specific gene target for pork detection is the porcine cytochrome b region of mitochondrial DNA (Tanabe, Miyauchi, et al. 2007; Tanabe, Hase, et al. 2007). Other gene targets commonly used are the genes of ND2, ND5, and 12S rRNA (Chisholm et al. 2005; Kesmen et al. 2009; Cahyadi et al. 2020). However, with the rising number of rat contamination, there has also been an increase in studies focusing on rat detection. The genetic marker for rat detection is the Mt-atp6 of *Rattus norvegicus* gene (Sihotang et al. 2023). Other gene targets for rat detection in meat products are ND5, cyt b 42, and mt-CoI genes (IkaWidyasa et al. 2015; Sihotang et al. 2021; Masnaini et al. 2023). Two protein markers of *Rattus norvegicus* have also been reported to be used for rat detection in meat produced from non-halal slaughter (Aini, Airin, and Raharjo, 2022). As the complexity of meat adulteration cases are increasing lately, thus a particular detection is needed to overcome these complex adulteration cases. Multiplex quantitative Real-time PCR (m-qPCR) is an evolution of PCR detection method, which not only can detect but also quantify the contamination. However, research on the detection and quantification of pork and rats simultaneously in meat and processed meat are yet limited.

Simultaneous detection of multiple species contamination in meat and processed meat can be conducted using multiplex PCR. The multiplex PCR uses several primers simultaneously in one reaction to amplify multiple target sequences (Indriati et al. 2019). It has been reported that multiplex PCR assay can be used to detect species substitutions of goat, cattle, chicken, and pig (Cahyadi et al. 2021). Multiplex PCR has been reported to discriminate the presence of beef and pork in meat samples using the cyt b gene as the marker (Indriati et al. 2019). It is described that

the primers of the cyt b gene can produce different lengths of DNA fragments based on the specific length of each species; therefore, it is utilized to discriminate two species simultaneously (Indriati et al. 2019). Therefore, this study aims to identify pork and rat contamination in meat and processed meat using two pairs of primers derived from porcine cyt b sequences and Mt-atp6 of *Rattus novergicus* gene sequences using multiplex PCR. Furthermore, the quantifications of pork and rat contaminations in the samples were performed in this study.

2. Materials and Methods

2.1 Sample collection

The samples in this study were taken from the East Jakarta region, as previous research had found rat DNA contamination in sausage samples from the street vendors in the East Jakarta area. The samples identified were processed meat obtained randomly from night market traders in East Jakarta and raw meat from meat grinding locations in East Jakarta. The samples consisted of 25 types of processed meat, including 7 sempols, 6 siomay, 7 meatballs, and 5 dimsum. Meanwhile, 5 samples of raw meat were collected from different grinding locations in East Jakarta. The identified samples were uncertified halal processed meats.

2.2 DNA extraction

DNA extraction began with sample preparation. Each sample was weighed at 25 mg. The kit used in this method is the gSYNCTM DNA Extraction kit (Geneaid, Taiwan). It includes GST buffer, proteinase K, GBS buffer, GD columns, W1 buffer, wash buffer, and elution buffer. The weighed sample was put into a 1.5 mL microtube, added with 200 μ L of GST buffer and 20 μ L of proteinase K, vortexed for 15 s, and incubated overnight at 60 °C. After incubation, the samples were centrifuged for 2 min at 12,000 rpm. An amount of 200 μ L of the supernatant was pipetted and

then put into a 1.5 mL microtube. 200 μ L of GSB buffer was added to the tube and then vortexed for 10 s. An amount of 200 μ L of absolute ethanol was put to the tube and then vortexed for 10 s. The sample solution was pipetted and then transferred to the GD column. The tube was centrifuged for 1 min at 12,000 rpm. The GS column was filled with 400 μ L of W1 buffer, and it was centrifuged for 30 s at 12,000 rpm. The supernatant was discarded, and the remaining part was filled with 600 μ L of wash buffer, and it was then centrifuged for 30 s at 12,000 rpm. The supernatant was discarded. After the centrifugation of GS column for 3 min at 12,000 rpm, the GS column was transferred into a new 1.5 mL microtube. The GS column was filled with 50 μ L of elution buffer and incubated at room temperature for 3 min. At 12,000 rpm, the GS Column tube was centrifuged for 30 s. The DNA was characterized using a spectrophotometer (Thermo ScientificTM NanoDrop One). DNA amplification was conducted using the qPCR method.

2.3 Multiplex-PCR

The results of DNA extraction were amplified using the Toyobo THUNDERBRID qPCR probe targeting the cytochrome b gene for pigs (Tanabe, Hase, et al. 2007) and the Mt-atp6 gene *Rattus norvegicus* for rats (Sihotang et al. 2023). The 20 μ L total PCR reaction included 2 μ L of DNA template, 0.6 μ L of primer, 0.4 μ L of probe, and 10 μ L of Toyobo THUNDERBRID qPCR probe. Up until 20 μ L of nuclease-free water (NFW) was supplied, the reaction volume remained maintained. Using a CFX96 Touch Deep Well Real-Time PCR, DNA was amplified for 45 cycles: denaturation at 95 °C for 15 s, annealing at 57 °C for 30 s, and extension at 60 °C for 30 s. The initial denaturation temperature was set for 1 minute. For the cytochrome b gene (pigs) and the Mt-atp6 gene (rats), there were two probes used in this amplification: one tagged with fluorescent HEX (rats) and the other with fluorescent FAM (pigs). The segment sequences of pig and rat primer

were used as a standard, as shown in Table 1. There were seven standards at concentrations of 1000 pg, 100 pg, 10 pg, 1 pg, 0.1 pg, 0.01 pg, and 0.001 pg.

3. Results

The result of the concentration of the DNA sample was in the range of 7.7 – 148.2 ng/ μ L, as shown in Table 2. The analysis showed that the lowest concentration was found in the meatball sample at 7.7 ng/ μ L, while the highest was found in the meat sample at 148.2 ng/ μ L. The DNA purity of the samples ranged between 1.82 – 2.06.

In this study, 6 of the total 30 samples assessed were contaminated with pork and rat DNA. The DNA samples were amplified using the method of absolute multiplex quantification real-time PCR (m-qPCR). The amplification results were presented as a standard curve for the cytochrome b gene (pigs) in Figure 1, a standard curve for the Mt-ATP6 gene (rats) in Figure 2, and the standard concentrations in Table 3. The amplification results of DNA samples are presented in Figure 3 and Table 4. The samples in this study include seven sempol, seven meatballs, six siomay, five dimsums, and meat from several grinding locations. Of the 30 samples, one sample was positive for pork ("Meatball 3") and five were positive for rats ("Sempol 3, Siomay 3, Meatball 6, Grinding 2, and Grinding 4").

Table 1 Sequences of primer and probes

Table 2 Concentration and Purity of DNA Samples

Figure 1 Standard Curve of Cytochrome B Gene (Pig)

Figure 2 Standard Curve of *Mt-atp6* Gene (Rat)

Table 3 C_q Value and Standard Concentration

Figure 3 Result of DNA Sample Amplification

Table 4 Cq Value and DNA Sample Concentration

4. Discussion

Cases of processed meat contaminated with other types of meat have occurred in Indonesia; therefore, appropriate methods are required to identify meat contamination. DNA-based methods, i.e., conventional PCR, real-time PCR, and qPCR, are often used to identify meat contaminations (pork and rats) (Chisholm et al. 2005; Kesmen et al. 2009; IkaWidyasa et al. 2015; Indriati et al. 2019; Salamah et al. 2019; Cahyadi et al. 2020; Cahyadi et al. 2021; Sunaryo et al. 2022). Addition to that, a method for detecting substitutions of several species simultaneously is also necessary (Indriati et al. 2019; Cahyadi et al. 2020; Cahyadi et al. 2021). Pork and rat were chosen in this study as they are common contamination found in the meat adulteration cases in Indonesia. Here in this study, we used a multiplex quantitative Real-Time PCR (m-qPCR) as a simultaneous detection method. It can detect multiple species at one time as current adulteration cases in Indonesia have a high probability of multiple species substitution. This method enhances efficiency by shortening the detection time of several species at once. This method has also not been widely explored on the meat substitution cases in Indonesia.

In the current investigation, our absolute multiplex quantitative real-time PCR (m-qPCR) has successfully proven to identify and quantify species substitutions in the meat and processed meat products simultaneously. The method used consist of two main steps, which are DNA extraction and DNA amplification stages. A spin column-based extraction kit was performed to extract the genomic DNA of the samples. The concentration produced using this method was 7.7-148.2 ng/ μ L. The DNA purity was 1.82-2.06, which meets the specification of 1.7-2.0 (Adriany et al. 2020; Sunaryo et al. 2023). In this study, the spin column method was used as it produces the purer DNA

than other methods (Andalia et al. 2023). It is known that the DNA yields from DNA extraction kit is usually lower and the purity is higher (Liao et al. 2017). The results in this study are also in line with that finding that the DNA concentration obtained in this study is also considered lower, however the purity is considered higher. The higher purity provided high-sensitivity detection methods.

Further, the DNA amplification method is the absolute quantification method (qPCR). In this method, the sample DNA is amplified with a positive control in the form of a standard. The standard consists of a combination of pork and rat DNA segment sequences with various concentrations, as shown in Table 1. Those DNA segment sequences targeted the mitochondrial DNA, in which mitochondrial DNA is commonly used to identify the species (Liao et al. 2017). The amplification resulted in two standard curves: the pig standard (FAM) in Figure 1 and the rat standard (HEX) in Figure 2. The standard curve provides information regarding reaction performance with various parameters, namely efficiency (E), R^2 , and slope.

The pig standard curve (FAM) shows the values of E at 87.5%, R^2 at 0.996, and slope at -3.662, while the rat standard curve (HEX) shows the values of E at 88.5%, R^2 at 0.991, and slope at -3.631. The 87.5% and 88.5% efficiency values indicate the relatively efficient qPCR amplification reactions. For the multiplex PCR, it is known that the amplification efficiency should be in the range of 90-110% (quantitative) and 80-120% (qualitative) (Broeders et al. 2014). Those broader range allow the reproducibility of amplification (Broeders et al. 2014). Even though the values are slightly below the ideal criteria of 90% – 110%, the values of 87.5% and 88.5% are still considered reasonable and can provide reliable results. The linearity of R^2 value ≥ 0.98 represents the ideal linearity for multiplex PCR (Broeders et al. 2014). The R^2 values of 0.996 and 0.991 obtained in our study indicated that the amplification data is strongly correlated with the linear

27 model on the standard curve. It described that the relationship between the logarithm of the initial amount of target DNA and the fluorescence uptake is linear. The slope values of -3.662 and -3.631 from our study suggested a successful amplification and indicate a relatively good efficiency level (Luque-Perez et al. 2013; Tan et al. 2020; Mariyani et al. 2021).

37 Of the 30 samples identified, one contained pig DNA, and five were positive for rat contamination. The sample that contained pig DNA was "Meatball 3" with a concentration of 2.451×10^{-4} pg and a Cq value of 22.69. Irwandi et al., 2020 reported the presence of pig DNA contaminations in meatballs, where two out of three samples tested positive for pig DNA. Similarly, 42 Purwantoro et al. 2022 detected pig DNA contamination in sausage samples, with one out of five 30 samples showing the presence of pig DNA. Our result is in line with that of Cahyadi et al. 2020 revealing how multiplex PCR used to detect multiple species contamination in one reaction.

Samples contaminated with rats were "Sempol 3" with a concentration of 3.603×10^{-11} pg and a Cq value of 38.37, "Meatball 6" with a concentration of 2.196×10^{-10} pg and a Cq value of 36.47, "Siomay 3" with a concentration of 4.908×10^{-11} pg and a Cq value of 38.04, "Grinding 2" with a concentration of 1.489×10^{-10} pg and a Cq value of 36.88, and "Grinding 4" with a concentration of 3.564×10^{-10} pg and a Cq value of 35.96. Sunaryo et al., 2022 also reported rat contamination in processed products, with one out of 30 sausage samples contaminated with rat DNA. Meanwhile, Susilowati 2019 discovered cases of non-halal meat contamination in grinding locations, with five out of 30 meat samples found to be contaminated with pork.

5. Conclusion

In this study, the multiplex quantitative Real-Time PCR (m-qPCR) brings a significant result in successfully detect and quantify the contaminated samples. Of the 30 samples, one contained

pork DNA, and five were positive for rat contamination. The sample contaminated with pig DNA was "Meatball 3" with a concentration of 2.451×10^{-4} pg and a Cq value of 22.69. Meanwhile, the samples contaminated with rats were "Sempol 3" with a concentration of 3.603×10^{-11} pg and a Cq value of 38.37, "Meatball 6" with a concentration of 2.196×10^{-10} pg and a Cq value of 36.47, "Siomay 3" with a concentration of 4.908×10^{-11} pg and a Cq value of 38.04, "Grinding 2" with a concentration of 1.489×10^{-10} pg and a Cq value 36.88, "Grinding 4" with a concentration of 3.564×10^{-10} pg and a Cq value of 35.96. The method serves as an effective technique for analyzing and detecting multiple species substitution at once. Furthermore, the multiplex quantitative Real-Time PCR (m-qPCR) can be widely used to assess, trace, and calculate the contamination for ensuring food quality and detecting complex food adulteration.

References

1. Adriany DT, Bakri AA, Bungalim MI. 2020. Comparison of DNA isolation methods on DNA purity for detection of White Spot Syndrome Virus (WSSV) on Bamboo Lobster (*Panulirus versicolor*). In: Prosiding Simposium Nasional VII Kelautan dan Perikanan 2020 Fakultas Ilmu Kelautan dan Perikanan. Makassar; p. 239–246.
2. Aini AN, Airin CM, Raharjo TJ. 2022. Protein markers related to non-halal slaughtering process of rat as mammal animal's model detected using mass spectrometry proteome analysis. Indonesian Journal of Chemistry. 22(1):867. <https://doi.org/10.22146/ijc.73656>.
3. Andalia N, Adriani, Wardani AF, Sahli IT, Yunus R, Solfaine R, Nikmatullah NA, Meri, Rusdin A, Safitri NM. 2023. Molecular Biology. Padang: PT Global Eksekutif Teknologi.

4. Anuar Ramli M, Salahudin A, Imran Abdul Razak M, Ammar Harith Idris M, Izzul Syahmi Zulkepli M. 2018. Halal meat fraud and safety issues in Malaysian and Indonesian market. [place unknown].
5. Broeders S, Huber I, Grohmann L, Berben G, Taverniers I, Mazzara M, Roosens N, Morisset D. 2014. Guidelines for validation of qualitative real-time PCR methods. *Trends Food Sci Technol.* 37(2):115–126. <https://doi.org/10.1016/j.tifs.2014.03.008>.
6. Cahyadi M, Fauziah NAD, Suwanto IT, Boonsupthip W. 2021. Detection of species substitution in raw, cooked, and processed meats utilizing multiplex-PCR assay. *Indones J Biotechnol.* 26(3):128–132. <https://doi.org/10.22146/ijbiotech.63472>.
7. Cahyadi M, Wibowo T, Pramono A, Abdurrahman ZH. 2020. A novel multiplex-PCR assay to detect three non-halal meats contained in meatball using mitochondrial 12S rRNA Gene. *Food Sci Anim Resour.* 40(4):628–635. <https://doi.org/10.5851/kosfa.2020.e40>.
8. Chisholm J, Conyers C, Booth C, Lawley W, Hird H. 2005. The detection of horse and donkey using real-time PCR. *Meat Sci.* 70(4):727–732. <https://doi.org/10.1016/j.meatsci.2005.03.009>.
9. IkaWidyasa Y, Sudjadi, Rohman A. 2015. Detection of rat meat adulteration in meat ball formulations employing Real Time PCR. *Asian J Anim Sci.* 9(6):460–465. <https://doi.org/10.3923/ajas.2015.460.465>.
10. Indriati M, Yuniarsih E, Raya J, Km L, Saketi C, Pandeglang Banten K. 2019. Multiplex PCR method of detecting pork to guarantee halal status in meat processed products. *Jurnal Ilmu Produksi dan Teknologi Hasil Peternakan.* 07(3):96–101. <https://doi.org/10.29244/jipthp.7.2.96-101>.

11. Irwandi I, Wardi ES, Dova S. 2020. Detection of pig gene contamination in packaged beef meatball products in Padang using the PCR. *Jurnal Akademi Farmasi Prayoga*. 5(2):1–12.
12. Kesmen Z, Gulluce A, Sahin F, Yetim H. 2009. Identification of meat species by TaqMan-based real-time PCR assay. *Meat Sci*. 82(4):444–449. <https://doi.org/10.1016/j.meatsci.2009.02.019>.
13. Lestari D, Rohman A, Syofyan S, Yuliana ND, Abu Bakar NKB, Hamidi D. 2022. Analysis of beef meatballs with rat meat adulteration using Fourier Transform Infrared (FTIR) spectroscopy in combination with chemometrics. *Int J Food Prop*. 25(1):1446–1457. <https://doi.org/10.1080/10942912.2022.2083637>.
14. Liao J, Liu YF, Yang L, Li FP, Sheppard AM. 2017. Development of a rapid mitochondrial DNA extraction method for species identification in milk and milk products. *J Dairy Sci*. 100(9):7035–7040. <https://doi.org/10.3168/jds.2017-12653>.
15. Luque-Perez E, Mazzara M, Weber TP, Foti N, Grazioli E, Munaro B, Pinski G, Bellocchi G, Van den Eede G, Savini C. 2013. Testing the robustness of validated methods for quantitative detection of GMOs across qPCR instruments. *Food Anal Methods*. 6(2):343–360. <https://doi.org/10.1007/s12161-012-9445-z>.
16. Mariyani M, Sismindari SU, Rumiati R. 2021. Validation of the real-time Polymerase Chain Reaction method for detecting pig (*Sus scrofa domestica*) and boar (*Sus barbatus*) DNA in beef sausages. Jogjakarta: Universitas Gadjah Mada.
17. Masnaini M, Achyar A, Chatri M, Putri DH, Ahda Y, Irdawati. 2023. Primer design and optimization of pcr methods for detecting mixed rat meat in food samples. *Proceedings of the 3rd International Conference on Biology, Science and Education (IcoBioSE 2021)*.:282–289. https://doi.org/10.2991/978-94-6463-166-1_37.

18. Maulani TR, Susilo H, Indriati M, Suhaemi A, Raya J, Km L, Saketi K, Pandeglang Banten K. 2020. Detection of pig DNA contamination with RT-PCR in sosis without halal labels from district Pandeglang. *Gorontalo Agriculture Technology Journal*. 3(2).
19. Mustaqimah DN, Septiani T, Roswiem AP. 2021. Detection of pork DNA in sausage using a Real Time-Polymerase Chain Reaction (RT-PCR). *Indonesian Journal of Halal*. 3(2):106–111.
20. Nida L, Pisestyani H, Basri C. 2020. Case study: adulteration of beef with wild boar meat in Bogor city. *Jurnal Kajian Veteriner*. 8(2):121–130.
<https://doi.org/10.35508/jkv.v8i2.2326>.
21. Owolabi IO, Olayinka JA. 2021. Incidence of fraud and adulterations in ASEAN food/feed exports: A 20-year analysis of RASFF's notifications. *PLoS One*. 16(11):e0259298.
<https://doi.org/10.1371/journal.pone.0259298>.
22. Purwantoro R, Suryandani H, Hudaya DA, Yuniarsih E, Rostianti T, Teknologi F, Universitas Mathla'ul Anwar Banten I. 2022. Detection of pork contamination in beef sausages using the multiplex PCR method in the Pandeglang district. *Teknotika*. 01(2).
23. Salamah N, Erwanto Y, Martono S, Rohman A. 2019. Real-Time PCR-based detection of bovine DNA by specific targeting on cytochrome-B. *Pharmaciana*. 9(2):201.
<https://doi.org/10.12928/pharmaciana.v9i2.14070>.
24. Sari F. 2017. Identification of pig species in food products of animal origin in traditional markers of Riau Province using Polymerase Chain Reaction. *Jurnal Riau Biologia* . 2(1):55–60.

25. Sihotang M, Sophian A, Purba M, Wilasti Y. 2023. Development of rat meat detection using Mt-atp6 *Rattus norvegicus* gene genetic marker. Curr Appl Sci Technol. 23(1). <https://doi.org/10.55003/cast.2022.01.23.006>.
26. Sihotang MAED, Erwinda YE, Suwarni E, Lusianti E. 2021. Primer design and in silico analysis for amplification of the Mt-Co1 gene in goat mice (*Rattus norvegicus*). Eruditio : Indonesia Journal of Food and Drug Safety. 1(2):20–29. <https://doi.org/10.54384/eruditio.v1i2.82>
27. Siswara HN, Erwanto Y, Suryanto E. 2022. Study of meat species adulteration in indonesian commercial beef meatballs related to halal law implementation. Front Sustain Food Syst. 6. <https://doi.org/10.3389/fsufs.2022.882031>.
28. Sunaryo H, Nikmatullah NA, Mufidah S. 2022. Detection of rat contamination in sausage samples with real time PCR. Farmasains.:57–64.
29. Sunaryo H, Wirman AP, Permanasari ED, Nikmatullah NA, Lestari D, Nurjanah D. 2023. Optimization of dna extraction methods in fresh meat (rat and chicken meat) based on incubation time. Indonesian Journal of Halal Research. 5(2):99–108. <https://doi.org/10.15575/ijhar.v5i2.21325>.
30. Suryawan GY, Suardana IW, Wandia IN. 2020. Sensitivity of Polymerase Chain Reaction in the detection of rat meat adulteration of beef meatballs in Indonesia. Vet World. 13(5):905–908. <https://doi.org/10.14202/vetworld.2020.905-908>.
31. Susilowati T. 2019. Detection of pig DNA contaminants in meat grinding samples at market Surya of Surabaya using real-time PCR. Surabaya: Universitas Islam Negeri Sunan Ampel.
32. Tan LL, Ahmed SA, Ng SK, Citartan M, Raabe CA, Rozhdestvensky TS, Tang TH. 2020. Rapid detection of porcine DNA in processed food samples using a streamlined DNA

extraction method combined with the SYBR Green real-time PCR assay. *Food Chem.* 309:125654. <https://doi.org/10.1016/j.foodchem.2019.125654>.

33. Tanabe S, Hase M, Yano T, Sato M, Fujimura T, Akiyama H. 2007. A real-time quantitative PCR detection method for pork, chicken, beef, mutton, and horseflesh in foods. *Biosci Biotechnol Biochem.* 71(12):3131–3135. <https://doi.org/10.1271/bbb.70683>.
34. Tanabe S, Miyauchi E, Muneshige A, Mio K, Sato C, Sato M. 2007. PCR method of detecting pork in foods for verifying allergen labeling and for identifying hidden pork ingredients in processed foods. *Biosci Biotechnol Biochem.* 71(7):1663–1667. <https://doi.org/10.1271/bbb.70075>.
35. Waluyo S, Malau J, Raekiansyah M, Yulian E, Hardiman I. 2023. Detection and quantification of pork contamination in processed meat samples using real-time PCR. *Al-Kauniah: Jurnal Biologi.* 16(1):46–52. <https://doi.org/10.15408/kauniah.v16i1.20203>.

Figure 1.




Figure 2.




Figure 3.




Table 1.

Target		Sequence
Pork (Porcine DNA)	Forward Primer	5'- CTTGCAAATCCTAACAGGCCTG -3'
	Reverse Primer	5'- CGTTTGCATGTAGATAGCGAATAAC -3'
	TaqMan MGB Probe	5'-(FAM)-ACAGCTTTCTCATCAGTTAC-(NFQ)(MGB) -3'
Rat (Mt-atp6 <i>Rattus norvegicus</i> gene)	RnATP6-161 Forward	5'-ACACCAAAGGACGAACCTG -3'
	RnATP6-161 Reverse	5'-AGAATTACGGCTCCTGCTCA -3'
	RnATP6-161 Probe	5'- [HEX]-TTCTAGGGCTTCTTCCCCAT-[QSY] -3'

Table 2.

NO	SAMPLE CODE	CONCENTRATION ng/μL	PURITY 260/280 nm
1	Sempol 1	57.2	1.84

2	Sempol 2	39.9	1.84
3	Sempol 3	11.5	1.98
4	Sempol 4	51.2	2.00
5	Sempol 5	20.0	2.00
6	Sempol 6	13.9	1.91
7	Sempol 7	22.2	2.00
8	Meatball 1	7.8	1.95
9	Meatball 2	98.4	2.01
10	Meatball 3	28.1	2.06
11	Meatball 4	36.5	2.01
12	Meatball 5	7.7	2.06
13	Meatball 6	75.2	2.03
14	Meatball 7	28.7	1.97
15	Siomay 1	59.7	1.82
16	Siomay 2	63.5	1.93
17	Siomay 3	13.9	1.90
18	Siomay 4	17.1	1.86
19	Siomay 5	20.5	1.87
20	Siomay 6	30.9	1.95
21	Dimsum 1	124.2	2.01
22	Dimsum 2	132.6	1.98
23	Dimsum 3	73.9	2.00
24	Dimsum 4	24.5	1.99
25	Dimsum 5	77.7	1.89
26	Grinding 1	110.1	2.06
27	Grinding 2	115.8	2.06
28	Grinding 3	100.7	2.01
29	Grinding 4	148.2	2.05
30	Grinding 5	120.1	1.98

Table 3.

NO	STANDART (Std)	Cq FAM (Babi)	Concentration FAM (Babi)	Cq HEX(Tikus)	Concentration HEX (Tikus)
1	Std-1	4.93	1.000E+03	4.02	1.000E+03
2	Std-2	9.63	1.000E+02	7.72	1.000E+02
3	Std-3	13.09	1.000E+01	12.35	1.000E+01
4	Std-4	15.59	1.000E+00	13.88	1.000E+00
5	Std-5	17.31	1.000E-01	16.82	1.000E-01
6	Std-6	20.81	1.000E-02	19.38	1.000E-02
7	Std-7	22.49	1.000E-03	22.31	1.000E-03

Table 4.

NO	Sample	Cq	Concentration	Cq	Concentration
----	--------	----	---------------	----	---------------

	Code	FAM (Babi)	FAM (Babi)	HEX(Tikus)	HEX (Tikus)
1	Sempol 1	N/A	N/A	N/A	N/A
2	Sempol 2	N/A	N/A	N/A	N/A
3	Sempol 3	N/A	N/A	38.37	3.603E-11
4	Sempol 4	N/A	N/A	N/A	N/A
5	Sempol 5	N/A	N/A	N/A	N/A
6	Sempol 6	N/A	N/A	N/A	N/A
7	Sempol 7	N/A	N/A	N/A	N/A
8	Meatball 1	N/A	N/A	N/A	N/A
9	Meatball 2	N/A	N/A	N/A	N/A
10	Meatball 3	22.69	2.451E-04	N/A	N/A
11	Meatball 4	N/A	N/A	N/A	N/A
12	Meatball 5	N/A	N/A	N/A	N/A
13	Meatball 6	N/A	N/A	36.47	2.196E-10
14	Meatball 7	N/A	N/A	N/A	N/A
15	Siomay 1	N/A	N/A	N/A	N/A
16	Siomay 2	N/A	N/A	N/A	N/A
17	Siomay 3	N/A	N/A	38.04	4.908E-11
18	Siomay 4	N/A	N/A	N/A	N/A
19	Siomay 5	N/A	N/A	N/A	N/A
20	Siomay 6	N/A	N/A	N/A	N/A
21	Dimsum 1	N/A	N/A	N/A	N/A
22	Dimsum 2	N/A	N/A	N/A	N/A
23	Dimsum 3	N/A	N/A	N/A	N/A
24	Dimsum 4	N/A	N/A	N/A	N/A
25	Dimsum 5	N/A	N/A	N/A	N/A
26	Grinding 1	N/A	N/A	N/A	N/A
27	Grinding 2	N/A	N/A	36.88	1.489E-10
28	Grinding 3	N/A	N/A	N/A	N/A
29	Grinding 4	N/A	N/A	35.96	3.564E-10
30	Grinding 5	N/A	N/A	N/A	N/A